

**VASCULAR REACTIVITY, CHEMICAL PROFILE,
TOXICOLOGICAL AND PHARMACOKINETIC
STUDIES OF *ANDROGRAPHIS PANICULATA* NEES.
EXTRACTS**

SRIRAMANENI RAGHAVA NAIDU

UNIVERSITI SAINS MALAYSIA

JULY 2009

**VASCULAR REACTIVITY, CHEMICAL PROFILE,
TOXICOLOGICAL AND PHARMACOKINETIC
STUDIES OF *ANDROGRAPHIS PANICULATA* NEES.
EXTRACTS**

BY

SRIRAMANENI RAGHAVA NAIDU

**Thesis submitted to fulfillment of the requirements for the degree of
Doctor of Philosophy**

July 2009

DEDICATION

**My Wife, Son (Anurag), Parents, Brother, Uncle, and Relatives for their love,
support and encouragement**

and

To Lord Saibaba

ACKNOWLEDGEMENTS

I take this opportunity to express my deep sincere gratitude and heartfelt thanks to my esteemed supervisor, Professor Dr. Mohd. Zaini Asmawi for his inspiration, guidance, constant encouragement, supervision, functional freedom, sound advice and personal care given throughout my research work.

I express my deep thanks to my co-supervisor Associate Prof. Dr. Amirin Sadikun for his invaluable support, advice, encouragement and constant supervision in completing my project successfully.

I am highly thankful to my best friends Venkat, Ram Mohan, Shesala Ravi, Rehmat, Annegowda, Naveen, Anand Swaroop, Omar, Ibrahim, Ali Jimale, Yam, Mahfoudh, Tharindu, colleagues from UCSI and ICT for helping me to get through the difficult times, and for all the emotional support, entertainment and caring they provided.

I am indebted to my lab technician Mr. Roseli Hassan his pivotal role in smooth progress of my project work. I earnestly thankful to pharmacology lab staff Mrs. Wong, Mr. Hassan, Mr. Selva, Mr. Wan and other technicians as well as staff of School of Pharmaceutical Sciences, Universiti Sains Malaysia who were contributed either directly or indirectly throughout the progress of my research work.

It gives me an immense pleasure to express my appreciation to Ministry of Science, Technology, and Environment for the research grant 304/PFARMASI/640043/k105 and 304/PFARMASI/6123003 for providing research chemicals and facilities.

Last in line but first in mind to my beloved wife, my son (Manu), parents, brother, uncle, and relatives for their love, encouragement and support without which this research works would be incomplete.

Raghava Naidu Sriramaneni

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iv
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATION & SYMBOLS	xix
LIST OF APPENDICES	xxiii
LIST OF PUBLICATIONS/PRESENTATIONS	xxiv
ABSTRAK	xxv
ABSTRACT	xxvii
CHAPTER 1 : INTRODUCTION	
1.0 Introduction	01
1.1 Vascular Smooth Muscle	01
1.2 Contraction of Blood Vessels	02
1.3 Smooth Muscle Relaxation	06
1.4 The Endothelium	07
1.4.1 The Physiology of Endothelium	07
1.5 Endothelium and Cardiovascular Disease	08
1.6 Vascular Mediators from the Endothelium: Endothelium-Derived Relaxant Factors (EDRF)	10
1.6.1 Prostacyclin	13
1.6.2 Endothelium-Derived Hyperpolarizing Factor (EDHF)	15
1.7 Hypertension	18
1.8 Some Pharmacologically Active Moieties from Natural Flora Used in Cardiovascular Malfunctions	20
1.9 Conventional Antihypertensive Agents	21
CHAPTER 2 : REVIEW OF LITERATURE	
2.0 <i>Andrographis paniculata</i>	24
2.1 Classification of <i>Andrographis paniculata</i>	26

2.2	Profiles of <i>Andrographis paniculata</i>	26
2.3	Ethnobotanical Review of <i>Andrographis paniculata</i>	27
2.4	Ethnobotanical uses of <i>Andrographis paniculata</i>	28
2.5	Pharmacological Activities of Important Active Constituents of <i>Andrographis paniculata</i>	29
2.6	Herbal Medicine	41
2.6.1	Nature as a Source of Drug Compounds	42
2.6.1 (A)	Alkaloids	42
2.7	Current research on <i>Andrographis paniculata</i> with Various Activities	44
2.8	Flavonoids	53
2.8.1	Vascular Effects of Flavonoids	55
2.8.2	Vasodilator Effects	55
2.8.3	Endothelium-Dependent Relaxation	57
2.9	Mechanism of Action	57
2.9.1	Inhibition of PKC	58
2.9.2	Inhibition of Ca ²⁺ Entry	58
2.9.3	Inhibition of Cyclic Nucleotide Phosphodiesterase	59
2.9.4	Inhibition of Tyrosine Kinases	60
2.9.5	Cardiac Effects	61
2.9.6	Antihypertensive Effects of Flavonoids	61
2.9.7	Coumarin	63
2.9.8	Terpenoids	63

CHAPTER 3: PREPARATION OF EXTRACTS OF ANDROGRAPHIS PANICULATA (NEES.) 66

3.0	Introduction	66
3.1	Objectives	67
3.2	Materials and Methods	67
3.2.1	Plant Material	67
3.2.2	Solvents & Reagents	67
3.2.3	Preparation of the Extract	67
3.3	Results	70
3.3.1	Yield of Extracts	70

3.4	Discussion	70
CHAPTER 4: SCREENING OF PETROLEUM ETHER, CHLOROFORM AND METHANOLIC EXTRACTS OF ANDROGRAPHIS PANICULATA ON RAT THORACIC AORTA		71
4.0	Introduction	71
4.1	Objectives	72
4.2	Materials and Methods	72
4.2.1	Materials	72
4.2.2	Plant Material and Preparation of Extracts	72
4.2.3	Experimental Animals	73
4.2.4	Organ Baths	73
4.2.5	Experimental Studies	73
4.2.5 (A)	Preparation and Setting up of Rat Aorta	73
4.2.5 (B)	Record of Isometric Vascular Tone	74
4.2.5 (C)	Calculation of Responses	75
4.2.6	Statistical Analysis	75
4.3	Results	76
4.3.1	The Effect of Petroleum Ether (PE) Extract of AP on Norepinephrine-Induced Contraction of Isolated Aortic Strip Preparations	76
4.3.2	The Effect of Methanolic Extract of AP on NE Induced Contraction of Isolated Aortic Strip Preparations	76
4.3.3	The Effect of <i>Andrographis Paniculata</i> Chloroform Extract (APCE) on NE Induced Contraction of Isolated Aortic Strip Preparations	79
4.4	Discussion	80

CHAPTER 5: CHEMICAL PROFILE OF <i>ANDROGRAPHIS PANICULATA</i> CHLOROFORM EXTRACT (APCE), ANDROFEAPHOLIDE (ANG) AND 14-DEOXY-11, 12-DIDEHYDROANDROGRAPHOLIDE (DDA).	81
5.0 Introduction	81
5.1 Objectives	84
5.2 Thin-Layer Chromatography (TLC)	84
5.3 Subfraction of AP ₁ from Chloroform Fraction of <i>Andrographis Paniculata</i> using Preparative Thin-Layer Chromatography (PTLC)	85
5.3.1 Materials	85
5.3.2 Method of Separation	85
5.3.3 Removal of The AP ₁ Compound band from the Preparative TLC	86
5.4 Results	86
5.4.1 The Retention Factor for Standard	87
5.4.2 The Retention Factor of Extracts	87
5.5 HPLC and NMR Profiles of Chloroform Extract of <i>Andrographis paniculata</i>	87
5.5.1 Analytical Column	87
5.5.2 Retention Time (R _t)	88
5.5.3 Peak Capacity	89
5.6 HPLC Profile of ANG, 14-Deoxyandrographolide (DA) and 14-Deoxy-11, 12-Didehydroandrographolide (DDA) and APCE.	90
5.7 Method	91
5.7.1 Preparation of Standard Solution	91
5.7.2 Preparation of Sample Solution	91
5.7.3 Procedure	92
5.8 Results and Discussion	92
5.9 NMR Profiles of Diterpenoid Lactones from <i>Andrographis Paniculata</i>	97
5.9.1 Methods	100
5.9.2 Results and Discussion	101

CHAPTER 6 :	CHRONIC EFFECTS OF DITERPENOID LACTONES FROM <i>ANDROGRAPHIS PANICULATA</i> CHLOROFORM EXTRACT IN SPONTANEOUSLY HYPERTENSIVE RATS	106
6.0	Introduction	106
6.1	Objectives	107
6.2	Materials and Methods	107
6.2.1	Materials	107
6.2.2	Experimental Animals	107
6.2.3	Effect of Ach and sodium nitroprusside (SNP) on vascular function	108
6.2.4	Plant material and preparation of extracts	108
6.2.5	Chronic Effects of APCE in SHR Rats	109
6.2.6	Effects in Hypertensive Rats	109
6.3	Parameters Studied	109
6.3.1	Noninvasive Measurement of Systolic Blood Pressure	109
6.3.2	Measurement of Rat Aortic Contraction	110
6.4	Data Presentation and Statistical Analysis	110
6.5	Results	111
6.5.1	Effect of oral Administration of APCE Daily For 4 Weeks on Systolic Blood Pressure of SHR Rats	111
6.5.2	Effect of 4 Weeks Daily oral Treatment with APCE on Acetylcholine Induced Relaxation of Aorta Free Contracted with Phenylephrine	112
6.5.3	Endothelium Denuded Aortic Relaxation to Sodium Nitroprusside (SNP)	114
6.5.4	Contractions to High K ⁺ and Phenylephrine (PE)	116
6.6	Discussion	116
CHAPTER7:	VASORELAXANT EFFECT INDUCED BY DITERPINOID LACTONES FROM <i>ANDROGRAPHIS PANICULATA</i> IN RAT AORTIC RINGS	119
7.0	Introduction	119
7.1	Objectives	121
7.2	Materials And Methods	121

7.2.1	Materials	121
7.2.2	Tissue Preparations	122
7.2.3	Tension Recording	123
7.3	Endothelium Dependent and Independent Vasorelaxation	123
7.4	Statistical Analysis	125
7.5	Results	125
7.5.1	Effects of DA on Ca ²⁺ -Induced Contraction in the Presence of High K ⁺ .	125
7.5.2	Effect of 14-Deoxyadnrographolide (DA) on the Dose Responce Curve of Endothelium Intact Arota to KCl and NE	129
7.5.3	Effects of 14-Deoxy-11, 12-Didehydroadnrographolide (DDA) on the Contractive Concentration– Response Curve of Endothelium Intact Arota to KCl and NE	131
7.5.4	Role of Endothelium in APCE-Induced Relaxation	134
7.5.5	Effects of APCE, DA, and DDA on the Phasic and Tonic Contraction Induced by NE	136
7.5.6	Influence of Different Factors on The Relaxant Effect of APCE, DA and DDA	140
7.6	Discussion	141
CHAPTER8: TOXICOLOGICAL SCREENING OF CHLOROFORM EXTRACT OF ANDROGRAPHIS PANICULATA IN EXPERIMENTAL ANIMALS		144
8.0	Introduction	144
8.1	Objectives	145
8.2	Materials and Methods	146
8.2.1	Preparation of Plant Extracts	146
8.3	Animals	146
8.3.1	Acute Toxicity (Determination of LD ₅₀)	146
8.3.2	Sub-acute Toxicity	147
8.3.3	Observations and Examination Methods	147
8.3.4	Blood Analysis	147
8.3.5	Biochemical Analysis	148

8.3.6	Tissue Analysis	148
8.3.7	Histopathological Examination	148
8.4	Statistical Analysis	149
8.5	Results	149
8.5.1	Sub-acute Toxicity	149
8.5.1 (A)	General Signs	149
8.5.1 (B)	Hematological Data Analysis	149
8.5.1 (C)	Serum Data Analysis	151
8.5.1 (D)	Tissue Organ Analysis	153
8.5.1 (E)	Body Weight and Systolic Blood Pressure (SBP)	154
8.5.1 (F)	Histopathological Examination	155
8.6	Discussion	167
CHAPTER 9: PHARMACOKINETIC STUDY OF CHLOROFORM EXTRACT OF <i>ANDROGRAPHIS PANICULATA</i>		171
9.0	Introduction	171
9.1	Objectives	173
9.2	Materials and Methods	174
9.2.1	Materials	174
9.2.2	Plant Extracts Preparation	174
9.2.3	Instruments	174
9.2.4	Chromatographic Condition And Sample Preparation	174
9.3	Study Protocol	175
9.3.1	Experimental Animals	175
9.3.2	Dosing of AP Chloroform Extract (APCE)	175
9.3.2 (A)	Sampling of Blood	175
9.4	Validation of HPLC Method	176
9.4.1	Linearity	176
9.4.2	Precision And Accuracy	176
9.4.3	Calculation	176
9.4.3 (A)	Quantification	176
9.4.3 (B)	Amount of Markers in Plasma	177

9.5	Calculation of other Kinetic Parameters	177
9.6	Results	178
9.6.1	Method Validation	178
9.7	Pharmacokinetic Application	183
9.8	Discussion	185
9.9	Conclusion	186
CHAPTER 10: SUMMARY		187
CHAPTER 11: FUTURE SCOPE OF WORK		191
REFERENCES		193
APPENDICES		226
PUBLICATIONS		

LIST OF TABLES

	Page
5.1 Summary of HPLC methods of ANG and other diterpenoids	83
5.2 ¹ H-NMR chemical shift corresponding to ANG in the absence and presence of APCE.	101
5.3 ¹ H-NMR chemical shift corresponding to DDA in the absence and presence of APCE.	102
6.1 Systolic Blood Pressure (SBP) of SHR after being treated for four weeks with APCE groups as compared to control.	111
8.1 Effect of oral administration of chloroform extract of <i>Andrographis paniculata</i> daily for 28 days on hematological profile of rats.	150
8.2 The effect of APCE on liver function enzyme tests on rat serum	151
8.3 The effect of APCE on heart and kidney indices on rat serum	152
8.4 Effect of APCE on organ weight profile in sub-acute toxicity.	153
8.5 Effect of oral administration of APCE daily for 28 days on systolic blood pressure (SBP) of rats.	154
9.1 Regression equations, correlation coefficient and linearity ranges of ANG and DDA in rat plasma	178
9.2 Accuracy and Precision (n=6) of the ANG and DDA in rat plasma.	182
9.3 Recovery (n=3) of the ANG and DDA in rat plasma.	182
9.4 Stability results (n = 3).of the ANG and DDA in rat plasma.	183
9.5 Concentration profile of ANG and DDA in rat serum samples (n=6).	184
9.6 Pharmacokinetic parameters following oral administration of APCE extract single dose (1000 mg/kg).	185

LIST OF FIGURES

	Page
1.1 Structure of artery, vein (Essentials of anatomy and physiology Martini & Bartholomew, 2008)	02
1.2 Receptor-mediated contraction and relaxation in different types of smooth muscle (Mineman and Wecker, 2004)	04
1.3 Mechanisms of contraction of vascular smooth muscle cells (Mineman and Wecker, 2004)	06
1.4 Biosynthesis of L-Arginine-nitricoxide (NO) pathway	12
1.5 Biosynthesis of prostaglandins, lipoxygenase and cytochrome P-450 pathway	15
2.1 Leaves and aerial parts of <i>Andrographis paniculata</i> .	25
3.1 Schematic flow chart diagram for extraction of dried powdered aerial parts of <i>Andrographis paniculata</i> (Nees.) with petroleum ether, chloroform and methanol	69
4.1 Contractile responses to norepinephrine (NE) of aortic rings in the presence of petroleum ether extract 0.25 mg/ml, 0.5 mg/ml and 1.0 mg/ml (n=8) ** p<0.01 and *** p<0.001	77
4.2 Contractile responses to norepinephrine (NE) of aortic rings in the presence of methanol extract 0.25 mg/ml, 0.5 mg/ml, 1.0 mg/ml and 2.0 mg/ml (n=8) * p<0.05, ** p<0.01 and *** p< 0.001	78
4.3 Contractile responses to norepinephrine (NE) of aortic rings in the presence of <i>Andrographis paniculata</i> chloroform extract (APCE) 20 μ g/ml, 40 μ g/ml, 80 μ g/ml and 160 μ g/ml (n=8) *** p<0.001	79
5.1 HPLC chromatogram of standard Andrographolide	94
5.2 HPLC chromatogram of standard DA	94
5.3 HPLC chromatogram of standard DDA	95
5.4 HPLC chromatogram of <i>Andrographis paniculata</i> chloroform extracts (APCE)	95
5.5 HPLC chromatogram of sub fraction of AP ₁	96
5.6 Chemical structure of andrographolide (ANG)	99
5.7 Chemical structure of 14-deoxy-11, 12-didehydroandrographolide (DDA)	100

5.8	¹ H-NMR Spectral analysis of Andrographolide (ANG)	103
5.9	¹ H-NMR Spectral analysis of 14-deoxy-11,12-didehydro andrographolide (DDA)	104
5.10. (A)	¹ H-NMR Spectral analysis of <i>Andrographis paniculata</i> chloroform extract (APCE)	105
5.10 (B)	¹ H-NMR Spectral analysis of <i>Andrographis paniculata</i> chloroform extract (APCE)	105
6.1	The effect of chloroform extract of AP (APCE) on acetylcholine-induced relaxation of epithelium intact SHR aorta pre-contraction with 1 μ M phenylephrine. Symbols represent mean \pm SEM of 8 experiments *** p<0.001, treatment versus control group	113
6.2	The effect of APCE on sodium nitroprusside-induced relaxation of endothelium denuded SHR aorta pre-contracted with 1 μ M phenylephrine. Symbols represent mean \pm SEM of 8 experiments *** P < 0.001, treatment versus control group	115
7.1	Effect of DA 10 μ M, 20 μ M and 40 μ M on dose response curve of Ca ²⁺ endothelium intact aorta. The aortic rings were pre-incubate d with 0.1% DMSO *** p< 0.001 vs control	126
7.2	Effect of DDA on the Ca ²⁺ dependent contraction aortic rings. The aortic rings were pre-incubated with 0.1% DMSO, DDA (10 μ M, 20 μ M and 40 μ M). *** P < 0.001 vs control	127
7.3	Effect of <i>Andrographis paniculata</i> chloroform extract (APCE) 10 μ g/ml, 20 μ g/ml, 40 μ g/ml and 80 μ g/ml on Ca ²⁺ dependent dose response curves. *** p< 0.001 vs control	128
7.4	Effect of DA 10 M, 20 M and 40 M on dose response curve of norepinephrine (NE) of endothelium intact aorta. *p<0.05 and *** p<0.001 vs. control	130
7.5	Effect of DDA 10 M, 20 M and 40 M on dose response curves of KCl on endothelium denuded aorta. ** p<0.01, *** p<0.001 vs control	132
7.6	Effect of DDA 10 M, 20 M and 40 M on concentration-response curves of norepinephrine (NE) of endothelium intact aorta. All data are expressed as means \pm SEM (n=8) ** and *** p<0.01 and 0.001 vs control	133
7.7	Effect of DA, DDA and APCE on the NE pre-contracted aortic rings in presence of endothelium. Results are presented as means of \pm SEM of eight experiments	135

7.8	Effect of DA, DDA, APCE and ANG on the NE pre-contracted aortic rings in absence of endothelium. Results are presented as means of \pm SEM of eight experiments	135
7.9	Inhibitory effects of APCE on the phasic and tonic contractions induced by NE. Phasic contraction was induced in Ca^{2+} free solution by the Ca^{2+} released from sarcoplasmic reticulum. Tonic contraction was induced by influx of extracellular Ca^{2+} through receptor-operated channels ** $p < 0.01$ vs tonic contraction	137
7.10	Inhibitory effects of DDA on the phasic and tonic contractions induced by NE. Phasic contraction was induced in Ca^{2+} free solution by the Ca^{2+} released from sarcoplasmic reticulum. Tonic contraction was induced by influx of extracellular Ca^{2+} through receptor-operated channels. ** $p < 0.01$ vs tonic contraction	138
7.11	Inhibitory effects of DA on the phasic and tonic contractions induced by NE. Phasic contraction was induced in Ca^{2+} free solution by the Ca^{2+} released from sarcoplasmic reticulum. Tonic contraction was induced by influx of extracellular Ca^{2+} through receptor-operated channels. ** $p < 0.01$ vs tonic contraction	139
7.11	Inhibitory effects of AP_1 on the phasic and tonic contractions induced by NE. Phasic contraction was induced in Ca^{2+} free solution by the Ca^{2+} released from sarcoplasmic reticulum. Tonic contraction was induced by influx of extracellular Ca^{2+} through receptor-operated channels. ** $p < 0.01$ vs tonic contraction	139
7.12	Effect of L-NAME (100 μM) in presence of DA, DDA, ANG and APCE. ($p > 0.05$) $n = 8$ experiments	140
7.13	Effect of indomethacin (10 μM) in presence of DA, DDA, ANG and APCE ($p > 0.05$) $n = 8$ experiments	141
8.1	Body weight gain (g) is expressed as difference between final and initial body weight. Each point represents mean \pm SEM 6 rats	154
8.2	Liver photomicrograph section of control (Tween-80) 10% (v/v) treated rat showing no visible lesions, cell swelling and maintaining liver architecture H & E, $\times 100$	155
8.3	Kidney (cortical part) photomicrograph section of control (Tween 80) 10% (v/v) treated rats showing normal vascular glomeruli, and tubular epithelium. Capillaries are filled with blood cells; some tubules contain single desquamated cells. H & E, $\times 100$	155
8.4	Heart photomicrograph section of control (Tween-80) 10% (v/v) treated rats showing absence of necrosis, deeply eosinophilic cytoplasm of myocytes. H&E, $\times 100$	156

8.5	Lungs photomicrograph section of control (Tween-80) 10% (v/v) treated rats showing presence of mild airway secretion in the lumen and lung parenchyma remains unaltered H&E, × 100	156
8.6	Spleen photomicrograph section of control (Tween 80) 10% (v/v) treated rats showing presence of hematopoietic cells, granulopoiesis seen in the centre of the photo erythroid cells and megakaryocytes also present H&E, ×100	157
8.7	Liver photomicrograph section of APCE 100 mg/kg treated rats showing hepatocytes are arranged in trabacules running radiantly from the central vein and are separated by sinusoids containing Kupffer cells with large spheroidal nucleus with distinctly marked nucleolus and peripheral chromatin distribution H&E, × 100	157
8.8	Kidney (cortical part) photomicrograph section of APCE 100 mg/kg treated rat showing renal glomeruli normal structure with renal tubules are lined with thick cubical epithelium. The tubules have relatively regular distinct lumen. Lobular organization of the glomerule and a flat epithelium lining the glomerular capsule can be seen H&E × 100	158
8.9	Heart photomicrograph section of APCE 100 mg/kg treated rat showing moderate necrosis deep eosinophilic cytoplasm H&E, ×100.	158
8.10	Lung photomicrograph section of APCE 100 mg/kg treated rats showing presence of mild airway secretion in the lumen and lung parenchyma remains unaltered H&E, ×100	159
8.11	Spleen photomicrograph section of APCE 100 mg/kg treated rats showing presence of hematopoietic cells, granulopoiesis, erythroid cells and megakaryocytes also present H&E, ×100	159
8.12	Liver photomicrograph section of APCE 300 mg/kg treated rats showing moderate necrosis hepatocytes arranged in trabacules running radiantly from the central vein and are separated by sinusoids containing Kupffer cells with large spheroidal nucleus with distinctly marked nucleolus and peripheral chromatin distribution H&E, ×100	160
8.13	Kidney (cortical part) photomicrograph section of APCE 300 mg/kg treated rats showing renal glomeruli normal structure with renal tubules are lined with thick cubical epithelium. The tubules have relatively regular distinct lumen. Lobular organization of the glomerule and a flat epithelium lining the glomerular capsule can be seen H&E, ×100	160
8.14	Heart photomicrograph section of APCE 300 mg/kg treated rats showing mild to moderate necrosis deep eosinophilic cytoplasm. H&E, ×100	161

8.15	Lung photomicrograph section of APCE 300 mg/kg treated rat showing presence of mild airway secretion in the lumen and lung parenchyma remains unaltered H&E, ×100	161
8.16	Spleen photomicrograph section of APCE 300 mg/kg treated rat showing presence of normal hematopoietic cells, granulopoiesis, erythroid cells and megakaryocytes also present H&E, ×100	162
8.17	Liver photomicrograph section of APCE 1000 mg/kg treated rats showing moderate infiltration and necrosis, hepatocytes arranged in a cord like structure and mild fatty degeneration is seen H&E, ×100	162
8.18	Kidney (cortical part) photomicrograph section of APCE 1000 mg/kg treated rat showing normal renal glomeruli, tightly filling the Bowman's capsule. The tubules have a relatively distinct lumen. Lobular organization of the glomerule and a flat epithelium lining the glomerular capsule can be seen. H&E, ×100	163
8.19	Heart photomicrograph section of APCE 300 mg/kg treated rats showing mild to moderate necrosis deep eosinophilic cytoplasm. H&E, ×100	163
8.20	Lung photomicrograph section of APCE 100 mg/kg treated rat showing presence of moderate airway secretion in the lumen and lung parenchyma remains unaltered. H&E, ×100	164
8.21	Spleen photomicrograph section of APCE 1000 mg/kg treated rat showing presence of venous sinuses, erythropoietic cells are scattered, granulocytes, macrophages and lymphocytes were also seen. H&E, ×100	164
8.22	Liver photomicrograph section of APCE 2000 mg/kg treated rat showing hexagonadal lobules with central veins and peripheral hepatic triads or tetrads embedded in connective tissue. Moreover liver architecture maintaining normal morphology	165
8.23	Kidney (medullary part) photomicrograph section of APCE 2000 mg/kg treated rat showing collecting tubules are lined with the relatively low simple cubic epithelium. The thick descending and ascending parts of Henle's loop and collecting coils H&E × 100	165
8.24	Lung photomicrograph section of APCE 100 mg/kg treated rat showing absence of airway secretion in the lumen and lung parenchyma remains unaltered. H&E, × 100	166
8.25	Heart photomicrograph section of APCE 2000 mg/kg treated rat showing absence of necrosis, deeply eosinophilic cytoplasm of myocytes H&E, ×100	166

8.26	Spleen photomicrograph section of APCE 2000 mg/kg treated rat showing granulopoiesis, erythroid cells and megakaryocytes are also present; both B & T cell regions are affected. H&E, ×100	167
9.1	Plasma calibration curve of ANG	179
9.2	Plasma calibration curve of DDA	179
9.3	HPLC chromatograms of blank rat plasma	180
9.4	HPLC chromatogram of ANG spiked in rat plasma (4.88 min)	180
9.5	HPLC chromatogram of DDA spiked in rat plasma (7.61 min)	181
9.6	HPLC chromatogram of APCE in rat plasma with ANG (4.88 min) and DDA (7.60 min)	181
9.7	Plasma concentration ANG and DDA versus time profiles determined after application of a single oral dose (1000 mg/kg) of <i>Andrographis paniculata</i> chloroform extract (APCE) in rats. Each point represents mean ± S.D. (n=6)	184

LIST OF ABBREVIATION & SYMBOLS

ACh	=	Acetylcholine
5-HT	=	5-hydroxytryptamine (Serotonin)
AC	=	Adenylyl cyclase
ACEI	=	Angiotensin converting enzyme inhibitors
ACN	=	Acetonitrile
ADP	=	Adenosine di-phosphate
AECB	=	Aqueous extract of <i>Caesalpinia benthamiana</i>
AhR	=	Aryl hydrocarbon receptor
ANG	=	Andrographolide
ANOVA	=	Analysis of variance
AP ₁	=	AP1- (sub-fraction)
AP ₃	=	14-deoxy-11, 12-didehydroandrographolide
AP ₄	=	Neoandrographolide
AP	=	<i>Andrographis paniculata</i>
APCE	=	<i>Andrographis paniculata</i> chloroform extract
ATP	=	Adenosine triphosphate
AUC	=	Area under the curve
BCG	=	Bacillus Calmette-Guein
CaCl ₂	=	Calcium chloride
cAMP	=	Cyclic Adenosine Monophosphate
cGMP	=	Cyclic Guanosine Phosphate
CK-MB	=	Creatine kinase
CO	=	Cardiac output
COX	=	Cyclooxygenase
CVD	=	Cardiovascular disease
CYPs	=	Cytochrome P450s
DA	=	14-deoxyandrographolide
DAG	=	Diacyl glycerol
DDA	=	14-deoxy-11, 12-didehydroandrographolide
DMSO	=	Dimethylsulphoxide

ECG	=	Electro cardio gram
EDNO	=	Endothelium-derived nitric oxide
EDRF	=	Endothelium-derived relaxant factors
EDTA	=	Ethylene diamine tetra acetic acid
EGCG(-)	=	Epigallocatechin- 3-gallate
ELT	=	Euglobulin lysis time
eNOS	=	Endothelial nitric oxide synthase
FO	=	Fish Oil
GM-CSF	=	Granulocyte macrophage colony stimulating factor
GMP	=	Guanosine monophosphate
GSTP	=	Glutathione S-transferase
GTP	=	Guanosine 5-triphosphate
H ₂ O ₂	=	Hydrogen peroxide
HCT	=	Haematocrit
HGB	=	Haemoglobin
HIV	=	Human immuno deficiency virus
HMP	=	Herbal medicinal plants
HOCl	=	Hypochlorous acid
HPBLs	=	Human peripheral blood lymphocytes
HPLC	=	High performance liquid chromatography
IP ₂	=	Inositol di phosphate
IP ₃	=	Inositol triphosphate
IP ₃	=	Inositol-1, 4, 5-trisphosphate
IP	=	Intraperitoneal
KCl	=	Potassium chloride
KH ₂ PO ₄	=	Potassium dihydrogen phosphate
LAD	=	Left anterior descending artery
LDH	=	Lactate dehydrogenase
LDL	=	Low-density lipoprotein
L-NAME	=	Nitro-L-arginine methyl ester
LPS	=	Lipopolysaccharide
LVEDP	=	Left ventricular end diastolic pressure
MAP	=	Mean arterial blood pressure

MARDI	=	Malaysian Agriculture Development Institute
MBP	=	Mean blood pressure
MCH	=	Mean corpuscular hemoglobin
MCHC	=	Mean corpuscular hemoglobin concentration
MCV	=	Mean cell volume
MLCK	=	Myosin light chain kinase
NaCl	=	Sodium chloride,
NaHCO ₃	=	Sodium bicarbonate
NE	=	Norepinephrine
NMR	=	Nuclear magnetic resonance
N	=	Nicotinic receptors
NO	=	Nitric oxide
NOS	=	Nitric oxide synthase
NSAID	=	Non-steroidal anti-inflammatory drugs
NSBP	=	Non-invasive systolic blood pressure
ODQ-1H	=	[1,2,4]oxadiazolo[4,2- α]quinoxalin-1-one
ODQ	=	Oxadiazole-[4,3-a]-quinoxalin-1-one
PAF	=	Platelet-activating factor
PBG	=	Peak blood glucose
PDE	=	Phosphodiesterase
PE	=	Phenylephrine
PGI ₂	=	Prostacyclin
PKC	=	Protein kinase C
PLC	=	Phospholipase C
PLT	=	Platelets count
PMA	=	Phorbol 12-myristate 13-acetate
PMNL	=	Polymorph-nuclear leukocytes
PMNs	=	Polymorphonuclear neutrophils
PPH	=	Postprandial hyperglycemia
PTFE	=	Polytetrafluoroethylene
PTLC	=	Preparative thin-layer chromatography
QPCR	=	Quantitative polymerase chain reaction
RBC	=	Red blood count

ROS	=	Reactive oxygen species
ROS	=	Reactive oxygen species
SBP	=	Systolic Blood Pressure
SD	=	Sprague-Dawley
sGC	=	Soluble guanylyl cyclase
SHR	=	Spontaneously hypertensive rats
SNP	=	Sodium nitroprusside
SOD	=	Super oxide dismutase
SPE	=	Solid phase extraction
TBA	=	Thiobarbituric acid
TCM	=	Traditional Chinese medicine
TIMP-1	=	Tissue inhibitors of metalloproteinase-1
TLC	=	Thin-Layer Chromatography
VEGF	=	Vascular endothelial growth factor
v/v	=	Volume in volume
WBC	=	White blood count
WE	=	Water extract
WHO	=	World Health Organization

LIST OF APPENDICES

		Page
Appendix A1	Effect of APCE (100 mg/kg) 28 day's treatment on heart, liver and kidney indices	227
Appendix A2	Effect of APCE (100 mg/kg) 28 day's treatment on heart, liver and kidney indices	228
Appendix A3	Control (10%Tween 80) treated group on heart, liver and kidney indices	229
Appendix A4	Control (10%Tween 80) treated group on heart, liver and kidney indices	230
Appendix A5	Effect of APCE (300 mg/kg) on heart, liver and kidney indices	231
Appendix A6	Effect of APCE (300 mg/kg) 28 day's treatment on heart, liver and kidney indices	232
Appendix A7	Effect of APCE (1000 mg/kg) 28 day's treatment on heart, liver and kidney indices	233
Appendix A8	Effect of APCE (2000 mg/kg) 28 day's treatment on heart, liver and kidney indices	234

LIST OF PUBLICATIONS

- 1 **Raghava Naidu. S.**, Amirin Sadikun & Mohd. Zaini Asmawi (2009). The Effect of extracts of *Andrographis paniculata* aerial parts on rat thoracic aorta. *Pharmacognosy Research [Phcog Res.]* 1 (2); 54-59.
- 2 **Raghava Naidu. S.**, Omar Z. Ameer, Ibrahim M. Salman, G. Venkatesh, Amirin Sadikun, and Mohd. Zaini Asmawi. (2009). Pharmacokinetic study of *Andrographis paniculata* on experimental animals. *Pharmacologyonline*, 1; 309-319.

LIST OF PRESENTATIONS

- 1 **Raghava Naidu.S**, Ibrahim M. Salman, Omar Z. Ameer, Amirin Sadikun, Mohd. Zaini Asmawi. Acute and Subacute Toxicological Study of Standardized Chloroform Extract of *Andrographis paniculata* on Experimental Animals. 13th Biological Graduate Conference. National University of Singapore, 15th – 18th December (2008).
- 2 **Raghava Naidu.S**, Asmawi. M.Z. and Amirin S. Chronic Treatment of Chloroform Extract of *Andrographis paniculata* Prevents the Endothelial Dysfunction in Spontaneously Hypertensive Rat Thoracic Aorta. Presented on Malaysian society for physiology and pharmacology, University Malaya, 6th April (2008).
- 3 **Raghava Naidu. S.**, Asmawi. M. Z. and Amirin. S. Vasorelaxant effect of chloroform extract of *Andrographis paniculata* on *in-vitro* rat thoracic aorta. Presented poster on workshop on the “Isolated Tissue Preparations and HPLC Training” June (2007) at IIUM & MSPP, Kuantan, Malaysia.

KAJIAN REAKTIVITI VASKULAR, PROFIL KIMIA, TOKSIKOLOGI DAN FARMAKOKINETIK EKSTRAK *ANDROGRAPHIS PANICULATA* (NEES.)

ABSTRAK

Tujuan kajian ini adalah untuk menilai reaktiviti vaskular, profil kimia, ketoksikan dan farmakokinetik ekstrak *Andrographis paniculata*. Ekstrak kloroform *Andrographis Paniculata* (APCE) didapati memberikan kesan vasorelaksasi poten keatas kontraksi aruhan norepinefrin (NE) pada aorta toraks tikus. Analisis HPLC dan ¹H-NMR APCE menunjukkan kehadiran andrografolida (ANG), 14-deoksiandrografolida (DA) dan 14-deoksi-11,12-didehidrografolida (DDA). Rawatan kronik selama 4 minggu dengan APCE 25, 50 dan 100 mg/kg/hari pada tikus hipertensif spontan (SHR) menunjukkan peningkatan relaksasi tergantung-endotelium dan tidak tergantung-endotelium terhadap asetilkolina (ACh) dan natrium nitroprusida (SNP) mungkin kerana pengaktifan nitrik oksida (NO) sintase dan juga perangsangan pengeluaran NO dalam sel-sel endotelium yang membawa kepada perencatan bagi lintasan kontraksi aruhan-Ca²⁺. Penurunan tekanan darah sistole (SBP) secara signifikan (p<0.001) mungkin disebabkan oleh tindakan vasodilatasi pada saluran darah.

APCE, sub-fraksi AP (AP₁), DA dan DDA bergantung dos merencat kedua-dua kontraksi tonik aruhan-NE dan kontraksi aruhan-K⁺ berkepekatan tinggi (80mM), mencadangkan yang APCE, AP₁, DA dan DDA bertindak sebagai penghalang saluran Ca²⁺ kepada kedua-dua saluran kendalian reseptor dan saluran tergantung potensi. DA, DDA dan APCE juga merencat kontraksi fasik aruhan-NE menandakan yang DA, DDA dan APCE merencat pembebasan Ca²⁺ daripada

retikulum sarkoplasma. Pengurangan kepekatan Ca^{2+} yang menyumbang kepada vasorelaksasi yang diaruhkan oleh DA, DDA dan APCE berkemungkinan melalui perencatan influks Ca^{2+} dan perencatan pembebasan Ca^{2+} dalam sel. Kesan perencatan DA, DDA dan APCE keatas kontrakasi fasik aruhan dos rendah NE mungkin bertindak melalui perencatan influks Ca^{2+} melalui lintasan bebas kalsium.

Akhirnya kajian ketoksikan akut dan ketoksikan kronik APCE 100, 300, 1,000 dan 2,000 mg/kg/hari tidak menunjukkan tanda-tanda ketoksikan sehingga ke akhir 28 hari jangkamasa penyelidikan. Tiada perubahan peningkatan berat badan mingguan dan profil hematologi serta perubahan profil makroskopik dan histopatologi organ dalaman semasa postmortem. Oleh itu, keputusan yang diperolehi mencadangkan yang APCE adalah tidak toksik sehingga 2,000 mg/kg. Dalam kajian farmakokinetik APCE menggunakan ANG dan DDA sebagai penanda menunjukkan farmakokinetik tak linear pada dos 1,000 mg/kg pada tikus.

**VASCULAR REACTIVITY, CHEMICAL PROFILE, TOXICOLOGICAL
AND PHARMACOKINETIC STUDIES OF *ANDROGRAPHIS PANICULATA*
NEES. EXTRACTS**

ABSTRACT

The aims of the study were to evaluate the vascular reactivity, chemical profile, toxicity and pharmacokinetic of *Andrographis paniculata* (AP) extracts. *Andrographis paniculata* chloroform extract (APCE) was found to be a potent vasorelaxant against norepinephrine (NE)-induced contraction of rat thoracic aorta. The HPLC and ¹H-NMR analysis of APCE revealed the presence of andrographolide (ANG), 14-deoxyandrographolide (DA) and 14-deoxy-11, 12-didehydroandrographolide (DDA). Chronic treatment for four weeks of APCE 25, 50 and 100 mg/kg/day in spontaneously hypertensive rats (SHR) demonstrated that it enhances the endothelium-dependent and endothelium-independent relaxation to acetylcholine (ACh) and sodium nitroprusside (SNP) presumably due to the activation of nitric oxide (NO) synthase and stimulation of the NO production in endothelial cells which lead to inhibition of Ca²⁺-induced contraction pathway. The systolic blood pressure (SBP) of SHR was significantly (p<0.001) reduced presumably due to its vasodilatory action on blood vessels.

APCE, sub-fraction of AP (AP₁), DA and DDA dose dependently inhibited both the NE-induced tonic contraction and high K⁺ (80 mM)-induced contraction, suggesting that APCE, AP₁, DA and DDA act as a Ca²⁺ channel blocker of both receptor-operated and potential-dependent channels. DA, DDA and APCE also dose dependently inhibited the NE-induced phasic contraction, suggesting that DA, DDA and APCE inhibits the Ca²⁺ release from sarcoplasmic reticulum. The reduction in

intracellular Ca^{2+} concentration that contribute to vasorelaxation induced by DA, DDA and APCE may be through inhibition of Ca^{2+} influx and inhibitions of intracellular calcium released. The inhibitory effect of DA, DDA and APCE on lower dose of NE-induced phasic contraction may act by inhibiting calcium influx through calcium-independent pathway.

Finally the acute and chronic toxicity studies of APCE at 100, 300, 1000 and 2000 mg/kg/day showed there was no visible sign of toxicity until the end of the 28 days study period. There were no significant changes observed on the weekly body weight gain and hematological profile as well as macroscopic and histopathological profile of the internal organs on post mortem. Therefore, the results obtained suggest that APCE is nontoxic up to 2000 mg/kg body weight. In pharmacokinetic study of APCE using ANG and DDA as markers showed non-linear pharmacokinetics at a dose 1000 mg/kg in rats.

CHAPTER 1

INTRODUCTION

1.0 Introduction

In the treatment of cardiac diseases several synthetic, semi-synthetic and natural drug molecules have been used from the past decades. Among the series of emerging drug candidates, few of active principles were also isolated from the natural flora and have been tested and used in various conditions of cardiovascular malfunctions. The physiology and etiology of cardiovascular tissues are important in the screening and development of new drug candidates.

1.1 Vascular smooth muscle

The etiology of cardiovascular disease depends on the structural integrity and health of the blood vessels of the cardiovascular system. The wall of an artery consist of three distinct layers namely tunica intima, tunica media and tunica adventitia. Tunica intima, the inner most layer of the artery wall, consists of a single layer of endothelial cells and connective tissue. The amorphous mucopolysaccharide ground substance containing elastin, collagen, and vascular smooth muscle cells is referred to as the tunica media layer. Tunica adventitia is the outermost layer surrounding the two inner layers and consists of strong fibrous tissue which maintains the shape of the vessel (Figure 1.1). Vascular smooth muscle is innervated primarily by the sympathetic nervous system through adrenergic receptors (adrenoceptors). Three types of adrenoceptors are present within vascular smooth muscle cells: alpha 1 (α_1), alpha 2 (α_2) and beta 2 (β_2). Norepinephrine is the main endogenous agonist for adrenoceptors.




Figure 1.1. Structure of artery and vein (Martini & Bartholomew, 2008).

1.2 Contraction of blood vessels

Smooth muscle contraction and the regulation of the contractile process has been the subject of intense studies for many years. Smooth muscle contraction is mainly composed of interlocked filaments held in place by a lattice work of fibres of dense bodies. The main contractile filaments are referred to as thick (contains myosin) and thin filaments (contains actin) owing to their microscopic appearance. Contraction occurs when these filaments slide over one another. This movement is mediated by the process of cross-bridge cycling. Variations in the free cytosolic calcium (Ca^{2+}) concentrations in vascular smooth muscle cells have been identified as the primary regulatory signal for smooth muscle contractions. The Ca^{2+} concentration in resting smooth muscle ranges between 80 to 270 nM while increase in the Ca^{2+} concentrations to 500 nM to 700 nM results in contraction (Webb, 2003).

Figure 1.2 (A) shows the presence of muscarinic (M_2) and muscarinic (M_3) receptors in most smooth muscle cells. Activation of M_3 receptors elicit contraction through stimulation of phospholipase C- β (PLC- β), which cleaves phosphatidylinositol-4, 5-bisphosphate into diacylglycerol (DAG) and inositol-1, 4, 5-trisphosphate (IP_3). The IP_3 mobilizes Ca^{2+} and triggers contraction. Beta (β)-adrenergic receptor activation stimulates adenylyl cyclase (AC) to generate cyclic adenosine-mono-phosphate (cAMP), which causes the relaxation of smooth muscle. M_2 receptors inhibit AC to prevent the relaxation effect on stimulation of the β -adrenergic receptor.

Figure 1.2 (B) illustrates that most peripheral blood vessels contain M_3 receptors on the endothelium, which trigger the synthesis of nitric oxide. Nitric oxide (NO) diffuses into the smooth muscle, where it mediates relaxation through the production of cyclic guanosine monophosphate (cGMP).

Figure 1.2 (C) shows the activation of M_1 receptors and nicotinic receptors (N) in parasympathetic ganglia causing the release of an inhibitory neurotransmitter (IN) from postganglionic neurons in gastrointestinal sphincters. This inhibitory neurotransmitter is usually adenosine triphosphate (ATP), NO or vasoactive intestinal peptide (VIP) and causes the sphincter smooth muscle to relax.




Figure 1.2. Receptor-mediated contraction and relaxation in different types of smooth muscle (Mineman and Wecker, 2004).

Vasoconstricting neurotransmitters and hormones bind to their receptors on the cell surface and initiate a series of processes leading to the contraction of vascular smooth muscle. Most receptors activate various types of guanosine 5-triphosphate (GTP) binding proteins (G-proteins), which are coupled to different ion channels and enzymes, and modulate their activities. These enzymes include both phospholipase C (PLC), which metabolises inositol diphosphate (IP_2) to produce inositol triphosphate (IP_3), diacylglycerol (DAG). Adenylate cyclase metabolises

ATP to produce cAMP. IP₃ releases Ca²⁺ from intracellular stores whereas DAG activates protein kinase C (PKC), which phosphorylates a number of proteins. In addition to the activation of the IP₃ metabolism, vasoconstrictors such as norepinephrine have been shown to depolarize the smooth muscle cells and consequently activate voltage operated Ca²⁺ channels in the plasma membrane of the smooth muscle, leading to an increased influx of extra cellular Ca²⁺. Moreover, the existence of receptor operated Ca²⁺ channels have been proposed in smooth muscle cells (Webb, 2003). The activation of this mechanism increases intracellular Ca²⁺, which is the primary signal for smooth muscle contraction (Bolton, 1979; Karaki *et al.*, 1984; Allen & Walsh, 1994). As a consequence of elevated intracellular Ca²⁺ concentration, Ca²⁺ binds to calmodulin to form Ca²⁺- calmodulin complex, which removes the auto inhibition of myosin light chain kinase (MLCK). The activated MLCK phosphorylates reversibly the light chain of myosin and activates the myosin ATPase. The phosphorylated myosin cyclically binds to actin filaments producing force or the shortening of the smooth muscle (Figure 1.3). The contractile force does not, however, depend directly on intracellular Ca²⁺, since the contractile force may be enhanced by increasing the responsiveness of the contractile machinery or the sensitivity of the myofilaments to intracellular Ca²⁺ (Webb, 2003). These modulatory mechanisms for changing the Ca²⁺ metabolism, serve an important role in the regulation of vascular smooth muscle tone (Somlyo *et al.*, 1999; Webb, 2003).




Figure 1.3. Mechanisms of contraction of vascular smooth muscle cells (Mineman and Wecker, 2004).

1.3 Smooth muscle relaxation

Relaxation of smooth muscle requires a fall in intracellular Ca²⁺ levels to resting levels and dephosphorylation of myosin. Smooth muscle relaxation occurs either as a result of removal of the contractile stimulus or by direct action of a substance that stimulates the inhibition of the contractile mechanism. This process is catalyzed by a specific myosin light chain phosphatase (MLCP) (Somlyo *et al.*, 1999). Alterations in the mechanisms that lead to reduction in intracellular Ca²⁺ levels and/or increase in MLCP activity may contribute to alterations in responsiveness of smooth muscle cells. Several mechanisms have been implicated in the sequestration or removal of cytosolic Ca²⁺. For an instance, the inhibition of sarcoplasmic reticular Ca²⁺ and Mg²⁺-ATPase activity which mediates the release of intracellular Ca²⁺ leads to reduction in cytosolic Ca²⁺ concentrations and hence causes relaxation of smooth muscle cells (Webb, 2003). In addition, the inhibition of receptor-operated and voltage-operated Ca²⁺ channels located in the plasma

membrane which are important in the Ca^{2+} influx and smooth muscle contraction, leads to the reduction in intracellular Ca^{2+} concentrations and hence causes smooth muscle relaxation (Webb, 2003).

1.4 Endothelium

1.4.1 Physiology of endothelium

The vascular endothelium is the largest endocrine organ in the body. It is approximately 14,000 square feet in surface area with a size of 6.5 tennis courts in area and five times the heart size in mass with a total weight of about 2 kg (Amudha *et al.*, 2002). The vascular endothelium under normal, healthy physiological conditions forms a continuous sheet of organized monolayer polyhedral cells. The endothelial cells are tightly interlocked so that passage of products from the blood occurs through the endothelial cell. These cells are both a passive filter and a metabolically active organ that synthesizes and release several vasoactive substances into the blood and into the underlying vascular smooth muscle cells, which regulates the vascular homeostasis.

The endothelium-derived vasoactive substances include vasodilators, such as NO, prostacyclin and yet unidentified endothelium-derived hyperpolarizing factor and also vasoconstrictors, such as free radicals, cyclooxygenase products and endothelin-I. Endothelium-derived vasoactive substances alter the vascular tone of the artery through myogenic mechanisms, local hormones or chemical substances, and/or metabolic by-products. Consequently, signal transmission causes muscle contraction or muscle relaxation and can occur through numerous pathways involving nerve signals, blood borne substances and locally generated substances.

For example, endothelium-derived vasoconstrictors typically bind to receptors on the smooth muscle cells and can elicit a contraction through enhancing intracellular Ca^{2+} concentrations. On the other hand, endothelium-derived vasodilators act in several ways to protect the integrity of the artery, chief among which are induction of vascular smooth muscle cell relaxation, inhibition of vascular smooth muscle cell growth, inhibition of platelet aggregation and thrombosis and inhibition of monocyte adhesion.

1.5 Endothelium and cardiovascular disease

Endothelial cells produce several biologically active substances that play key role in local regulation of blood flow, blood pressure and vascular tone (Furchgott & Vanhoutte, 1989; Luscher *et al.*, 1995; Gewaltig *et al.*, 2002; Endemann & Schiffrin, 2004). It is not surprising that alterations in physiological functions of endothelial cells have the potential to contribute directly to the impaired vascular homeostasis and hence to pathogenesis of cardiovascular disease. In support of this hypothesis, data from clinical and experimental studies have demonstrated that impairment of endothelial function either initiates or associates with the development and progression of cardiovascular disease (Endemann & Schiffrin, 2004). Endothelial dysfunction can be determined by the reduction in the activity of endothelium-derived vasodilators, mainly NO by local increases in antagonists (endothelium-derived contractile factors) to these vasodilators, or by other associates of these two factors. On the other hand, considerable evidence suggests that the manifestation of endothelial dysfunction occur before the development of cardiovascular disease (CVD) and endothelial function has been reported to be impaired even in the offspring of CVD parents (Taddei *et al.*, 1992; Taddei *et al.*, 1996; Amudha *et al.*,

2002). This shows that the onset of endothelial dysfunction may be an important pathogenic event preceding the development of clinically evident vascular disease.

Over the last decade, extensive research has focused on determining the presence and nature of endothelial dysfunction in experimental models of CVD and in patients with CVD (Amudha *et al.*, 2002). Investigations into the mechanisms of endothelial dysfunction both in hypertension and diabetes mellitus have demonstrated that reduced bioactivity and/or bioavailability of endothelium-derived nitric oxide plays an important role (Luscher & Noll, 1995; Endemann & Schiffrin, 2004). Several factors including low-density lipoprotein (LDL) cholesterol oxidation, increased production of reactive oxygen species (ROS) and decreased production of nitric oxide via endothelial nitric oxide synthase (eNOS) have been identified as important etiological factors in the reduced bioavailability and/or bioactivity of endothelium-derived nitric oxide and subsequently, in the development of endothelial dysfunction (Amudha *et al.*, 2002; Endemann & Schiffrin, 2004; Kalinowski & Malinski, 2004; Forestermann, 2005). An important implication of the free radical-oxidative stress hypothesis of endothelial dysfunction is that a wide range of antioxidants, including ascorbic acid and α -tocopherol, may inhibit ROS and reestablish endothelial function (Taddie *et al.*, 1998; Endemann & Schiffrin, 2004). Up to date most interventions attempting to improve endothelial dysfunction have targeted one or more of the numerous risk factors that can cause endothelial damage (Amudha *et al.*, 2002).

Many pharmacological agents have been suggested to achieve vascular protection through various mechanisms such as reduction in the blood pressure, inhibition of hyperglycemia and reduction in oxidative stress. Beneficial changes to the endothelium from these interventions might result from promotion of vascular relaxation, inhibition of vasoconstriction, reduction in the production of free radicals or other mechanisms that protect the endothelium from injury (Amudha *et al.*, 2002). The plants showing nitric oxide production can be a promising candidates for vasorelaxation, which may have potential lead molecule for preventing and treating cardiovascular diseases such as hypertension and atherosclerosis (Park *et al.*, 2009).

1.6 Vascular mediators from the endothelium: Endothelium-derived relaxant factors (EDRF)

The discovery of endothelium-derived relaxing factor and its identification as nitric oxide, a highly reactive free radical gas, is one of the most exiting discoveries of biomedical research in the last two decades (Furchgott & Zawadzki, 1980; Plamer *et al.*, 1987). Over the past few years, NO has become established as a universal intercellular messenger that serves a variety of biomodulatory functions in physiological as well as pathological conditions. NO is synthesized from L-arginine by the enzymatic action of nitric oxide synthase (NOS), i.e., endothelial, neuronal and inducible, exist in mammalian cells. The endothelial subtype accounts for the majority of the basal and stimulated NO synthesis in endothelial cells throughout the vasculature (Figure 1.4). The average half-life of NO in tissue is about 3 to 6 seconds whereas in blood it is 1 to 2 seconds.

The synthesis of NO can be stimulated by an increase in endothelial Ca^{2+} concentration following physical and chemical stimuli such as shear stress and hypoxia, activation of cell surface receptors by a variety of endogenous substances like acetylcholine (ACh) and bradykinin, or application of Ca^{2+} channel agonists (Vanhoutte *et al.*, 1995; Luscher & Noll, 1995; Hansen & Nedergaard, 1999). After endothelial NOS is turned on by Ca^{2+} flux, it biosynthesizes NO in bursts for about a minute and it is turned off by phosphorylation. On the other hand, it has also been (Schulz & Triggle, 1994) reported that there is a continuous and spontaneous basal release of NO from the endothelium, the amount of which regulates the arterial tone. Thus, impaired mechanism and/or inhibition of the NO production from endothelium causes dramatic decrease in blood flow and can certainly induce profound and sustained hypertension. This view is supported by the findings that chronic inhibition of NOS activity leads to a present of hypertension in experimental animals (Pechanova *et al.*, 2004).




Figure 1.4. Biosynthesis of L-Arginine – Nitricoxide (NO) pathway.

It has been suggested that there is a diminished basal NO synthesis in the vasculature of patients with hypertension and diabetes as well as in experimental hypertension and diabetes (Endemann & Schiffrin, 2004). On the other hand, considerable number of studies suggests that the synthesis and release of NO is unaffected or it may even be enhanced, but excessive production of super oxide anions leads to increased incapacitation of NO and subsequently, attenuates endothelium-dependent vasodilatation in hypertensive and diabetic arteries (Tschudi *et al.*, 1996; Maffei *et al.*, 2002). In this perspective, it is suggested that other than synthesis/release of NO, activation of eNOS may also lead to release of superoxide anions in higher quantities in hypertensive and diabetic arteries (Milstein & Katuski,

1999; Forstermann, 2005). In physiological conditions, the concentration of superoxide radicals remains low within the organism as a result of its reaction with superoxide dismutase (SOD) enzyme. However, in hypertension and diabetes mellitus, there may be an increase in the production of these radicals or deficiency of SOD (Gewaltig & Kodja, 2002; Endemann & Schiffrin, 2004).

NO plays different roles depending on the site of its production. NO synthesized by eNOS diffuses out in all directions. About 10% to 30% of NO diffuses to the wall of blood vessels and triggers a cascade of events leading to smooth muscle relaxation. NO relaxes vascular smooth muscle via the activation of soluble guanylate cyclase that converts guanosine monophosphate (GMP) to cyclic GMP (Moncada *et al.*, 1991; Hansen & Nedergaard, 1999). In smooth muscle cells, cGMP has been reported to activate the cGMP-dependent protein kinase that regulates several pathways involved in Ca^{2+} homeostasis with the end result being a reduction in the concentration of intracellular Ca^{2+} available for contraction and a decrease in the sensitivity of contractile proteins to Ca^{2+} (Moncada & Higgs, 1993; Hansen & Nedergaard, 1999). In addition, NO has been reported to hyperpolarize vascular smooth muscle via cGMP-dependent mechanisms as well as directly activating Ca^{2+} activated K^+ channels (Kca) and Na^+/K^+ adenosine 5'-triphosphate (ATPase) activity (Gupta *et al.*, 1994; Cohen *et al.*, 1995).

1.6.1 Prostacyclin

Prostacyclin (PGI_2) is the major relaxant prostanoid produced by the vascular endothelial cells. The formation of PGs begins with the liberation of arachidonic acid from cell membrane phospholipids by phospholipase A_2 . Arachidonic acid is

then converted into PGG₂ and PGH₂ by the enzyme cyclooxygenase (COX). Finally, PGH₂ is converted to PGI₂ by the action of PGI₂ synthase (Gryglewski, 1995). Although other PGs (PGE₂, PGF_{2α} and PGD₂) are also synthesized in the endothelial cells, PGI₂ is the major vasodilator prostanoid in all the vascular cells. Physiologically, PGI₂ is a local rather than circulating hormone, because its blood levels are too low to have any general effects. The production of PGs can be blocked by non-steroidal anti-inflammatory drugs, such as indomethacin and aspirin, which inhibit both isoforms i.e. COX 1 & COX 2 - activity (Akpaffiong & Taylor, 1998; Nascimento *et al.*, 2003). Recently another catalytically active COX variant, COX-3, derived from an alternative splicing of the COX-1 gene, has been identified in the brain (Chandrasekharan *et al.*, 2002).

Endothelial cells release PGI₂ in response to shear stress, hypoxia, and stimulation of various receptors on endothelial membrane (Luscher & Noll, 1995). Moreover, it is suggested from *in vitro* experiments that NO activates PGI₂ synthesis by activation of prostaglandin-H synthase and possibly by increasing cyclooxygenase activity by a cGMP-independent pathway (Gryglewski, 1995). PGI₂ exerts its vasodilatory actions by binding to membrane receptors on the smooth muscle, which activate adenylate cyclase and subsequently increase the intracellular concentration of cAMP. The elevation of intracellular cAMP leads to the reduction of intracellular Ca²⁺ concentration and to decrease in the sensitivity of contractile proteins to Ca²⁺ (Cohen & Vanhoutte, 1995). When compared with the inhibition of eNOS, the blockade of COX has negligible impact on blood pressure. On the other hand, the inhibition of COX has been found to enhance endothelium-mediated vasodilatation in hypertensive and diabetic rat arteries (Taddei *et al.*, 1997;

Akpaffiong & Taylor, 1998; Nascimento *et al.*, 2003). Therefore, hypertension and diabetes appear to be associated with an imbalance in endothelial production of COX-derived vasodilator and vasoconstrictor factors (figure 1.5).




Figure 1.5. Biosynthesis of prostaglandins, lipoxigenase and cytochrome P- 450 pathway.

1.6.2 Endothelium-derived hyperpolarizing factor (EDHF)

Defined as a group of yet unidentified substances which produce vascular smooth muscle hyperpolarization and relaxation. The nature of the responses attributed to EDHF is still unresolved, but the evidence from several sources suggests that there are multiple EDHFs, and that the chemical mediators of the EDHF response may vary with the vascular bed (Edwards & Weston, 1998; McGuire *et al.*, 2001). In recent years, the most popular candidates for EDHF have been the non-prostanoid products of the metabolism of arachidonic acid, namely epoxyeicosatrienoic acids (Campbell *et al.*, 2003; Archer *et al.*, 2003; Gauthier *et al.*, 2005). EDHF has been reported to be diffusible factor, which causes the opening of

K⁺ channels in the smooth muscle membrane (Quilley *et al.*, 1997; Edwards & Weston, 1998; McGuire *et al.*, 2001; Triggle *et al.*, 2003). Involvement of adenosine 5'-triphosphate (ATP)-sensitive K⁺ channels (K_{ATP}) and Na⁺/K⁺-ATPase has been reported in some vessels, but in the majority of the studies EDHF has been suggested to act through Ca²⁺ activated K⁺ channels (McGuire *et al.*, 2001). The action of EDHF can be inhibited by K⁺ channel blockers or by depolarizing the smooth muscle with increasing extra cellular high K⁺ concentrations (Adeagbo *et al.*, 1993; Ueda *et al.*, 2005).

The release of EDHF is as similar as that of NO and PGI₂ which is initiated by an increase in intracellular free Ca²⁺ concentration in the endothelial cell (Cohen & Vanhoutte, 1995; Luscher & Noll, 1995). Consequently, many autocoids and hormones that release NO and PGI₂ have also been shown to release EDHF (Cohen & Vanhoutte 1995). The mechanisms whereby hyperpolarization causes relaxation remain controversial (Triggle *et al.*, 2003). Most likely, the hyperpolarization of the smooth muscle cell membrane reduces Ca²⁺ influx through voltage dependent Ca²⁺ channels, which allows the Ca²⁺ sequestration and removal of lower intracellular free Ca²⁺ concentration. The importance of EDHF in endothelium-dependent relaxations has been reported to increase as the artery size decreases (McGuire *et al.*, 2001; Triggle *et al.*, 2003). It is also suggested that EDHF-mediated relaxation is upregulated in the pathophysiological states, such as hypertension and diabetes, which are associated with reduced bioavailability of NO (Ding *et al.*, 2000; McGuire *et al.*, 2001; Endemann & Schiffrin, 2004). Decreased endothelium-mediated hyperpolarization has been observed in many forms of experimental hypertension

and diabetes (McGuire *et al.*, 2001; Triggle *et al.*, 2003; Endemann & Schiffrin, 2004).

Hydrogen sulphide (H₂S) is increasingly being recognized as an important signalling molecule in the cardiovascular and nervous systems (Csaba Szabó (2007)). The production of H₂S from L-cysteine is catalysed primarily by two enzymes, cystathionine γ -lyase and cystathionine β -synthase. Evidence is accumulating to demonstrate that inhibitors of H₂S production or therapeutic H₂S donor compounds exert significant effects in various animal models of inflammation, reperfusion injury and circulatory shock. H₂S can also induce a reversible state of hypothermia and suspended-animation-like state in rodents (Csaba Szabó (2007)). Intravenous bolus injection of H₂S transiently decreased the blood pressure of the rats by 12 ± 30 mmHg which was antagonized by prior blockade of K_{ATP} channels (Zhao *et al.*, 2001). H₂S also relaxed rat aortic tissues *in vitro* in a K_{ATP} channels-dependent manner (Zhao *et al.*, 2001).

Over the last decade, studies have unraveled many aspects of endogenous production and physiological functions of carbon monoxide (CO). The majority of endogenous CO is produced in a reaction catalyzed by the enzyme heme oxygenase (HO). Inducible HO (HO-1) and constitutive HO (HO-2) are mostly recognized for their roles in the oxidation of heme and production of CO and biliverdin, whereas the biological function of the third HO isoform, HO-3, is still unclear (Wu & Wang, 2005). The interaction of CO and ion channels constitutes an important mechanism for the biological effect of CO. Most noticeable among the CO-targeted ion channels are K⁺ channels. This superfamily is composed of voltage- dependent K_v, ATP-

sensitive KATP, and calcium activated KCa channels (Wu & Wang, 2005). The interaction of CO and K⁺ channels may be the dominant force in driving the CO-induced vasorelaxation in specific types of blood vessels, especially peripheral resistance and cerebral arterioles. In other types of blood vessels, CO effects on K⁺ channels may become less important compared with activation of the cGMP pathway by CO (Wu & Wang, 2005).

Barkoudah et al. (2004) recently showed that application of a CO-releasing compound dilated arteriolar branches of the middle cerebral artery from piglets by activating BKCa channels in vascular smooth muscle cells. However, removal of endothelium or blocking the sGC-cGMP pathway abolished CO-induced vasorelaxation. Release of NO from endothelium or activation of sGC-cGMP pathway in vascular SMC allowed CO to cause vascular dilation (Wu & Wang, 2005).

1.7 Hypertension

Hypertension is one of the leading causes of cardiovascular and cerebrovascular complications, the most common reason to visit physician offices, and number one reason for drug prescriptions all over the world (Kearney *et al.*, 2005). Hypertension is characterized by a normal cardiac output and elevated arterial pressure. The etiology of hypertension is multifactorial and the precise mechanisms are not completely understood. Hypertension is a consequence of the interaction between genetics and the internal environment of the body.

Two forms of hypertension have been described:

1. Primary or essential hypertension
2. Secondary hypertension

Essential hypertension is a far more common condition and accounts for 95% of all cases of hypertension whereas secondary hypertension accounts for only 5%. Endothelial dysfunction and vascular smooth muscle dysfunction are initiating and perpetuating factors in essential hypertension (Luscher & Noll, 1990; Amudha *et al.*, 2002; Endemann & Schiffrin, 2004). In hypertension, the cardiovascular system exhibits several important physiological and morphological changes, chief among which are increase in the blood pressure, cardiac hypertrophy (increased cardiac muscle mass of blood vessels), impairment in vascular contraction and relaxation (Taddie *et al.*, 1998; Ludwig *et al.*, 2002; Sabbatini *et al.*, 2002; Endeman & Schiffrin, 2004). However, inspite of several studies on animal models of hypertension as well as in hypertensive human subjects, the exact mechanisms responsible for these pathological changes remain uncertain. It is suggested that the narrowing of the blood vessels contributed to increased peripheral vascular resistance and consequently to elevated blood pressure. This view is supported by the regression observed in the abnormal structure of blood vessels in hypertension towards physiological value with antihypertensive treatments (Ludwig *et al.*, 2002; Sabbatini *et al.*, 2002).

In recent past, the pharmacological research is focused on multidisciplinary drug discovery approach on isolation, synthesis and evaluation of emerging vasodilating drug molecules with maximum therapeutic effect with minimum or no

side effects. The plants which can enhance the NO-cGMP pathway and a direct action on the vascular smooth muscle through a dephosphorylation of myosin light chain kinase, resulting in vasodilation.

1.8 Some pharmacologically active moieties from natural flora used in cardiovascular malfunctions

Several coumarin derivatives have been shown to possess cardiovascular properties. Many of them are selective coronary vasodilators, an effect that may be related to a Ca^{2+} -antagonistic activity. Carbochromen (3-diethylaminoethyl-7-ethoxycarbonylmethoxy-4-methylcoumarin) is a potent specific coronary vasodilator which has been used for many years in the treatment of angina pectoris. Although the exact mechanism of action remains still unknown, it has been reported that carbochromen coronary effects could be mediated by an increased release of prostaglandins. Khellin (2-methyl-5, 8-dimethoxyfurochromone) is an active principle obtained from *Ammi visnaga* L. has strong vasodilator and spasmolytic activities. It probably decreases the availability of Ca^{2+} required for smooth muscle activation acting at multiple sites (Toimil *et al.*, 2002).

Vasorelaxing activity of the aqueous extract of *Caesalpinia benthamiana* (AECB) roots was tested using isolated rat aortic rings precontracted by phenylephrine (PE). Interaction of AECB with NO generation was also investigated by quantitative polymerase chain reaction (QPCR) analysis and its antioxidant properties were assessed by using human polymorphonuclear neutrophils (PMNs) in a cellular pathophysiological model of oxidative burst. Scavenging activities versus

superoxide anion ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), and hypochlorous acid ($HOCl$) were evaluated in the cell-free system (Alexis *et al.*, 2008).

The *in vitro* and *ex vivo* suppressive effects of *Andrographis paniculata* on nitric oxide production in mouse peritoneal macrophages is elicited by Bacillus Calmette-Guein (BCG) and stimulated by lipopolysaccharide (LPS). Incubation of BCG-induced macrophages with the methanol extract of *A. paniculata* reduced LPS stimulated NO production. The diterpene lactones andrographolide and neoandrographolide were isolated as active components from the extract. These compounds suppressed NO production in a concentration-dependent manner in the concentration range from 0.1 to 100 μ M and their IC_{50} values were 7.9 and 35.5 μ M. Neoandrographolide at doses of 5 and 25 mg/kg/day suppressed NO production by 35% and 40%. However, andrographolide did not reduce NO production on oral administration at the same doses. These results indicate that neoandrographolide, which inhibited NO production both *in vitro* and *ex vivo* may play an important role in the use of *A. paniculata* as an anti-inflammatory crude drug (Javzan *et al.*, 2002).

1.9. Conventional antihypertensive agents

Antihypertensive agents act at one or more of the four anatomic control sites and produce their effects by interfering with normal mechanisms of blood pressure regulation. A useful classification of these agents categorizes them according to the principal regulatory site or mechanism on which they act (Chobanian *et al.*, 2003).

- 1) Diuretics, which lowers blood pressure by depleting the body sodium ion concentration and reducing blood volume and by other mechanisms

Four basic types of diuretics are used in treatment of hypertension.

- a) Thiazide diuretics: Hydrochlorothiazide, chlorthiazide, indapamide *etc*
- b) High ceiling diuretics: Furosemide, bumetanide, ethacrynic acid
- c) Aldosterone antagonist: Spironolactone
- d) Angiotensin II receptor: Losartan, irbesartan, valsartan, telmisartan.

2) Drugs modifying sympathetic nervous system activity

a) Centrally acting drugs: Clonidine and methyldopa are believed to act upon vasomotor centers in the brain to decrease peripheral sympathetic nervous system tone. The drugs decrease cardiac output and peripheral resistance.

b) Inhibitors of neurotransmitter release and/or storage, guanethidine are actively transported into adrenergic nerve endings and inhibit norepinephrine release with nerve stimulation. Nerve uptake is necessary for the drug's antihypertensive actions.

c) Adrenergic receptor blockers

- i) α -adrenergic receptor antagonists: Prazosin, doxazosin, terazosin

Prazosin's antihypertensive actions are mentioned via competitive blockade of α_1 -receptor in peripheral arterioles, reducing vascular resistance. Prazocin is selective α_1 -receptor blocker its first dose produce severe hypotension.

- ii) β -adrenergic receptor antagonist: Propranolol, acebutolol, atenolol, metoprolol

β -adrenergic receptor antagonists reduce blood pressure by reducing heart rate and myocardial contractility (reduced cardiac output). These are considered as effective agents for mild to moderate hypertension.

3) Combination alpha/beta adrenergic receptor blockers: Labetalol, trandate

Labetalol exhibits both selective α_1 -adrenoceptor blockade and non-selective β -adrenoceptor blockade. The antihypertensive action of the drugs is results from a decrease in peripheral resistance with little or no decrease on cardiac out put. The drug is most useful for the treatment of mild to moderate hypertension.

3) Vasodilators

- a) Direct vasodilators: Hydralazine, minoxidil, diazoxide, sodium nitroprusside

Direct vasodilators act upon vascular smooth muscle to produce a relaxation of vascular tone and a decrease in peripheral resistance.

- b) Calcium entry blockers: Isradipine, nifedipine, diltiazem, amlodipine, felodipine, lercanidipine and verapamil hydrochloride.

The calcium entry blockers inhibit calcium entry into myocardial and vascular smooth muscle cells. The drugs can decrease peripheral resistance and decrease cardiac output the relative sensitivity of vascular smooth muscle and cardiac tissue vary with the three prototype agents. They are used for treatment of mild or moderate hypertension.

4) Angiotensin converting enzyme inhibitors (ACE Inhibitors): Captopril, esinopril, ramipril, perindopril.

CHAPTER 2

REVIEW OF LITERATURE

2.0 *Andrographis paniculata*

Malaysia is known as one of the 12 mega-diversity centres harbouring a multitude of medicinal plant species each presumably studded with as yet unknown genetic and chemical variations of economic importance. Several medicinal plants occurring in Malaysia species are used over several centuries in the traditional systems of medicine. Nearly 75% of the herbal drugs and perfumery products used in the world are available in natural state. Therefore, the rich and varied plant diversity, especially the genetic diversity of medicinal and aromatic plants, is one of the country's important strengths and is the bedrock for all future bio-industrial developments. Unfortunately, the renowned medicinal plant wealth of Malaysia has seldom been subjected to genetic scrutiny keeping in mind the latent and patentable properties and economic utility of the selected plant types. As severe habitat losses and consequent endangerment and extinction of known and hitherto lesser known species of economic value are not uncommon in this country, it is imperative that heritable variations within the otherwise unimproved natural populations of prospective taxa are studied for selection, improvement and development of suitable cultivars. Otherwise called bio-prospecting, this line of research is essential to fish out useful genes and gene products for commercialisation in the now unfolded patent regime. Knowledge of the genetic diversity is also a prerequisite for any *in situ* and *ex situ* conservation schemes (Hamrick *et al.*, 1991) as it is not practical to conserve all genotypes of a given species against the mass extinction spasm projected for the 21st century (Raven, 1999).