# CLINICAL IMPRESSION AND ULTRASOUND FEATURES IN THE PREOPERATIVE EVALUATION OF OVARIAN MASSES

#### by

#### DR CHEONG YEE TSING

Dissertation Submitted In
Partial Fulfilment Of The
Requirements For The Degree Of
Master Of Medicine
(Obstetrics and Gynaecology)

UNIVERSITI SAINS MALAYSIA NOVEMBER 2001

# **TABLE OF CONTENTS**

	Page
ACKNOWLEDGEMENTS	iii
1. INTRODUCTION TO:	
1.1 THE STATE OF PAHANG	iv
1.2 HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAH	HANG vi
1.3 DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY	Υ,
HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAH	HANG vii
2. ABSTRAK	xiii
3. ABSTRACT	xv
4. INTRODUCTION	1
5. MATERIALS AND METHODS	23
6. RESULTS	32
7. DISCUSSION	50
8. CONCLUSION .	77
9. LIMITATIONS	79
10. RECOMMENDATIONS	80
11. REFERENCES	81
12. APPENDIX	96

### **ACKNOWLEDGEMENTS**

I would like to acknowledge the invaluable guidance given to me by Dr Ghazali Ismail, Consultant Obstetrician and Gynaecologist, Head of the Department of Obstetrics and Gynaecology, Hospital Tengku Ampuan Afzan, Kuantan and Dr Mokhtar Awang, Consultant Obstetrician and Gynaecologist, Hospital Tengku Ampuan Afzan, Kuantan throughout the preparation for this dissertation and my posting in Kuantan.

I would also like to express my appreciation of the effort of Dr Awang Nila Ismail, Head and Consultant, Department of Obstetrics and Gynaecology, Hospital Universiti Sains Malaysia who painstakingly checked and corrected this dissertation. This book would not have been possible without their help.

I am also much indebted to everyone who has helped me — medical colleagues who have made helpful suggestions. I wish to thank each and every one of them for their efforts, and last but not least my parents for their love and encouragement.

#### INTRODUCTION TO THE STATE OF PAHANG

Pahang is the largest state in Peninsular Malaysia and is situated in the eastern coastal region. The state's 35,964 square kilometres encompasses a remarkable range of Malaysia's many different environments from the majestic peaks and cool hill regions of the state's western region to the miles of soft sand beach along the South China Sea. It is home to Taman Negara (National Park), all the major hill resorts in the country, and has Gunung Tahan, the highest peak in the peninsula.

This is a state that has a rich cultural tradition, of an ancient history. Stone artifacts point to man having existed here since the Stone Age. Later Indian empires, the Sri Vijaya and the Majapahit, spread their rule, customs and traditions over the people. Then came the Siamese and later the Melaka Sultante annexed the state. Others were to follow until finally the British brought the state into its ambit with the creation of the Federated Malay States. Today, the one million population constitutes a mixture of Malays, Chinese, Indians and aborigines, the Orang Asli.

The state capital, Kuantan, is really your first stop to a fascinating holiday. It is a contemporary town, growing fast yet retaining its ancient charms. About 48 km from Kuantan is Sungai Lembing, an old steel and tin mining town, where the deepest lode mine in the world was once very active. Relics of European architecture can still be found in the town.

All this has left Pahang with a heritage of colour, pomp and pageantry. Along with it the state inherited a beautiful coastline that is dotted with picturesque beaches and thousands of villages. The state is now one of the leading tourist destinations in the country.

Pahang provides the visitor with almost unspoilt natural splendours. There is myriad of exotic flora and fauna to bewilder you. There are pristine waterfalls along with palm-fringed sandy beaches. The state has an air of unspoilt beauty, a tropical holiday, an unforgettable journey that spells adventure.

# HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAHANG

Hospital Tengku Ampuan Afzan (HTAA) started its service in the year 1904. Over the years, it has undergone several stages of development and is now an eight-storey complex which is well equipped with medical facilities.

HTAA is the main hospital for the State of Pahang which is one of the thirteen component states of Malaysia. The hospital consists of 700 official beds and around 1,200 staff. It serves the town of Kuantan which is the capital town of the State of Pahang. It also serves as the referral centre for the seven outlying district hospitals, outpatient clinics and the private medical practitioners.

Since 1999, HTAA has become the teaching hospital for the Faculty of Medicine, International Islamic University (Universiti Islam Antarabangsa, UIA) and undertakes undergraduate teaching.

HTAA is recognized by the Health Ministry of Malaysia and local universities for the training of postgraduate students in the field of Obstetrics and Gynaecology, Surgery, Medicine, Paediatrics, Orthopaedics, Otorhinolaryngology and Family Medicine.

# Department of Obstetrics and Gynaecology Hospital Tengku Ampuan Afzan, Kuantan, Pahang

The Department of Obstetrics and Gynaecology is staffed by five Consultants/Specialists, six postgraduate students/registrars in training, five medical officers and a minimum of ten house officers.

The department has 36 antenatal beds and 44 postnatal beds. New cases are seen in the screening room after which they will either be admitted to the appropriate ward or observed in the screening room where we also provide day care services. The first class ward has 12 beds for antenatal and postnatal patients. There are 10 beds in the labour room which caters for approximately 9,000 deliveries per year. Two beds are used for the management of pre-eclampsia cases. With the introduction of the 'husband friendly' policy in February 1999, three beds are reserved for patients whose husbands wish to accompany them throughout the labour.

There are 36 gynaecology beds and we performed over 1,000 gynaecological operations per year covering a wide variety of cases as shown in the statistics. There are two operative sessions per week for elective cases. The facility for Caesarean Sections is

shared between the general operating theatre and the maternity operating theatre which is located in a block adjacent to the labour ward.

We run a weekly antenatal clinic, gynaecological clinic and subspecialty clinics like combined clinic (every two weeks), infertility clinic, colposcopy clinic and clinic seeing oncology and molar pregnancy cases.

The department is actively involved in audit and quality control assessment. It also has an ongoing commitment to both undergraduate and postgraduate education. The hospital is associated with the Faculty of Medicine, International Islamic University (Universiti Islam Antarabangsa, UIA) and undertakes undergraduate teaching.

The trainee post in the Department of Obstetrics and Gynaecology is recognized by the Health Ministry of Malaysia and local universities for training for the specialist qualification (Master of Medicine, Obstetrics & Gynaecology, M.MED O&G). The trainee post is also recognized for training by the Royal College of Obstetricians and Gynaecologists.

The trainees are resident in the hospital when on duty. With an average number of eight calls a month, they receive adequate experience in the management of emergency cases. All patients attended by trainees are also under the clinical supervision of practitioners of full specialist status.

Hospital Tengku Ampuan Afzan has an established education programme, fully equipped Postgraduate Medical Centre with good library facilities. The Department of Obstetrics and Gynaecology holds regular teaching programmes e.g. weekly teaching sessions for the trainees and journal clubs, regular joint obstetrics/neonatal paediatric meetings and clinico-pathological conferences.

The trainees participate in running the family planning clinics and receive instruction in neonatal resuscitation which is conducted by the Paediatric Department regularly. They are also involved in clinical research projects and in the training of medical students, nurses and midwives.

### The timetable for the work schedule is as follows:

	AM	PM
MONDAY	ANTENATAL	TEACHING SESSION
	CLINIC	FOR TRAINEES
TUESDAY	GYNAECOLOGICAL	HOUSE OFFICER
	CLINIC	TEACHING
WEDNESDAY	ELECTIVE	CAESAREAN SECTION
	OPERATIONS	AUDIT (every two weeks)
THURSDAY	SPECIAL	PERINATAL MORBIDITY
	CLINIC	AND MORTALITY
		MEETING (every two weeks)
FRIDAY	ELECTIVE	ELECTIVE
	OPERATIONS	OPERATIONS
SATURDAY	HOSPITAL CME	·
	(Continuing Medical	
	Education)	

# Statistics for the year 1999

#### **Obstetric Unit**

Total number of deliveries	8647	
Mode of deliveries	•	
Spontaneous vertex	7123	
Breech	181	
Forceps	57	
Vacuum	81	
Caesarean Sections	1205	
Total	8647	
Multiple Pregnancy		
Twins	65	
Triplets	1	
Total	66	

# **Gynaecological Unit**

Number of clinic attendance4235Number of admissions3325Number of operations1354

# **Gynaecological Operations (YEAR 1999)**

Type of operation	Number of cases
Total abdominal hysterectomy (with or without	102
oophorectomy)	
Vaginal hysterectomy (with or without pelvic floor repair)	14
Dilatation and curettage	671
Diagnostic laparoscopy	60
Sterilization: Mini-laparotomy	42
Laparoscopic	49
Cystectomy/oophorectomy	65
Myomectomy	21
Salpingectomy	107
Cervical cerclage	6
Wertheim's operation	7
Staging of cancer of cervix	12
Marsupialisation	21
Hysteroscopy	53
Others	124
Total	1354

#### **ABSTRAK**

Objektif Untuk menilai kebolehan ultrasound dan gambaran klinikal dalam membezakan ketulan ovari samada benign atau malignant.

Rekabentuk Satu kajian prospektif.

Tempat Jabatan Ginekologi, Hospital Tengku Ampuan Afzan, Kuantan, Pahang.

Subjek 126 orang pesakit wanita yang telah menjalani pembedahan untuk ketulan ovari dari 1 Februari 1999 hingga 31 Januari 2000

Ukuran pencapaian Sensitiviti, spesifisiti, kadar ramalan positif dan negatif ultrasound dan gambaran klinikal dalam membezakan ketulan ovari benign atau malignant. Keputusan 120 orang pesakit dimasukkan ke dalam kajian ini setelah mengeluarkan 6 pesakit yang tidak mempunyai patologi ovari. 97 pesakit (81%) telah mempunyai ketulan ovari benign manakala 23 pesakit (19%) mempunyai ketulan ovari malignant. Gambaran klinikal telah menunjukkan kadar sensitiviti dan spesifisiti yang tinggi, diikuti oleh skor ultrasound 2 hingga 5 dalam membezakan antara ketulan ovari benign atau malignant. Kedua-duanya mempunyai sensitiviti dan spesifisiti yang lebih dari 70% dan ianya lebih tinggi dari ultrasound.

Ujian klinikal dan ultrasound adalah kaedah yang Kesimpulan dalam dilakukan menilai boleh ketulan ovari sebelum pembedahan dilakukan. Ketepatan diagnostik boleh dipertingkatkan adanya dengan penanda tumour atau ultrasound yang lebih jitu. Tetapi ini adalah lebih sukar dan lebih mahal untuk dilakukan.

#### **ABSTRACT**

# CLINICAL IMPRESSION AND ULTRASOUND FEATURES IN THE PREOPERATIVE EVALUATION OF OVARIAN MASSES

Objective To evaluate the ability of clinical impression and ultrasound features to discriminate a benign from a malignant ovarian mass.

**Design** A prospective study.

Setting Department of Gynaecology, Hospital Tengku Ampuan
Afzan, Kuantan, Pahang.

Subjects 126 women admitted consecutively to the gynaecology department for surgical exploration of an ovarian mass between 1<sup>st</sup> February 1999 and 31<sup>st</sup> January 2000.

Main outcome measures The sensitivities, specificities, positive and negative predictive values of ultrasound features and clinical impression for discriminating a benign from a malignant ovarian mass.

Results 120 constituted the study population after excluding 6 cases of non-ovarian pelvic pathology. 97 cases (81%) were diagnosed as benign ovarian tumours and 23 cases (19%) as malignant. Clinical impression gave the highest level of sensitivity and specificity, followed by ultrasound score of 2-5 in the discrimination between benign and malignant ovarian masses. Both achieved sensitivity and specificity of greater than 70% and showed superiority over the individual ultrasound features suggestive of malignancy.

Conclusions Clinical examination and ultrasound scanning are methods which can be applied easily in our centre in the preoperative evaluation of ovarian masses. The diagnostic accuracy may be improved by including tumour markers or more detailed ultrasound assessment, this would be at the expense of simplicity and increased financial cost.

# CLINICAL IMPRESSION AND ULTRASOUND FEATURES IN THE PREOPERATIVE EVALUATION OF OVARIAN MASSES

#### by

#### DR CHEONG YEE TSING

Dissertation Submitted In
Partial Fulfilment Of The
Requirements For The Degree Of
Master Of Medicine
(Obstetrics and Gynaecology)

UNIVERSITI SAINS MALAYSIA NOVEMBER 2001

# **TABLE OF CONTENTS**

	Page
ACKNOWLEDGEMENTS	iii
1. INTRODUCTION TO:	
1.1 THE STATE OF PAHANG	iv
1.2 HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAH	HANG vi
1.3 DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY	Υ,
HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAH	HANG vii
2. ABSTRAK	xiii
3. ABSTRACT	xv
4. INTRODUCTION	1
5. MATERIALS AND METHODS	23
6. RESULTS	32
7. DISCUSSION	50
8. CONCLUSION .	77
9. LIMITATIONS	79
10. RECOMMENDATIONS	80
11. REFERENCES	81
12. APPENDIX	96

### **ACKNOWLEDGEMENTS**

I would like to acknowledge the invaluable guidance given to me by Dr Ghazali Ismail, Consultant Obstetrician and Gynaecologist, Head of the Department of Obstetrics and Gynaecology, Hospital Tengku Ampuan Afzan, Kuantan and Dr Mokhtar Awang, Consultant Obstetrician and Gynaecologist, Hospital Tengku Ampuan Afzan, Kuantan throughout the preparation for this dissertation and my posting in Kuantan.

I would also like to express my appreciation of the effort of Dr Awang Nila Ismail, Head and Consultant, Department of Obstetrics and Gynaecology, Hospital Universiti Sains Malaysia who painstakingly checked and corrected this dissertation. This book would not have been possible without their help.

I am also much indebted to everyone who has helped me — medical colleagues who have made helpful suggestions. I wish to thank each and every one of them for their efforts, and last but not least my parents for their love and encouragement.

#### INTRODUCTION TO THE STATE OF PAHANG

Pahang is the largest state in Peninsular Malaysia and is situated in the eastern coastal region. The state's 35,964 square kilometres encompasses a remarkable range of Malaysia's many different environments from the majestic peaks and cool hill regions of the state's western region to the miles of soft sand beach along the South China Sea. It is home to Taman Negara (National Park), all the major hill resorts in the country, and has Gunung Tahan, the highest peak in the peninsula.

This is a state that has a rich cultural tradition, of an ancient history. Stone artifacts point to man having existed here since the Stone Age. Later Indian empires, the Sri Vijaya and the Majapahit, spread their rule, customs and traditions over the people. Then came the Siamese and later the Melaka Sultante annexed the state. Others were to follow until finally the British brought the state into its ambit with the creation of the Federated Malay States. Today, the one million population constitutes a mixture of Malays, Chinese, Indians and aborigines, the Orang Asli.

The state capital, Kuantan, is really your first stop to a fascinating holiday. It is a contemporary town, growing fast yet retaining its ancient charms. About 48 km from Kuantan is Sungai Lembing, an old steel and tin mining town, where the deepest lode mine in the world was once very active. Relics of European architecture can still be found in the town.

All this has left Pahang with a heritage of colour, pomp and pageantry. Along with it the state inherited a beautiful coastline that is dotted with picturesque beaches and thousands of villages. The state is now one of the leading tourist destinations in the country.

Pahang provides the visitor with almost unspoilt natural splendours. There is myriad of exotic flora and fauna to bewilder you. There are pristine waterfalls along with palm-fringed sandy beaches. The state has an air of unspoilt beauty, a tropical holiday, an unforgettable journey that spells adventure.

# HOSPITAL TENGKU AMPUAN AFZAN, KUANTAN, PAHANG

Hospital Tengku Ampuan Afzan (HTAA) started its service in the year 1904. Over the years, it has undergone several stages of development and is now an eight-storey complex which is well equipped with medical facilities.

HTAA is the main hospital for the State of Pahang which is one of the thirteen component states of Malaysia. The hospital consists of 700 official beds and around 1,200 staff. It serves the town of Kuantan which is the capital town of the State of Pahang. It also serves as the referral centre for the seven outlying district hospitals, outpatient clinics and the private medical practitioners.

Since 1999, HTAA has become the teaching hospital for the Faculty of Medicine, International Islamic University (Universiti Islam Antarabangsa, UIA) and undertakes undergraduate teaching.

HTAA is recognized by the Health Ministry of Malaysia and local universities for the training of postgraduate students in the field of Obstetrics and Gynaecology, Surgery, Medicine, Paediatrics, Orthopaedics, Otorhinolaryngology and Family Medicine.

# Department of Obstetrics and Gynaecology Hospital Tengku Ampuan Afzan, Kuantan, Pahang

The Department of Obstetrics and Gynaecology is staffed by five Consultants/Specialists, six postgraduate students/registrars in training, five medical officers and a minimum of ten house officers.

The department has 36 antenatal beds and 44 postnatal beds. New cases are seen in the screening room after which they will either be admitted to the appropriate ward or observed in the screening room where we also provide day care services. The first class ward has 12 beds for antenatal and postnatal patients. There are 10 beds in the labour room which caters for approximately 9,000 deliveries per year. Two beds are used for the management of pre-eclampsia cases. With the introduction of the 'husband friendly' policy in February 1999, three beds are reserved for patients whose husbands wish to accompany them throughout the labour.

There are 36 gynaecology beds and we performed over 1,000 gynaecological operations per year covering a wide variety of cases as shown in the statistics. There are two operative sessions per week for elective cases. The facility for Caesarean Sections is

shared between the general operating theatre and the maternity operating theatre which is located in a block adjacent to the labour ward.

We run a weekly antenatal clinic, gynaecological clinic and subspecialty clinics like combined clinic (every two weeks), infertility clinic, colposcopy clinic and clinic seeing oncology and molar pregnancy cases.

The department is actively involved in audit and quality control assessment. It also has an ongoing commitment to both undergraduate and postgraduate education. The hospital is associated with the Faculty of Medicine, International Islamic University (Universiti Islam Antarabangsa, UIA) and undertakes undergraduate teaching.

The trainee post in the Department of Obstetrics and Gynaecology is recognized by the Health Ministry of Malaysia and local universities for training for the specialist qualification (Master of Medicine, Obstetrics & Gynaecology, M.MED O&G). The trainee post is also recognized for training by the Royal College of Obstetricians and Gynaecologists.

The trainees are resident in the hospital when on duty. With an average number of eight calls a month, they receive adequate experience in the management of emergency cases. All patients attended by trainees are also under the clinical supervision of practitioners of full specialist status.

Hospital Tengku Ampuan Afzan has an established education programme, fully equipped Postgraduate Medical Centre with good library facilities. The Department of Obstetrics and Gynaecology holds regular teaching programmes e.g. weekly teaching sessions for the trainees and journal clubs, regular joint obstetrics/neonatal paediatric meetings and clinico-pathological conferences.

The trainees participate in running the family planning clinics and receive instruction in neonatal resuscitation which is conducted by the Paediatric Department regularly. They are also involved in clinical research projects and in the training of medical students, nurses and midwives.

### The timetable for the work schedule is as follows:

	AM	PM
MONDAY	ANTENATAL	TEACHING SESSION
	CLINIC	FOR TRAINEES
TUESDAY	GYNAECOLOGICAL	HOUSE OFFICER
	CLINIC	TEACHING
WEDNESDAY	ELECTIVE	CAESAREAN SECTION
	OPERATIONS	AUDIT (every two weeks)
THURSDAY	SPECIAL	PERINATAL MORBIDITY
	CLINIC	AND MORTALITY
		MEETING (every two weeks)
FRIDAY	ELECTIVE	ELECTIVE
	OPERATIONS	OPERATIONS
SATURDAY	HOSPITAL CME	·
	(Continuing Medical	
	Education)	

# Statistics for the year 1999

#### **Obstetric Unit**

Total number of deliveries	8647	
Mode of deliveries		
Spontaneous vertex	7123	
Breech	181	
Forceps	57	
Vacuum	81	
Caesarean Sections	1205	
Total	8647	
Multiple Pregnancy		
Twins	65	
Triplets	1	
Total	66	

# **Gynaecological Unit**

Number of clinic attendance4235Number of admissions3325Number of operations1354

# **Gynaecological Operations (YEAR 1999)**

Type of operation	Number of cases
Total abdominal hysterectomy (with or without	102
oophorectomy)	
Vaginal hysterectomy (with or without pelvic floor repair)	14
Dilatation and curettage	671
Diagnostic laparoscopy	60
Sterilization: Mini-laparotomy	42
Laparoscopic	49
Cystectomy/oophorectomy	65
Myomectomy	21
Salpingectomy	107
Cervical cerclage	6
Wertheim's operation	7
Staging of cancer of cervix	12
Marsupialisation	21
Hysteroscopy	53
Others	124
Total	1354

#### **ABSTRAK**

Objektif Untuk menilai kebolehan ultrasound dan gambaran klinikal dalam membezakan ketulan ovari samada benign atau malignant.

Rekabentuk Satu kajian prospektif.

Tempat Jabatan Ginekologi, Hospital Tengku Ampuan Afzan, Kuantan, Pahang.

Subjek 126 orang pesakit wanita yang telah menjalani pembedahan untuk ketulan ovari dari 1 Februari 1999 hingga 31 Januari 2000

Ukuran pencapaian Sensitiviti, spesifisiti, kadar ramalan positif dan negatif ultrasound dan gambaran klinikal dalam membezakan ketulan ovari benign atau malignant. Keputusan 120 orang pesakit dimasukkan ke dalam kajian ini setelah mengeluarkan 6 pesakit yang tidak mempunyai patologi ovari. 97 pesakit (81%) telah mempunyai ketulan ovari benign manakala 23 pesakit (19%) mempunyai ketulan ovari malignant. Gambaran klinikal telah menunjukkan kadar sensitiviti dan spesifisiti yang tinggi, diikuti oleh skor ultrasound 2 hingga 5 dalam membezakan antara ketulan ovari benign atau malignant. Kedua-duanya mempunyai sensitiviti dan spesifisiti yang lebih dari 70% dan ianya lebih tinggi dari ultrasound.

Ujian klinikal dan ultrasound adalah kaedah yang Kesimpulan dalam dilakukan menilai boleh ketulan ovari sebelum pembedahan dilakukan. Ketepatan diagnostik boleh dipertingkatkan adanya dengan penanda tumour atau ultrasound yang lebih jitu. Tetapi ini adalah lebih sukar dan lebih mahal untuk dilakukan.

#### **ABSTRACT**

# CLINICAL IMPRESSION AND ULTRASOUND FEATURES IN THE PREOPERATIVE EVALUATION OF OVARIAN MASSES

Objective To evaluate the ability of clinical impression and ultrasound features to discriminate a benign from a malignant ovarian mass.

**Design** A prospective study.

Setting Department of Gynaecology, Hospital Tengku Ampuan
Afzan, Kuantan, Pahang.

Subjects 126 women admitted consecutively to the gynaecology department for surgical exploration of an ovarian mass between 1<sup>st</sup> February 1999 and 31<sup>st</sup> January 2000.

Main outcome measures The sensitivities, specificities, positive and negative predictive values of ultrasound features and clinical impression for discriminating a benign from a malignant ovarian mass.

Results 120 constituted the study population after excluding 6 cases of non-ovarian pelvic pathology. 97 cases (81%) were diagnosed as benign ovarian tumours and 23 cases (19%) as malignant. Clinical impression gave the highest level of sensitivity and specificity, followed by ultrasound score of 2-5 in the discrimination between benign and malignant ovarian masses. Both achieved sensitivity and specificity of greater than 70% and showed superiority over the individual ultrasound features suggestive of malignancy.

Conclusions Clinical examination and ultrasound scanning are methods which can be applied easily in our centre in the preoperative evaluation of ovarian masses. The diagnostic accuracy may be improved by including tumour markers or more detailed ultrasound assessment, this would be at the expense of simplicity and increased financial cost.

#### INTRODUCTION

The management of a woman found to have an adnexal mass on bimanual pelvic examination is a problem commonly faced by gynaecologists. The differential diagnosis is wide and varied and includes both benign and malignant gynaecological diseases in addition to pathologies affecting the bowel and urinary tract. Among these different benign and malignant conditions, the definitive diagnosis of an ovarian mass is a common problem in gynaecologic patients.

An adnexal mass often involves ovarian substance because of the propensity of the ovary for neoplasia. Although the overwhelming majority of adnexal masses are benign, it is important to determine preoperatively whether a patient is at a high risk for ovarian malignancy so that the appropriate preparation and intervention can be planned.

Ninety percent of all ovarian tumours are benign, although this varies with age. Among surgically managed cases, the frequency of malignant tumours is 13% in premenopausal women and 45% in postmenopausal women. The lifetime risk of developing ovarian cancer is 1.4% which is higher than that for either cancer of the cervix (1.25%) or the endometrium (1.1%). Deaths from ovarian

#### INTRODUCTION

The management of a woman found to have an adnexal mass on bimanual pelvic examination is a problem commonly faced by gynaecologists. The differential diagnosis is wide and varied and includes both benign and malignant gynaecological diseases in addition to pathologies affecting the bowel and urinary tract. Among these different benign and malignant conditions, the definitive diagnosis of an ovarian mass is a common problem in gynaecologic patients.

An adnexal mass often involves ovarian substance because of the propensity of the ovary for neoplasia. Although the overwhelming majority of adnexal masses are benign, it is important to determine preoperatively whether a patient is at a high risk for ovarian malignancy so that the appropriate preparation and intervention can be planned.

Ninety percent of all ovarian tumours are benign, although this varies with age. Among surgically managed cases, the frequency of malignant tumours is 13% in premenopausal women and 45% in postmenopausal women. The lifetime risk of developing ovarian cancer is 1.4% which is higher than that for either cancer of the cervix (1.25%) or the endometrium (1.1%). Deaths from ovarian

cancer outnumber those from carcinoma of the cervix and body of the uterus combined (Anderson et al. 1997).

The presumptive diagnosis of an ovarian mass will often determine the type of surgery and the preoperative preparation required, and may influence selection of the institution and the seniority and expertise of the surgeon (Davies et al. 1993). The prognosis for women with ovarian cancer may be greatly influenced by appropriate first line surgery (Oram & Jacobs 1987; Gillis et al. 1991). The initial laparotomy is not only important for the accurate determination of the extent of disease, but also the best opportunity for maximum debulking (Hacker 1987).

If the clinician could accurately differentiate a malignant from a benign mass, patients with masses believed to be malignant could seek proper oncologic consultation preoperatively. At the time of operation, the appropriate incision would allow careful staging. The preoperative diagnosis of ovarian cancer might reduce the need for repeat laparotomies and could lead to referral to or consultation with gynaecologic oncologists capable of proceeding with radical procedures. This surgery demands specific skills and experience. The amount of residual malignant tissue after primary surgery is among the best prognostic factors in ovarian cancer (Levin et al.

1993). Therefore, accurate preoperative prediction of the benign or malignant character of an ovarian tumour is essential for optimal preoperative and intraoperative surgical management, which may provide, in the case of malignancy, the best chance of a long disease free interval and cure (Schutter et al. 1994).

Minimal access surgery (MAS) has become an important part of the gynaecologist's repertoire, especially when used for removing a persistent or symptomatic benign ovarian cyst (Mettler et al. 1993; Parker & Berek 1993). However, if there is a suspicion of malignancy, most gynaecologists would perform a laparotomy via a midline incision. The incidence of finding a malignant cyst at MAS in women where preoperative investigations wrongly suggested that the mass was benign ranges from 0.04% to 3.7% (Peterson et al. 1990; Hulka et al. 1992; Nezhat et al. 1992; Canis et al. 1994). The role of MAS in early ovarian cancer has not been clearly defined. It may be detrimental to survival of the patient if the laparoscopic management results in inadequate staging, the piecemeal removal of a potentially curable early stage cancer, or suboptimal clearance of tumour in more advanced disease (Crawford et al. 1995).

In spite of the known benefits of thorough surgical staging and cytoreductive surgery, many women do not receive appropriate

treatment at the time of the initial surgical diagnosis (Young et al. 1983). This is partly because of the skilled and lengthy surgery that radical cytoreduction often requires and partly a consequence of the difficulty in making an accurate preoperative diagnosis of ovarian cancer. It is postulated that methods of improving the preoperative diagnosis of ovarian malignancy would lead to more women receiving first line surgery from appropriately trained and experienced personnel, either locally or after referral to a gynaecologist with a special interest in gynaecological oncology (Davies et al. 1993).

An accurate preoperative diagnosis of ovarian cancer and a method for better preoperative discrimination of patients with an ovarian mass are therefore of great value and much needed.

The standard evaluation for adnexal masses includes history, physical examination and ultrasound. In an international multicenter study conducted by Schutter et al. (1994), the individual accuracy of pelvic examination, ultrasound and serum CA 125 in discriminating between benign and malignant pelvic masses among postmenopausal patients was approximately the same (76, 74 and 77%, respectively). Buist et al. (1992) assessed the diagnostic potential of pelvic examination, the study showed that

the test accuracy of pelvic examination (70%) was even better than that for computed tomography and ultrasonography (both 65%) and magnetic resonance imaging (50%).

Ultrasound examination of the pelvis and abdomen has become the standard diagnostic test for evaluating adnexal masses and is by far the most commonly used imaging modality for evaluation of a patient suspected of having ovarian malignancy. Its principal value in this setting involves confirming the mass, differentiating ovarian from uterine or tubal origin, delineating the internal appearance of the mass, and defining any associated abdominal findings. It is being explored as a screening method for carcinoma of the ovary, in the attempt to detect early disease. The overall rate of false positive was 9.3% and the likelihood of a positive scan being a primary ovarian cancer was 1:67 (Campbell et al. 1989).

Whether ultrasound can differentiate between benign and malignant pelvic masses has been the subject of many studies. Scoring systems have been evolved to reduce the number of false positives. One of the scoring systems used (Davies et al. 1993) where ultrasound scans were performed and assessed for the following features suggestive of malignancy: multiloculated cysts; evidence of solid areas; bilateral lesions; presence of ascites and

evidence of metastases. An ultrasound score of 3 (when two or more of the above features were noted) achieved a sensitivity of 87%, specificity of 74% and positive predictive value of 58%.

The serum CA 125 test is a helpful expedient in the diagnosis and monitoring of epithelial ovarian carcinoma. Einhorn et al. (1986) have suggested that preoperative serum CA 125 levels may be of value in differentiating a benign from a malignant pelvic mass. However, CA 125 as a single parameter does not distinguish sufficiently benign from malignant masses preoperatively, because it can be elevated in menstruating women, during pregnancy, by liver cirrhosis, and in heart failure (Niloff et al. 1984). Raised serum CA 125 levels are also found in association with benign ovarian cysts, endometriosis and pelvic infection.

Finkler et al. (1988) assessed the diagnostic procedures: serum CA 125, clinical examination and ultrasound interpretation and they found that in conjunction with such tests, measurement of serum CA 125 significantly increased diagnostic accuracy and may thus have an important role in the preoperative evaluation of women with ovarian masses.

Due to the limited budget in our centre, tumour markers are not routinely done to all patients presented with an adnexal mass. However, evaluation of the patient using clinical judgement and ultrasound scan can be applied easily in our centre.

The purpose of this study is to evaluate the ability of ultrasound findings and clinical impression to discriminate a benign from a malignant mass in patients presented to the gynaecology department with an ovarian mass. Sensitivity, specificity, positive and negative predictive values were calculated for tests used in patients undergoing surgical exploration of ovarian masses.

# Literature Review Of The Ability Of Other Methods To Discriminate A Benign From A Malignant Ovarian Mass

Trans-vaginal ultrasonography, Doppler ultrasound and Serum CA 125 have all been shown to be important diagnostic tools in the preoperative evaluation of ovarian masses. However, due to the limited budget and unavailability of the facility in our center, the use of these tests are not possible in all patients presented with an ovarian mass.

# Trans-vaginal ultrasonography

Since the mid-1980s, with technical advances in instrumentation and imaging, trans-vaginal ultrasonography has gained in popularity. The advantages of this approach are two-fold:

- 1. The ultrasound transducer is closer to the objects of interest and not separated from them by layers of fat and muscle. The pelvic organs can therefore be studied with higher ultrasound frequencies, producing images of greater resolution and enhanced quality.
- 2. The need for a full bladder is obviated.

Although trans-vaginal ultrasonography has better resolution than abdominal ultrasonography and provides detailed images of the ovaries and masses confined to the true pelvis, there are some disadvantages to trans-vaginal ultrasonography. It is more invasive and postmenopausal vaginal atrophy may limit access and patient acceptability.

The ultrasound frequencies used improve resolution, but decrease penetration to 10cm. Thus ovaries situated high or lateral in the pelvis, or greatly enlarged so that they are no longer in the pelvis, may not be seen on scan. Co-incident abdominal palpation or bimanual pelvic examination (or a trans-abdominal scan) should ensure that such ovaries are detected (Timor-Tritsh et al. 1988).

With the advent of high-frequency trans-vaginal ultrasonography, new opportunities are presented to better define ovarian lesions. Scoring system using trans-vaginal sonographic characterization of pelvic/ovarian lesions has been developed to maximize the discrimination between benign and malignant entities. Test results have improved using trans-vaginal ultrasound examination compared with the transabdominal method (Sassone et al. 1991).

# Doppler

Whilst ultrasound imaging has provided a non-invasive method of studying ovarian morphology, pulsed Doppler combined with real-time ultrasound (the duplex method) has the potential to examine patterns of blood flow and hence of identifying functional changes. The duplex technique will show a low pulsatility index (PI) in arteries with vascular impedance.

Colour combined with pulse Doppler has been used to investigate the pelvic circulation under various physiological and pathological conditions (Kurjak et al. 1989). Changes in vascularity (which may be an indication of malignancy) can be identified as fluctuating areas of colour. Applying pulsed Doppler to these areas gives a measurement of blood flow and impedance. Previous studies have malignancy associated may be with shown that ovarian characteristic patterns of flow in the newly formed vessels within the tumour (Bourne et al. 1989; Kurjak et al. 1989, 1991).

It has recently become possible to obtain colour Doppler signals using a trans-vaginal probe. This technique not only assesses morphological features of malignancy, but also functional changes that may be associated with malignancy such as the presence of neovascularisation. Two studies have shown that ovarian cancers

may be distinguished from benign adenomas using this technique (Kurjak et al. 1989; Bourne et al. 1989).

With the introduction of trans-vaginal colour Doppler ultrasound in gynaecology, it is now possible for us to evaluate the haemodynamics in benign and malignant tumours. Earlier reports suggested that pulsatility index (Bourne et al. 1989) or resistance index (Kurjak et al. 1990) might represent independent parameters allowing the differentiation of benign from malignant tumours. More recent evidence from the analysis of larger populations suggests that neither pulsatility index nor resistance index are sufficient parameters for the preoperative classification of tumours, if taken singularly. Kurjak et al. (1993) highlighted the importance of a multiparameter evaluation including the number and location of vessels.

Most malignancies are characterized by a central location of vessels. These malignant tumours tend to initiate tumour neovascularity in the centre of the lesion whereas peripherally located vessels probably originate from pre-existing host vessels. The detection of centrally located vessels is therefore highly predictive of malignancy. New scoring systems should therefore include parameters such as conventional morphology, the location

of vessels, their arrangement and haemodynamics (Kurjak et al. 1993).

The techniques of pulsed and colour Doppler play an increasingly important role in the differential diagnosis of a pelvic mass and will probably serve to improve the specificity of ultrasound for ovarian cancer (Davies & Oram 1991). Although the accuracy of differentiation between ovarian carcinoma and benign pelvic masses might be further improved by the technique of Doppler ultrasound with colour flow imaging (Bourne et al. 1989; Kawai et al. 1992; Weiner et al. 1992), this technique requires expensive technology and thus will not be immediately available to most of the centres.

## Tumour markers

Quantitative and qualitative changes in numerous circulating substances have been associated with epithelial ovarian cancer. This may reflect an alteration in ovarian function or surface molecular structure, or a 'general' response to malignancy (Smith & OI 1984b, Bast & Knapp 1987).

Changes in circulating enzymes (van Kley et al. 1981; Awais 1978; Gauduchon et al. 1983; Cramer et al. 1989), hormones (Heinonen et al. 1982; Backstrom et al. 1983), non-specific inflammatory proteins (Lukomska et al. 1981), placental and fetal antigens (Donaldson et al. 1980; Nouwen et al. 1985; Doellgast & Homesley 1984) have all been identified in women with ovarian cancer. Examples of the serum markers for ovarian cancer are as follows:

## **Enzymes**

- 1. Galactosyl transferase
- 2. Alpha 1-fucosidase
- 3. Amylase
- 4. Lactic dehydrogenase
- 5. Cystine aminopeptidase

# Feto-placental markers

- 1. Alpha-feto protein
- 2. Human chorionic gonadotrophin
- 3. Placental alkaline phosphatase
- 4. Carcinoembryonic antigen

#### Hormones

- 1. Progesterone
- 2. Oestrogen

### Miscellaneous

- 1. Circulating immune complexes
- 2. d-dimer of fibrin

Using monoclonal antibody techniques, antigen with improved specificity and sensitivity for epithelial ovarian cancer has been defined – though the ultimate tumour-specific antigen remains elusive. The tumour marker shown to combine the highest sensitivity and specificity for the disease has been CA 125 which is also the most extensively studied ovarian tumour-associated antigen (Bast et al. 1983).

## **CA 125**

CA 125, a high molecular-weight glycoprotein, is recognized by a murine monoclonal antibody, OC 125. Immunohistochemical studies have demonstrated CA 125 expression to be a feature of cells derived from the embryonal coelomic epithelium and Mullerian duct (Kabawat *et al.* 1983) and both benign and malignant pathologies affecting tissues with these embryological origins have been associated with increased CA 125 expression (Niloff *et al.* 1984).

Serum CA 125 levels measured by radioimmunoassay are greater than 35U/ml in over 80% of women with epithelial ovarian cancer (especially of the non-mucinous type) (Bast et al. 1983; Hawkins et al. 1989). Bast et al. (1983) have also shown that only 1% of healthy female blood donors have serum CA 125 levels of greater than 35U/ml. However, raised serum CA 125 levels are also found

in association with benign ovarian cysts, endometriosis and pelvic infection in addition to cancers of the endometrium, fallopian tube, breast and colon. Consequently, the accuracy of CA 125 measurement in differentiating benign from malignant pelvic pathology is limited (Niloff et al. 1984).

In view of the association of CA 125 with pathologies other than ovarian cancer and therefore a low specificity and low positive predictive value, methods of improving its specificity are being investigated.

# CA 125. clinical examination and ultrasound

In the study by Finkler et al. (1988), the ability to differentiate a malignant from a benign ovarian mass was assessed for the following diagnostic procedures: serum CA 125, clinical examination and ultrasound interpretation. Results of diagnostic tests in combination were most informative. CA 125 added predictive value to each of the other diagnostic tests used in this study.

The study conducted by Tingulstad et al. (1996) showed superiority of CA 125 over ultrasound and menopausal status in their ability to discriminate a benign from a malignant pelvic mass. At cut-off level

of 70U/ml, CA 125 has a sensitivity of 70%, specificity of 95% and positive predictive value of 87% in predicting malignancy. Results from other studies also showed that serum CA 125 estimation improved the accuracy of diagnosing ovarian cancer preoperatively when combined with pelvic ultrasonography (Gadducci et al. 1988) or menopausal status (Vasilev et al. 1988).

## CA 125 and other tumour markers

Combinations of CA 125 with other tumour markers for discriminating between benign and malignant diseases have also been used (Gadducci et al. 1991).

The use of a panel of monoclonal antibodies directed against a spectrum of tumour-associated antigens is being reviewed as a method of improving the specificity of serum CA 125 for ovarian cancer. These antigens may be different epitopes on the same complex or totally separate antigens.

At Duke University Medical Centre, United States, Bast et al. (1990) looked for CA 125, CA 15-3, TAG 72, PLAP, HMFG 1, HMFG 2 and NB/70K expression in serum from patients with known ovarian cancer and from women with a 'false positive' elevation of CA 125. His findings indicate CA 15-3 and TAG 72 to be the more

useful antigens for differentiating benign from malignant disease. Serum levels of all three antigens were elevated in 77% of women with ovarian cancer compared with only 5% of women with benign disease. The improved specificity for ovarian cancer using the combination of serum CA 125, CA 15-3 and TAG 72 has been confirmed by others. Einhorn *et al.* (1989) found 98% specificity for malignancy when serum levels of all three antigens were raised, compared with 83% for CA 125 used as a single test.

The increase in specificity using such a method may however occur at the expense of sensitivity. A fall in sensitivity for ovarian cancer to 53% has been documented using a combination of serum CA 125, CA 15-3 and TAG 72 (Jacobs et al. 1990).

A general conclusion of several studies (Soper et al. 1990) is that some additional information can be obtained from combinations of serum tumour markers but that other diagnostic modalities are needed for optimal and correct preoperative judgement of a pelvic mass.

There has been recent interest in developing scoring systems to improve the preoperative discrimination between benign and malignant ovarian masses (Davies et al. 1993). Most scoring

systems have included either tumour markers (Cruickshank et al. 1987; Vasilev et al. 1988; Einhorn et al. 1989; Bast et al. 1990) or ultrasonography (Deland et al. 1979; Hermann et al. 1987).

A study by Jacobs et al. (1990) of 143 women undergoing surgery for a pelvic mass assessed age, ultrasound score, menopausal status, a clinical impression score and the serum CA 125 level as discriminators between benign and malignant ovarian disease. The most useful individual criteria were the serum CA 125 level and the ultrasound score.

In order to improve the accuracy in discriminating benign from malignant ovarian disease, Jacobs et al. (1990) defined a risk of malignancy index (termed RMI). The RMI being the product of the serum CA 125 level (U/mI), the ultrasound score (U) and the score for menopausal status (M):

 $RMI = U \times M \times serum CA 125$ 

Postmenopausal status was defined as more than one year of amenorrhoea or age older than 50 years in women who had had a hysterectomy. Women who did not meet these criteria were classified as premenopausal. The ultrasound examination was performed via the abdominal route with the full bladder technique.

The presence of a multilocular cystic lesion, solid areas, bilateral lesions, ascites and intra-abdominal metastases, scored 1 point each. A total ultrasound score (U score) was calculated for each patient.

Based on the data obtained, the RMI was calculated for each patient. The calculation was based on a simplified regression equation where total ultrasound score of 0 gave U=0, score 1 U=1, and score  $\geq 2$  U=3; premenopausal status gave M=1, postmenopausal M=3. The serum level of CA 125 was applied directly into the calculation.

A RMI of 200 achieved a high sensitivity of 85% and a specificity of 97% for diagnosing ovarian cancer. Davies et al. (1993) undertook a study to verify the accuracy of RMI in discriminating benign from malignant ovarian disease. Tested on a new population of women, the RMI retained the high sensitivity for diagnosing ovarian cancer seen in the original report describing its derivation. The study confirmed that the RMI is more accurate than the individual criteria in diagnosing ovarian cancer, and was comparable with other scoring systems.

The main advantage of the RMI, compared with other approaches such as blood flow characterization with Doppler ultrasound (Bourne et al. 1989; Kurjack et al. 1989; Fleischer et al. 1991; Kurjack et al. 1991; Weiner et al. 1992) or the use of a panel of different tumour markers (Einhorn et al. 1986; Vasilev et al. 1988; Zi-Xia et al. 1988; Gadducci et al. 1992; McGuckin et al. 1992; Jacobs et al. 1993), is that the RMI can be applied easily in less specialized departments of gynaecology.

## **MATERIALS AND METHODS**

Participants of the study were women admitted consecutively to the Department of Gynaecology, Hospital Tengku Ampuan Afzan, Kuantan, Pahang for surgical exploration of an ovarian mass between 1<sup>st</sup> February 1999 and 31<sup>st</sup> January 2000. Among those women who presented with a pelvic mass at the Department of Gynaecology, Hospital Tengku Ampuan Afzan, Kuantan, Pahang between 1<sup>st</sup> February 1999 and 31<sup>st</sup> January 2000, 126 women were diagnosed to have ovarian mass. The method of detection was either by physical examination or by ultrasonography. They were subsequently admitted for surgical exploration of the ovarian mass and these 126 consecutive patients were enrolled in the study.

A standardized data sheet was used to record clinical and test data (see Appendix). Data collected on each patient included age, parity, ethnic group, family history of ovarian cancer, menopausal status, usage of oral contraceptive pills, clinical impression, ultrasound interpretation and final pathologic diagnosis.

Family history of ovarian cancer was defined as one or more first degree relatives with ovarian cancer. Postmenopausal status was defined as women with menopausal symptoms and more than one year of amenorrhoea or age older than fifty years in women who had had a hysterectomy. Women who did not meet these criteria were classified as premenopausal.

The clinical impression of the ovarian mass was judged based on history and physical examination, both general and pelvic. The clinical assessment and physical examination were performed by the participating consultant gynaecologist. The examination was considered to be abnormal when a palpable mass of any size was found to be clinically distinguishable as being separate from the uterus and gastrointestinal tract. The participating clinician was specifically asked to define the ovarian mass as either benign or malignant without knowledge of the ultrasound interpretation. Report of clinical impression of the ovarian mass as either benign or malignant was then recorded preoperatively in all cases.

All women admitted for surgical exploration of ovarian masses had an ultrasound scan done preoperatively within one week before surgical exploration. All ultrasound scans were performed by consultant gynaecologists or under their supervision using a