

# Crystal structure of cyclohexylammonium thiocyanate

Abdulaziz A. Bagabas,<sup>a,†</sup> Sultan B. Alhoshan,<sup>a</sup> Hazem A. Ghabbour,<sup>b</sup> C. S. Chidan Kumar<sup>c,d,§</sup> and Hoong-Kun Fun<sup>b,c,\*¶</sup>

<sup>a</sup>National Petrochemical Technology Center (NPTC), Materials Science Research Institute (MSRI), King Abdulaziz City for Science and Technology (KACST), PO Box 6086, Riyadh 11442, Saudi Arabia, <sup>b</sup>Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, PO Box 2457, Riyadh 11451, Saudi Arabia, <sup>c</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and <sup>d</sup>Department of Chemistry, Alva's Institute of Engineering & Technology, Mijar, Moodbidri 574225, Karnataka, India. \*Correspondence e-mail: hfun.c@ksu.edu.sa

Received 6 December 2014; accepted 14 December 2014

Edited by J. Simpson, University of Otago, New Zealand


In the title salt,  $C_6H_{11}NH_3^+ \cdot SCN^-$ , the cyclohexylammonium ring adopts a slightly distorted chair conformation. The ammonium group occupies an equatorial position to minimize 1,3 and 1,5 diaxial interactions. In the crystal, the components are linked by  $N-H \cdots N$  and  $N-H \cdots S$  hydrogen-bonding interactions, resulting in a three-dimensional network.

**Keywords:** crystal structure; cyclohexylammonium; distorted chair; hydrogen bonding.

**CCDC reference:** 1039130

## 1. Related literature

For the synthesis and uses of the title compound, see: Baluja *et al.* (1960); Coddens *et al.* (1986); Goel (1988); Mathes *et al.* (1948) 1955); Mathes & Stewart (1955); Morrison & Ratcliffe (1953); Stewart (1951); For the structures of other cyclohexylammonium salts, see: Bagabas *et al.* (2014); Shimada *et al.* (1955); Smith *et al.* (1994); Odendal *et al.* (2010).



## 2. Experimental

### 2.1. Crystal data

$C_6H_{14}N^+ \cdot NCS^-$	$Z = 18$
$M_r = 158.26$	Cu $K\alpha$ radiation
Trigonal, $R\bar{3}$	$\mu = 2.71 \text{ mm}^{-1}$
$a = 23.4036 (6) \text{ \AA}$	$T = 296 \text{ K}$
$c = 8.3373 (2) \text{ \AA}$	$0.98 \times 0.25 \times 0.11 \text{ mm}$
$V = 3954.8 (2) \text{ \AA}^3$	

### 2.2. Data collection

Bruker APEXII CCD diffractometer	10355 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2009)	1670 independent reflections
$T_{\min} = 0.176$ , $T_{\max} = 0.748$	1530 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.051$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.100$	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
1670 reflections	
103 parameters	

**Table 1**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$N1-H1N \cdots S1^i$	0.89 (2)	2.62 (2)	3.4955 (14)	167 (2)
$N1-H2N \cdots N2$	0.90 (2)	1.94 (2)	2.822 (2)	170.4 (18)
$N1-H3N \cdots S1^{ii}$	0.87 (2)	2.573 (18)	3.4214 (14)	165.5 (19)

Symmetry codes: (i)  $x - y + \frac{2}{3}, x + \frac{1}{3}, -z + \frac{4}{3}$ ; (ii)  $-x + \frac{1}{3}, -y + \frac{2}{3}, -z + \frac{2}{3}$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL2013 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

## Acknowledgements

The authors extend their appreciation to King Abdulaziz City for Science and Technology (KACST) for funding this work through project number 29–280. CSCCK thanks Universiti Sains Malaysia (USM) for a postdoctoral research fellowship.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SJ5432).

† Additional correspondence author, e-mail: abagbas@kacst.edu.sa.

§ Thomson Reuters ResearcherID: C-3194-2011.

¶ Thomson Reuters ResearcherID: A-3561-2009.

## References

- Bagabas, A. A., Aboud, M. F. A., Shemsi, A. M., Addurihem, E. S., Al-Othman, Z. A., Chidan Kumar, C. S. & Fun, H.-K. (2014). *Acta Cryst.* **E70**, o253–o254.
- Baluja, G., Chase, B. H., Kenner, G. W. & Todd, A. (1960). *J. Chem. Soc.* pp. 4678–4681.
- Bruker (2009). *SADABS, APEX2 and SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Coddens, M. E., Furton, K. G. & Poole, C. F. (1986). *J. Chromatogr. A*, **356**, 59–77.
- Goel, A. B. (1988). US Patent 4 775 735.
- Mathes, R. A. & Stewart, F. D. (1955). US Patent 2 704 752.
- Mathes, R. A., Stewart, F. D. & Swedish, F. Jr (1948). *J. Am. Chem. Soc.* **70**, 3455.
- Morrison, A. L. & Ratcliffe, F. (1953). GB Patent 697 473.
- Odendal, J. A., Bruce, J. C., Koch, K. R. & Haynes, D. A. (2010). *CrystEngComm*, **12**, 2398–2408.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shimada, A., Okaya, Y. & Nakaura, M. (1955). *Acta Cryst.* **8**, 819–822.
- Smith, H. W., Mastropaolo, D., Camerman, A. & Camerman, N. (1994). *J. Chem. Crystallogr.* **24**, 239–242.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Stewart, F. D. (1951). US Patent 2 547 722.

## supporting information

*Acta Cryst.* (2015). E71, o62–o63 [https://doi.org/10.1107/S2056989014027297]

## Crystal structure of cyclohexylammonium thiocyanate

**Abdulaziz A. Bagabas, Sultan B. Alhoshan, Hazem A. Ghabbour, C. S. Chidan Kumar and Hoong-Kun Fun**

### S1. Chemical context

The title compound,  $C_6H_{11}NH_3^+ \cdot SCN^-$ , has previously been synthesized by reacting an aqueous solution of cyclohexylamine (CHA) with ammonium thiocyanate ( $H_4N^+ \cdot SCN^-$ ) at 85–90 °C, followed by extraction of  $C_6H_{11}NH_3^+ \cdot SCN^-$  with benzene, and then recrystallization from ethanol (Mathes *et al.*, 1948). This compound is used as an animal repellent and also as an insecticide or fungicide (Stewart, 1951). It is also a starting material for the preparation of other compounds (Baluja *et al.*, 1960; Morrison *et al.*, 1953; Stewart, 1951), an accelerator and activator for rubber vulcanization (Mathes *et al.*, 1955) and an accelerator for the curing of polyepoxide-polyimine materials (Goel, 1988). It has also been used as a stationary phase for gas chromatography (Coddens *et al.*, 1986). Nevertheless, the crystal structure of this important compound has not been determined. We report here the crystal structure of  $C_6H_{11}NH_3^+ \cdot SCN^-$  together with a new room-temperature synthesis using a salt metathesis reaction. This is a simpler process and results in a higher yield than the one published in the literature. Furthermore, we aim to use this compound to prepare new metal complexes based on both the cyclohexylammonium cation and the thiocyanate anion.

### S2. Structural commentary

The asymmetric unit of the title compound (Fig. 1), contains one cyclohexylammonium cation (C1–C6/N1) and one thiocyanate anion (S1/C7/N2). The cyclohexylammonium ring adopts a slightly distorted chair conformation, with puckering parameters:  $Q = 0.5669(18) \text{ \AA}$ ,  $\theta = 177.95(18)^\circ$ , and  $\varphi = 161(5)^\circ$ . For an ideal chair configuration,  $\theta$  has a value of 0 or 180°. The ammonium functional group is at an equatorial position to minimize 1,3 and 1,5 di-axial interactions. The bond lengths and bond angles are in normal ranges and are comparable with those reported earlier for similar compounds (Bagabas *et al.*, 2014; Shimada *et al.*, 1955; Smith *et al.*, 1994; Odendal *et al.*, 2010).

### S3. Supramolecular features

In the asymmetric unit, a strong N1–H2N $\cdots$ N2 hydrogen bond links the cation and the anion. In the crystal structure, these contacts are supported by intermolecular N1–H1N $\cdots$ S1 and N1–H3N $\cdots$ S1 hydrogen bonds (Symmetry codes:  $x-y+2/3, x+1/3, -z+4/3$ ;  $-x+1/3, -y+2/3, -z+2/3$ ) with S1 as a bifurcated acceptor (Fig. 2) to produce a three-dimensional network.


### S4. Synthesis and crystallization

The title compound,  $C_6H_{11}NH_3^+ \cdot SCN^-$ , was prepared by exchanging counter ions in a salt metathesis reaction between sodium thiocyanate (NaSCN) and cyclohexylammonium chloride ( $C_6H_{11}NH_3^+ \cdot Cl^-$ ) in ethanolic medium, where the precipitation of sodium chloride (NaCl) is the driving force for the reaction. In typical reaction, 100 mmol of NaSCN was dissolved in 350 ml absolute ethanol, while 100 mmol  $C_6H_{11}NH_3^+ \cdot Cl^-$  was dissolved separately in 250 ml absolute ethanol. Combining these two solutions at room temperature resulted in white precipitate of NaCl, as confirmed by X-ray powder

diffraction (PXRD), which was filtered off through F-size fritted filter. The filtrate was left for the solvent to evaporate to dryness at room temperature. About 200 ml absolute ethanol was added to the residual solid to dissolve the desired product,  $C_6H_{11}NH_3^+.SCN^-$ . It was noticed at this step that a small amount of the residual solid did not dissolve and it was separated by filtration. This undissolved material was NaCl, again identified by PXRD. Slow evaporation at room temperature over a period of 10 days of the ethanolic solution resulted in colorless crystals of  $C_6H_{11}NH_3^+.SCN^-$  (yield = around 100%) suitable for single crystal X-ray diffraction studies. It is advisable to recrystallize this light-sensitive material in the dark. The chemical composition of the desired product was also confirmed by C, H, N, S elemental microanalysis: (%C: 52.67 exp.; 53.11 cal.), (%H: 9.30 exp.; 8.93 cal.), (%N: 17.85 exp.; 17.70 cal.), and (%S: 20.18 exp.; 20.25 cal.). Potassium thiocyanate (KSCN) can be used instead of NaSCN for executing the reaction, where KCl precipitates and  $C_6H_{11}NH_3^+.SCN^-$  produces with around 98% yield.


### S5. Refinement details

The nitrogen-bound H-atoms were located in a difference Fourier map and were refined freely (NH = 0.87 (2), 0.90 (2) and 0.89 (2) Å). Other H atoms were positioned geometrically (C—H 0.97–0.98 Å) and refined using a riding model with  $U_{iso}(H) = 1.2 U_{eq}(C)$



**Figure 1**

The molecular structure of the title compound with atom labels and 50% probability displacement ellipsoids. The strong N—H...N hydrogen bond linking the cation and the anion is shown as a dashed line.



**Figure 2**

Crystal packing of the title compound, showing the N–H⋯N and N–H⋯S hydrogen bonding interactions (see Table 1) as dashed lines producing a three-dimensional network

### Cyclohexylammonium thiocyanate

#### Crystal data

$C_6H_{14}N^+ \cdot NCS^-$

$M_r = 158.26$

Trigonal,  $R\bar{3}:H$

$a = 23.4036(6) \text{ \AA}$

$c = 8.3373(2) \text{ \AA}$

$V = 3954.8(2) \text{ \AA}^3$

$Z = 18$

$F(000) = 1548$

$D_x = 1.196 \text{ Mg m}^{-3}$

Cu  $K\alpha$  radiation,  $\lambda = 1.54178 \text{ \AA}$

Cell parameters from 3689 reflections

$\theta = 3.8\text{--}71.3^\circ$

$\mu = 2.71 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Needle, colourless

$0.98 \times 0.25 \times 0.11 \text{ mm}$

#### Data collection

Bruker APEXII CCD  
diffractometer

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2009)

$T_{\min} = 0.176$ ,  $T_{\max} = 0.748$

10355 measured reflections

1670 independent reflections

1530 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$

$\theta_{\max} = 72.1^\circ$ ,  $\theta_{\min} = 3.8^\circ$

$h = -28 \rightarrow 28$

$k = -28 \rightarrow 28$

$l = -7 \rightarrow 9$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.038$  $wR(F^2) = 0.100$  $S = 1.06$ 

1670 reflections

103 parameters

0 restraints

Hydrogen site location: mixed

H atoms treated by a mixture of independent  
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0622P)^2 + 1.6831P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} < 0.001$  $\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.06708 (2)	0.40416 (2)	0.39763 (4)	0.04932 (18)
N1	0.22496 (6)	0.30544 (6)	0.62467 (16)	0.0396 (3)
N2	0.15453 (8)	0.36894 (8)	0.5327 (2)	0.0628 (4)
C1	0.17544 (6)	0.24495 (6)	0.71255 (15)	0.0339 (3)
H1A	0.1541	0.2582	0.7939	0.041*
C2	0.12316 (7)	0.19773 (7)	0.59728 (17)	0.0433 (3)
H2A	0.1437	0.1862	0.5119	0.052*
H2B	0.1005	0.2187	0.5496	0.052*
C3	0.07361 (7)	0.13543 (8)	0.6863 (2)	0.0525 (4)
H3A	0.0494	0.1465	0.7626	0.063*
H3B	0.0422	0.1041	0.6102	0.063*
C4	0.10721 (9)	0.10360 (7)	0.7747 (2)	0.0570 (4)
H4A	0.1264	0.0871	0.6973	0.068*
H4B	0.0746	0.0664	0.8364	0.068*
C5	0.16092 (8)	0.15210 (8)	0.88648 (19)	0.0502 (4)
H5A	0.1836	0.1312	0.9345	0.060*
H5B	0.1410	0.1643	0.9721	0.060*
C6	0.21055 (7)	0.21389 (7)	0.79664 (18)	0.0422 (3)
H6A	0.2427	0.2452	0.8715	0.051*
H6B	0.2339	0.2025	0.7184	0.051*
C7	0.11856 (7)	0.38340 (7)	0.47760 (17)	0.0426 (3)
H3N	0.2412 (9)	0.2942 (10)	0.545 (2)	0.055 (5)*
H2N	0.2062 (9)	0.3280 (9)	0.587 (2)	0.046 (4)*
H1N	0.2572 (10)	0.3321 (10)	0.691 (3)	0.061 (5)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0485 (2)	0.0679 (3)	0.0414 (3)	0.03650 (19)	0.00383 (14)	0.00618 (15)
N1	0.0367 (6)	0.0371 (6)	0.0419 (7)	0.0162 (5)	-0.0010 (5)	0.0025 (5)

N2	0.0598 (8)	0.0673 (9)	0.0734 (10)	0.0408 (8)	-0.0064 (7)	0.0012 (7)
C1	0.0320 (6)	0.0350 (6)	0.0344 (7)	0.0165 (5)	0.0010 (5)	0.0007 (5)
C2	0.0362 (7)	0.0460 (7)	0.0398 (8)	0.0145 (6)	-0.0040 (5)	-0.0015 (6)
C3	0.0374 (7)	0.0488 (8)	0.0518 (9)	0.0070 (6)	0.0015 (6)	-0.0043 (7)
C4	0.0615 (9)	0.0362 (7)	0.0624 (10)	0.0162 (7)	0.0150 (8)	0.0050 (7)
C5	0.0567 (9)	0.0488 (8)	0.0500 (9)	0.0299 (7)	0.0048 (7)	0.0119 (6)
C6	0.0379 (7)	0.0437 (7)	0.0469 (8)	0.0218 (6)	-0.0034 (6)	0.0032 (6)
C7	0.0416 (7)	0.0437 (7)	0.0417 (8)	0.0207 (6)	0.0031 (6)	-0.0006 (6)

*Geometric parameters (Å, °)*

S1—C7	1.6486 (15)	C3—C4	1.517 (3)
N1—C1	1.4978 (16)	C3—H3A	0.9700
N1—H3N	0.87 (2)	C3—H3B	0.9700
N1—H2N	0.90 (2)	C4—C5	1.520 (2)
N1—H1N	0.89 (2)	C4—H4A	0.9700
N2—C7	1.148 (2)	C4—H4B	0.9700
C1—C2	1.5132 (18)	C5—C6	1.524 (2)
C1—C6	1.5142 (18)	C5—H5A	0.9700
C1—H1A	0.9800	C5—H5B	0.9700
C2—C3	1.527 (2)	C6—H6A	0.9700
C2—H2A	0.9700	C6—H6B	0.9700
C2—H2B	0.9700		
C1—N1—H3N	109.8 (13)	C2—C3—H3B	109.2
C1—N1—H2N	110.7 (11)	H3A—C3—H3B	107.9
H3N—N1—H2N	108.9 (17)	C3—C4—C5	111.65 (13)
C1—N1—H1N	109.9 (13)	C3—C4—H4A	109.3
H3N—N1—H1N	110.1 (18)	C5—C4—H4A	109.3
H2N—N1—H1N	107.5 (16)	C3—C4—H4B	109.3
N1—C1—C2	109.91 (11)	C5—C4—H4B	109.3
N1—C1—C6	109.33 (11)	H4A—C4—H4B	108.0
C2—C1—C6	112.22 (11)	C4—C5—C6	111.14 (13)
N1—C1—H1A	108.4	C4—C5—H5A	109.4
C2—C1—H1A	108.4	C6—C5—H5A	109.4
C6—C1—H1A	108.4	C4—C5—H5B	109.4
C1—C2—C3	109.83 (11)	C6—C5—H5B	109.4
C1—C2—H2A	109.7	H5A—C5—H5B	108.0
C3—C2—H2A	109.7	C1—C6—C5	110.12 (11)
C1—C2—H2B	109.7	C1—C6—H6A	109.6
C3—C2—H2B	109.7	C5—C6—H6A	109.6
H2A—C2—H2B	108.2	C1—C6—H6B	109.6
C4—C3—C2	111.86 (13)	C5—C6—H6B	109.6
C4—C3—H3A	109.2	H6A—C6—H6B	108.2
C2—C3—H3A	109.2	N2—C7—S1	179.71 (16)
C4—C3—H3B	109.2		
N1—C1—C2—C3	178.72 (12)	C3—C4—C5—C6	-54.67 (18)

---

C6—C1—C2—C3	56.83 (16)	N1—C1—C6—C5	-179.85 (12)
C1—C2—C3—C4	-54.73 (17)	C2—C1—C6—C5	-57.63 (16)
C2—C3—C4—C5	54.43 (18)	C4—C5—C6—C1	55.66 (17)

---

*Hydrogen-bond geometry (Å, °)*

---

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N...S1 <sup>i</sup>	0.89 (2)	2.62 (2)	3.4955 (14)	167 (2)
N1—H2N...N2	0.90 (2)	1.94 (2)	2.822 (2)	170.4 (18)
N1—H3N...S1 <sup>ii</sup>	0.87 (2)	2.573 (18)	3.4214 (14)	165.5 (19)

---

Symmetry codes: (i)  $x-y+2/3, x+1/3, -z+4/3$ ; (ii)  $-x+1/3, -y+2/3, -z+2/3$ .