

**STANDARDIZATION AND EVALUATION OF
THE *CATHARANTHUS ROSEUS* EXTRACT**

NASIRA SAIF UR REHMAN

UNIVERSITI SAINS MALAYSIA

2012

**STANDARDIZATION AND EVALUATION OF
THE *CATHARANTHUS ROSEUS* EXTRACT**

NASIRA SAIF UR REHMAN

Thesis is submitted in fulfillment of the
requirements for the Degree of
Doctor of philosophy

UNIVERSITI SAINS MALAYSIA

2012

With the name of Allah, Who is utmost kind and merciful

To my family, Suny, Sabo and Sado

ACKNOWLEDGEMENTS

All praise goes to almighty Allah who showed me the path to seek the knowledge. Only Allah is 'Aleem ul Hakeem' and man knows negligible. "Nor shall men compass aught of his knowledge, except as He willeth".

I would like to say thanks to those who contributed their precious time to help me during my study with alacrity. With conceited zealous I pay respect and gratitude to my supervisor Professor Dr. Saringat Haji Baie for his multidimensional and vigilant supervision. I am thankful to my co supervisor professor Ishak Mat for his encouragement. Dr. Amin Malik Shah bin Abdul Malik Shah deserves my gratitude for providing me everything I needed in the laboratory. I also wish to express my thanks to Professor Dr. Zhari Ismail for his generous permission to work on his HPLC.

I owe my thanks to Dr. Nor Azlina Khalil, Advanced Medical and Dental Institute (AMDI) for her help in animal handling and dissection during toxicity study. Many thanks extended to Mrs. Siti Aminah for her help to perform studies on cell lines. Thanks go to Miss. Norzaina (IPPT), for providing me ample supply of leaves of *Catharanthus roseus* on her visit to Malaka on every Hari Raya. I considered this as a very unique Hari Raya gift.

I could never forget to acknowledge the most valuable support provided by Mr. Hamdan Mohammad Razak in Pharmaceutical Chemistry Lab, Mr. Abdel Rahim, Mr. Muath and Miss. Norshirin Idrees in Immunopharmacology laboratory and Miss Lia Laila during study on animals in Pharmaceutical Technology lab. I gratefully acknowledge the technical assistance of Mr. Shamsuddin, Mr. Abdu Rahim, Mr. Ibrahim, Mr. Rosali, and Mr. Firdaus for their help to keep my work running. I am indebted to my colleague Dr. Nadeem Irfan Bukhari for providing me useful information. From the core of my heart I am

thankful to my laboratory fellow Mr. Malikarjun for helping me in Statistics. The help of my dear friends on various occasions is gratefully acknowledged.

I would like to submit my gratitude to the Dean of the School of Pharmaceutical Sciences Dr. Syed Azhar Syed Sulaiman and all staff members of the School of Pharmaceutical Sciences. I also like to express my gratitude to IPS for arranging valuable workshops, training programmes and providing me the privilege of getting award of USM fellowship.

I have considered my years of study in USM as a crucial learning period. Many Special thanks to the USM for giving me opportunity to get admission and get provided with the quality knowledge and skill which would not be possible for me otherwise. During my studies I have benefited from the well set electronic Hamzah Sendut Library. Thank you for subscribing the best journals of the world for literature survey, otherwise that would not be possible to complete in time. Thanks to the staff of the library for arranging lectures in the library for teaching me how to use the electronic library and how to use “endnote” and many other programmes.

I enjoyed my stay in USM, being a mini world on its own, having students from all over the world. My Malay friends deserve many thanks and admiration for their hospitality and frequent invitations to their homes and serving me home cooked very delicious Malay traditional dishes. I never missed social life here. I got opportunities of attending many weddings and enjoyed their traditional ways. I am grateful to Professor Saringat Haji Baie, Dr. Amin Malik Shah, Dr. Mohammad Izham and their families for inviting me on food parties on different occasions such as wedding, Akika, naming ceremony, berbuka and Hari Raya. I had the privilege of being treated very special. You never let me miss my home.

I am greatly obligated to my teacher Professor Dr. Tasneem Ahmad, Director of Center for Bioequivalence, Karachi University, Pakistan, I acknowledge his help, valuable advice and permission to work in his laboratory. Finally, never enough thanks and love goes to my family. They allowed me to go abroad for study when they direly needed my presence among them.

Nasira Saif ur Rehman

TABLE OF CONTENTS

TITLE	PAGE
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	v
LIST OF TABLES	xx
LIST OF FIGURES	xxii
LIST OF EQUATIONS	xxv
LIST OF ABBREVIATIONS	xxvii
LIST OF SYMBOLS	xxxii
ABSTRAK	xxxiii
ABSTRACT	xxxv

CHAPTER 1

GENERAL INTRODUCTION

1.1	CANCER	1
1.1.1	Oncogenes and tumor suppressor genes	2
1.1.2	Molecular natural cell cycle clock	3
1.1.3	A brief history of treatment of cancer	3
1.2	DRUG DISCOVERY OF NATURAL FLORA	5
1.2.1	Lead structures of the synthetic compounds	5
1.2.2	“Natural Inhibitor of Carcinogenesis”	6
1.3	<i>CATHARANTHUS ROSEUS (C. ROSEUS)</i>	6
1.3.1	Nomenclature of <i>Catharanthus roseus</i>	6

1.3.2	Ethno medical uses of <i>Catharanthus roseus</i>	8
1.4	VINCA ALKALOIDS AND OTHER COMPOUNDS PRESENT IN <i>CATHARANTHUS ROSEUS (C. ROSEUS)</i>	9
1.4.1	Discovery of anticancer alkaloids from <i>C. roseus</i>	12
1.4.2	Earliest chemotherapeutic agents	13
1.4.3	Cancers treated by <i>Vinca</i> alkaloids	13
1.4.4	Sulfate and ditartrate salts of <i>Vinca</i> alkaloids	15
1.5	STRUCTURE OF VINCA ALKALOIDS	15
1.5.1	Structural difference and clinical activity	17
1.5.2	Derivatives of <i>Vinca</i> alkaloids	17
1.6	MODE OF ACTION OF VINCA ALKALOIDS	18
1.6.1	Target of <i>Vinca</i> alkaloids for anticancer activity	19
1.6.2	<i>Vinca</i> alkaloids as mitotic inhibitors during cell cycle	19
1.6.3	Activity of <i>Vinca</i> alkaloids in high and low concentrations	22
1.6.4	Special <i>Vinca</i> -specific high-affinity and low-affinity sites	23
1.7	CELL RESISTANCE AGAINST VINCA ALKALOIDS	23
1.7.1	Classical multiple drug resistance (MDR) of cancer cells to natural hydrophobic drug	24
1.7.1(a)	Transmembrane pump of ABC transporter (P-gp)	25
1.7.1(b)	Multi drug resistance protein	26
1.7.1(c)	P450 3A4 (CYP 3A4) enzyme	26
1.7.1(d)	Alteration in tubulin properties	27
1.8	DRAW BACK OF PURIFIED VINCA ALKALOIDS	27

1.8.1	Low therapeutic index	27
1.8.2	Multiple drug resistance	27
1.8.3	Development of neurotoxicity	28
1.8.4	Dose-limiting toxicity	28
1.9	ORAL ADMINISTRATION OF CYTOTOXIC DRUG	28
1.9.1	General introduction of oral bioavailability	29
1.9.2	Bioavailability of lipophilic and hydrophilic drugs	29
1.9.3	Factors affecting bioavailability	30
1.9.4	Barriers against diffusion of drug molecules	30
1.9.5	Bioavailability of oral cytotoxic compounds	31
1.9.6	The higher doses of <i>Vinca</i> alkaloids can escape efflux	32
1.10	USE OF <i>C. ROSEUS</i> EXTRACT AS ANTICANCER (C. ROSEUS) EXTRACT AS ANTICANCER DRUG	32
1.10.1	Report of the world health organization (WHO)	32
1.10.2	Potential therapeutic indications of <i>C. roseus</i> extract	33
1.10.2(a)	Oxygen radical absorbance capacity (ORAC)	33
1.10.2(b)	<i>C. roseus</i> extract inhibits Pgp and CYP3A4 activity	33
1.10.2(c)	Antiangiogenesis activity	34
1.10.2(d)	<i>C. roseus</i> extract inhibited the progression of tumor	34
1.10.2(e)	<i>C. roseus</i> extract causes immune suppression	34
1.10.2(f)	Cytotoxicity of <i>C. roseus</i> extract against cancer cells	34
1.10.3	Quantification of the total extract	35

1.11	THEORETICAL FRAMEWORK	35
1.12	Contribution of variety of chemical types in <i>Catharanthus roseus</i> <i>C. roseus</i> extract	35
1.11.2	Synergy“, „polyvalence“ and „plurality“	36
1.12	JUSTIFICATION OF THE STUDY	38
1.13	SCOPE AND OBJECTIVES	38

CHAPTER 2

HPLC METHOD VALIDATION AND STABILITY STUDY

2.1	INTRODUCTION	40
2.1.1	High performance liquid chromatography	40
2.1.2	HPLC method for <i>Vinca</i> alkaloids	40
2.1.3	Reversed phase chromatography	41
2.1.4	Validation of HPLC method	41
2.1.4(a)	Basic steps for validation of method	42
2.1.4(b)	Basic points considered in preparation of dilutions	42
2.1.5	Procedure of validation	43
2.1.5(a)	Construction of the calibration curve to assess linearity	43
2.1.5(b)	Intra-day and inter-day recovery study	43
2.1.6	Specifications for method validation	44
2.1.7	Parameters for validation of HPLC Methods	46
2.1.7(a)	Accuracy	46

2.1.7(a)(i)	Accuracy assessment by standard curve	46
2.1.7(a)(ii)	Accuracy assessment by recovery study	47
2.1.7(b)	Precision and sensitivity	48
2.1.7(c)	Linearity	49
2.1.7(d)	Range	49
2.1.7(e)	Specificity/selectivity	50
2.1.7(f)	Limit of detection	50
2.1.7(g)	Limit of quantification	51
2.1.7(h)	Robustness	51
2.1.7(i)	Reproducibility	52
2.1.7(j)	Repeatability	52
2.1.8	Stability	52
2.1.9	Objective of method validation study	52
2.2	EXPERIMENTAL FOR HPLC	53
2.2.1	Materials	53
2.2.2	HPLC Instrumentation	53
2.2.2(a)	Mobile phase	53
2.2.2(b)	Preparation of standard solutions	54
2.2.2(c)	HPLC method and conditions	54
2.2.2(d)	Construction of calibration curve	54
2.2.3	Method validation	54
2.2.3(a)	Determination of Linearity and range	54
2.2.3(b)	Evaluation of accuracy and precision	55

2.2.3(c)	Determination of recovery	55
2.2.3(d)	Determination of limit of quantification (LOQ) and limit of detection (LOD)	56
2.3	RESULTS	56
2.3.1	HPLC method validation	57
2.3.1(a)	Specificity	57
2.3.1(b)	Linearity	60
2.3.1(c)	Recovery and accuracy	63
2.3.1(d)	The precision	64
2.3.1(e)	Limit of detection and quantification	65
2.3.2	Stability of VBL and VCR	66
2.4	DISCUSSION	66
2.4.1	HPLC	66
2.4.2	Stability	67
2.5	CONCLUSION	67

CHAPTER 3

QUANTIFICATION OF VINBLASTINE/ VINCRISTINE IN THE EXTRACT OF *CATHARANTHUS ROSEUS* LEAVES BY HPLC METHOD

3.1	INTRODUCTION	69
3.1.1	Methods of analysis of the alkaloids in plant extract	69
3.1.2	HPLC methods for analysis of alkaloids	69
3.1.3	Method of choice for analysis of alkaloids in <i>C. roseus</i>	70

3.1.4	HPLC techniques and instrumentation	72
3.1.5	Validation of Analytical procedure	72
3.1.6	External standard method	73
3.1.7	Material preparation	73
3.2	EXPERIMENTAL: MATERIAL AND METHOD	74
3.2.1	Plant	74
3.2.2	Test material Preparations	74
3.2.3	Preparation of standard, test and buffer solutions	77
3.2.4	HPLC method- development and method-validation	77
3.2.5	Quantification of contents of VBL and VCR	78
3.2.6	Identification of peaks of VBL and VCR	78
3.2.7	Objective	78
3.3	RESULTS	79
3.3.1	Chromatograms of standard solutions	79
3.3.2	Identification of the peaks in the extract	79
3.3.3	Results at wavelength 297nm	82
3.3.3(a)	Peaks of mixed standards	82
3.3.3(b)	Ethanol extract E ₆ un-spiked and spiked	84
3.3.3(c)	Water extract WI	86
3.3.4	Results at wavelength 262 nm	87
3.3.4(a)	Peaks of mixed standards	87
3.3.4(b)	Extract E ₆ , un-spiked and spiked	87

3.3.4(c)	Overlapped chromatograms of WI, spiked and unspiked	87
3.4	DISCUSSION	93
3.5	CONCLUSION	94

CHAPTER 4

IN-VITRO STUDY OF THE CYTOTOXIC ACTIVITY OF THE EXTRACT OF *CATHARANTHUS ROSEUS*

4.1	INTRODUCTION	95
4.1.1	General design and precision of bioassay	95
4.1.2	Cell viability	96
4.1.3	Cytotoxicity testing	96
4.1.3(a)	Method of anti proliferation activity	97
4.1.3(b)	Selection of solvent and precautions	98
4.1.3(c)	Cell harvesting	98
4.1.3(d)	Cell preparation	99
4.1.4	Techniques to determine the cell growth inhibition	99
4.1.5	Selectivity	99
4.2	EXPERIMENTAL: MATERIAL AND METHOD	100
4.2.1	Preparation of crude extract	100
4.2.2	Cell lines	100
4.2.3	Preparation of medium	101
4.2.4	Determination of percentage of cell viability (CV)	101
4.2.4(a)	Cell harvesting and cell preparation for CV testing	101

4.2.4(b)	Preparation of test solutions from crude extract	102
4.2.4(c)	Procedure of cell viability test	102
4.2.4(d)	Statistical Analysis	103
4.2.5	Cytotoxicity Test	103
4.2.5(a)	Preparation of test solutions from the extract	103
4.2.5(b)	Procedure of cytotoxicity test	103
4.2.5(c)	Methyl Thiazoldiphenyl Tetrazolium (MTT) Assay	104
4.2.5(d)	Statistical analysis	105
4.3	RESULTS	106
4.3.1	Determination of the effect of water extracts on 3T3 (normal colon) cells viability	106
4.3.2	Determination of the effect of water extracts on HT29 (colon cancer cells) viability	110
4.3.3	Determination of the effect of ethanol extracts on HT29 (colon cancer cells) viability	113
4.3.4	Determination of IC ₅₀	117
4.3.5	Effect of extract on colon cancer cells HCT 116	118
4.3.6	Effect of extract on colon cancer cells HT 29	119
4.3.7	Effect of extract on breast cancer cells T47D	119
4.4	DISCUSSION	123
4.5	CONCLUSION	123

CHAPTER 5

IN VIVO SINGLE DOSE ACUTE ORAL TOXICITY STUDY OF THE *CATHARANTHUS ROSEUS* EXTRACT

5.1	INTRODUCTION	124
5.1.1	Preclinical safety study	124
5.1.2	Bioassay	125
5.1.3	Animal biological system, as surrogate to human system	125
5.1.4	Toxicity test system	126
5.1.5	Maximum tolerated dose (MTD)	126
5.1.6	MTD from animal to human	126
5.1.6(a)	Dose conversion factor	127
5.1.6(b)	Equation to convert doses from animal to animal	127
5.1.7	Selection of animal model	128
5.1.8	Experimental design	129
5.1.9	Ethical allegation	130
5.1.10	Single dose acute oral toxicity testing	131
5.1.11	Limit test	131
5.1.12	Dose and number of animals for acute oral toxicity test	132
5.1.13	Fixed Dose Procedure (FDP)	132
5.1.14	Testing dose above than 2000 mg/kg body weight	135
5.1.15	Procedure of Single Dose Acute Oral Toxicity (AOT)	135
5.1.16	Preparation and administration of test material	136
5.1.17	Gavages	136
5.1.18	Post treatment clinical observation	137

5.1.19	Anaesthesia and Euthanasia	138
5.1.20	Necropsy	139
5.1.21	Disposal of euthanasia animals	139
5.2	EXPERIMENTAL: MATERIAL AND METHOD	139
5.2.1	Animal	140
5.2.2	Limit test	140
5.2.3	Animal model for acute oral toxicity test	140
5.2.4	Caging, housing, care and maintenance	141
5.2.5	Control and test animals marking	141
5.2.6	Administration of test substance	142
5.2.6(a)	Test animal	142
5.2.6(b)	Control animal	142
5.2.7	Animal diet	143
5.2.8	Experimentation for single dose acute oral toxicity test	144
5.3	RESULTS	145
5.3.1	Post treatment cage side observations	145
5.3.1(a)	General behavior and clinical observations	145
5.3.1(b)	Hunched posture in test female animal	146
5.3.2	Weight loss and weight gain observations	150
5.3.3	Comparison in test and control animals	153
5.3.4	Mortality or signs of abnormality	153
5.3.5	Clinical examination before Euthanasia	154
5.3.6	Euthanasia	154

5.3.7	Necropsy examination	154
5.4	DISCUSSION	155
5.5	CONCLUSION	156

CHAPTER 6

EX-VIVO ANTI- ANGIOGENESIS ACTIVITY OF THE *CATHARANTHUS ROSEUS* EXTRACT

6.1	INTRODUCTION	157
6.1.1	Angiogenesis	157
6.1.2	Mechanism of angiogenesis	157
6.1.3	Tumor angiogenesis	158
6.1.4	Anti-angiogenesis assays	158
6.1.5	Rat aorta model of angiogenesis	160
6.1.6	Growth of new micro-vessels	161
6.2	EXPERIMENTAL: MATERIAL AND METHOD	162
6.2.1	Rats	162
6.2.2	Preparation of control and test Solutions	162
6.2.3	Preparation of three-dimensional aortic ring culture	163
6.2.4	Procedure of rat aorta angiogenesis assay	163
6.3	RESULTS	164
6.4	DISCUSSION	170
6.5	CONCLUSION	170

CHAPTER 7

GENERAL DISCUSSION AND SUGGESTIONS FOR FURTHER STUDY

7.1	GENERAL DISCUSSION	171
7.1.1	Clinical practice in treatment of cancer	171
7.1.2	Use of Crude extracts in treatment of cancer	171
7.1.3	The Study of <i>C. roseus</i> extract as anticancer drug	172
7.1.4	Oral administration of <i>Vinca</i> alkaloids	172
7.1.5	Anticancer activity of <i>Catharanthus roseus</i> extract	172
7.1.6	Use of extract in modern medicine	173
7.2	THE STUDY OF <i>C. ROSEUS</i> EXTRACT	173
7.2.1	Method validation and stability	174
7.2.2	Quantification	175
7.2.3	The <i>in vitro</i> cytotoxicity determination	175
7.2.4	The <i>in vivo</i> toxicity study	176
7.2.5	The <i>ex vivo</i> anti angiogenesis activity	176
7.3	DISCUSSION	176
7.4	CONCLUSION	177
7.5	SUGGESTIONS FOR FURTHER STUDY	177
	REFERENCES	179
	APPENDICES	

A 2.1 Mean area \pm SD of HPLC method validation assay for VBL	199
A 2.2 Mean area \pm SD of HPLC method validation assay for VCR	200
A 2.3 HPLC validation: standard curve for VCR (6 replicates, n=7)	201
A 2.4 HPLC method validation another set of 6 replicates (n=7)	204
A 2.5 HPLC validation: standard vinblastine VBL (6 replicates, n=7)	207
A 2.6 Second set of standard curves of VBL 6 replicates n=7	210
A 3.1 Mean area \pm SD of the extracts of the <i>Catharanthus roseus</i> at 262 nm	213
A 3.2 Mean area \pm SD of the extracts of the <i>Catharanthus roseus</i> at 297 nm	216
A 4.1 Optical density data of 3T3 normal colon cells, exposed to aqueous extracts W ₄ (Sample 4) and W ₂ (Sample 2)	219
A 4.2 Standard deviation for normal colon cells 3T3 data at different concentration and time, exposed to aqueous extracts W ₄ (Sample 4) and W ₂ (Sample 2)	219
A 4.3 Growth curve of 3T3 normal colon cancer cells exposed to aqueous extracts W ₄ (Sample 4) and W ₂ (Sample 2)	221
A 4.4 Data of HT29 colon cancer cells growth when exposed to aqueous extract W ₄ (Sample 4) and W ₂ (Sample2)	224
A 4.5 Calculation of standard deviation and construction of growth curves: HT29 colon Cancer cells exposed to aqueous extract W ₄ (Sample 4) and W ₂ (Sample2)	224

A 4.6 HT29 colon cancer cell growth curves, exposed to aqueous extract W ₄ (Sample 4) and W ₂ (Sample2)	226
A 4.7 Data of HT29 Cell growth when exposed to ethanol extract E ₄ (Sample 4) and E ₂ (Sample2)	229
A 4.8 Calculation of standard deviation and construction of growth curves: HT29 colon cancer cells exposed to ethanol extracts E ₄ (Sample 4) and E ₂ (Sample2)	229
A 4.9 HT29 colon cancer cells growth curves: HT29 colon cancer cells exposed to ethanol extract E ₄ (Sample 4) and E ₂ (Sample2)	231
A 4.10 Percent inhibition and standard deviation (T47D colon cancer cells HT29 and breast cancer cells HCT116)	234
A 5.1 Approval letter of animal ethics committee to perform single dose acute oral toxicity test	235
List of oral and poster presentations	236

LIST OF TABLES

No.	Title	Page
1.1	A brief history of clinical cytotoxic drugs	5
1.2	Important alkaloids reported from <i>Catharanthus roseus</i>	10
1.3	Frequently used salts of anticancer alkaloids from <i>C. roseus</i>	11
1.4	Cancers treated by <i>Vinca</i> alkaloids	14
2.1	Specifications for validation assay	45
2.2	The linear regression coefficient (R^2) of VBL	61
2.3	The linear regression coefficient (R^2) of VCR	62
2.4	HPLC method validation for vinblastine (VBL)	63
2.5	HPLC method validation for vincristine (VCR)	63
2.6	Accuracy, precision and Coefficient of correlation (CV) of VBL	64
2.7	The values of Regression equation: R^2 , LOD and LOQ for VCR	65
2.8	The values of Regression equation: R^2 , LOD and LOQ for VBL	65
3.1	UV range for <i>C. roseus</i> alkaloids on HPLC	71
3.2	Contents of vinblastine and vincristine present in the extract	92
3.3	Total contents of VCR and VBL present in the extract	92
4.1	Percentage of normal colon (3T3) cell viability in presence of aqueous extracts W_4 (sample 4) and W_2 (sample 2)	107
4.2	Percentage of colon cancer (HT29) cell viability in presence of aqueous extracts W_4 (sample 4) and W_2 (sample 2) after 24, 48 and 76 hours of treatment	110
4.3	Percentage of colon cancer (HT29) cell viability in presence of ethanol extracts E_4 (sample 4) and E_2 (sample 2) after 24, 48 and 76 hours of treatment	113

4.3	Percentage of colon cancer (HT29) cell viability in presence of ethanol extracts E ₄ (sample 4) and E ₂ (sample 2) after 24, 48 and 76 hours of treatment	113
4.4	IC ₅₀ of WI, WII, W ₆ , and E ₆	118
4.5	IC ₅₀ of extracts WI, WII, W ₆ and E ₆ on human colon cancer cells HCT-116 measured by the MTT assay	120
4.6	<i>In vitro</i> cytotoxicity of <i>C. roseus</i> extracts on human colon cancer cells HT-29 measured by the MTT assay	121
4.7	<i>In vitro</i> cytotoxicity of <i>C. roseus</i> extracts on human breast cancer cells T-47D measured by MTT assay	122
5.1	Oral LD ₅₀ for acute oral toxicity testing	134
5.2	Recommended gavages size and volume of oral solution	137
5.3	Animal marking	141
5.4	Gavages and volume of test solution	143
5.5	Quantity of normal feed	144
5.6	Day 1, pre and post-treatment cage-side observations	147
5.7	Daily post treatment clinical observations for 14 days	148
5.8	Daily check for food intake for first 7 days and weight loss at day 7 and day 14	149
5.9	Weight loss and gain in test (T) and control (C) animals	151
5.10	A summary of general conditions of animal and weight gain and loss after administration of test substance	152
5.11	Fixed oral LD ₅₀ cut-off values	156
6.1	Angiogenesis models	159

LIST OF FIGURES

No	Title	Page
1.1	<i>Catharanthus roseus</i>	7
1.2	Chemical structure of <i>Vinca</i> alkaloids	16
1.3	The cell cycle clock	21
2.1A	Chromatogram (A) peak of blank	58
2.1B	Chromatogram (B) Peak of vinblastine	59
2.1C	Chromatogram (C) Peak of vincristine	59
2.2	Standard curve (n=6) of vinblastine (VBL)	61
2.3	Standard curve (n=6) of vincristine (VCR)	62
3.1	Flow chart of the extraction process	76
3.2A	Chromatogram (A) Peak of VBL, Retention time 3.94 minutes	80
3.2B	Chromatogram (B) Peak of VCR, Retention time 3.52 minutes	80
3.2 C	Chromatogram (C) Peak of mixed standards, vinblastine (VBL) Rt 3.997minutes and vincristine (VCR) Rt 3.58 minutes	81
3.3A	Retention time (Rt) of vincristine (VCR) was 3.26 minutes	83
3.3B	Retention time (Rt) of vindoline (VDL) was 3.65 minutes, vinblastine (VBL) was 3.98 minutes and catharanthine CTR) was 4.14 minutes	83
3.4A	Chromatogram (A) of un-spiked extract E ₆ solution (1mg/ml)	84
3.4B	Chromatogram (B) of extract E ₆ spiked with mixed solution of standards of vincristine (VCR), vindoline (VDL), vinblastine (VBL), and catharanthine (CTR)	85
3.4C	Chromatograms (C): overlapped peaks of extract E ₆ (A) spiked with mixed standards vincristine (VCR), vindoline (VDL), vinblastine (VBL), and catharanthine (CTR) (B) to see the position of peaks (Un-spiked is black and spiked is red)	85

3.5	A comparison of the chromatogram of the solution of extract WI with overlapped chromatogram of extract spiked with mixed standards vincristine (VCR), vindoline (VDL), vinblastine (VBL), and catharanthine (CTR) at 297 nm (Un-spiked is black and spiked is red)	86
3.6A	Chromatogram (A) Peak of vincristine (VCR) observed at 262 nm	88
3.6B	Chromatogram (B) Peaks of vindoline (VDL), vinblastine (VBL) and catharanthine (CTR) observed at 262 nm	88
3.7A	Chromatogram (A) Peaks of un-spiked ethanol extract E ₆ (262nm)	89
3.7B	Chromatogram B: Extract E ₆ , spiked with solution of mixed standards vincristine (VCR), vindoline (VDL), vinblastine (VBL), and catharanthine (CTR) at 262 nm	89
3.7C	Chromatogram C: The overlapping of peaks of ethanol extracts E ₆ and standard peaks allow alkaloids to be amplified and viably identified at 262nm	90
3.8	Chromatogram showing overlapping of peaks of WI and mixed standards. Overlapping allows alkaloids to be amplified and viably identified for quantification at 262 nm	90
4.1A	Percentage viability of 3T3 (normal cells) in aqueous extracts W ₄ (sample 4) and W ₂ (sample2) after 24 hours of treatment	108
4.1B	Percentage viability of 3T3 (normal cells) in aqueous extracts W ₄ (sample 4) and W ₂ (sample2) after 48 hours of treatment	108
4.1C	Percentage viability of 3T3 (normal cells) in aqueous extracts W ₄ (sample 4) and W ₂ (sample2) after 72 hours of treatment	109
4.2A	Percentage viability of HT29 colon cancer cells in aqueous extracts W ₄ (sample 4) and W ₂ (sample 2) after 24 hours of treatment	111
4.2B	Percentage viability of HT29 colon cancer cells in aqueous extracts W ₄ (sample 4) and W ₂ (sample 2) after 48 hours of treatment	111
4.2C	Percentage viability of HT29 colon cancer cells in aqueous extracts W ₄ (sample 4) and W ₂ (sample 2) after 72 hours of treatment	112

4.3A	Percentage viability of HT29 colon cancer cells in ethanol extracts E ₄ (sample 4) and E ₂ (sample 2) after 24 hours of treatment	114
4.3B	Percentage viability of HT29 colon cancer cells in ethanol extracts E ₄ (sample 4) and E ₂ (sample 2) after 48 hours of treatment	114
4.3C	Percentage viability of HT29 colon cancer cells in ethanol extracts E ₄ (sample 4) and E ₂ (sample 2) after 72 hours of treatment	115
4.4	Cancer cell morphology after 48 hours of treatment with extracts of <i>Catharanthus roseus</i> (500µg/ml). A, D, G: untreated cell control. B, E, H: treatment with sample of aqueous extract W ₂ . C, F, I: treatment with sample of aqueous extract W ₄ . A, B, C: HT29 colon cancer cells treated with aqueous extract. D, E, F: 3T3 normal colon cells treated with aqueous extract. G, H, I: HT29 colon cancer cells treated with ethanol extract.	116
4.5	Log- dose versus response of extracts on human colon cancer cells HCT-116	120
4.6	Log- dose versus response of extracts on human colon cancer cells HT-29	121
4.7	Log- dose versus response of extracts on human breast cancer cells T47 D	122
5.1	Test procedure with starting dose (2000 mg/kg body weight)	133
6.1 A&B	The images of rat aorta angiogenesis assay treated with ethanol extract E ₆ , showing absence of growth of new blood vessels	166
6.2 A&B	The images of rat aorta angiogenesis assay treated with extract WI showing absence of growth of new blood vessels	167
6.3 A&B	Images of negative control obtained on the day 5: sprouting of new blood vessels	168
6.4 A&B	Images of positive control suramine on day 5: prohibition of growth of micro vessels	169

LIST OF EQUATIONS

Equation	Page
Detection limit = $3.3 \sigma/s$	49
$Y = mx + c$	51
$m = \Delta y / \Delta x$	52
$Z_i = Y_i / x_i$	53
$Z_i = \mu_z + D_i + e_i$	53
$\text{LOD} = \frac{3.3 \sigma_D}{m}$	56
$\text{LOD} = \frac{10 \sigma_D}{m}$	56
$\% \text{ CV} = (\text{standard deviation of array } \sigma / \text{average of array}) \times 100$	62
$\% \text{ CV} = (\sigma / \bar{y}) \times 100$	62
$Y = mx + c$	62
$\text{LOD} = \frac{3.3 \sigma_D}{m}$	63

$$\% \text{ accuracy} = (\text{Obtained Value} / \text{True value}) \times 100 \quad 63$$

$$\% \text{ Precision} = (\sigma / \mu) \times 100 \quad 63$$

$$Y = mx + c \quad 71$$

$$\text{The \% of cell viability} = \frac{\text{Optical Density of treated cells}}{\text{Optical Density of control cells}} \times 100 \quad 102$$

$$\% \text{ inhibition} = \frac{(\text{Control OD} - \text{sample OD})}{\text{Control OD}} \times 100 \quad 103$$

$$\text{Viable cells (\%)} = \frac{(\text{Total cells} - \text{dead cells})}{\text{Total cells}} \times 100 \quad 103$$

$$\text{The \% of cell viability} = \frac{\text{OD of treated cells}}{\text{OD of control cells}} \times 100 \quad 110$$

$$\text{Viability} = (\text{OD sample} - \text{OD blank}) / (\text{OD control} - \text{OD blank}) \quad 113$$

$$\% \text{ of growth inhibition} = (1 - \text{viability}) \times 100 \quad 113$$

LIST OF ABBREVIATIONS

Abbreviations	Description
AEC	Animal Ethics Committee
aFGF	acidic fibroblast growth factor
ACPE	Accreditation Council for Pharmacy Education
ANOVA	Analysis of Variance
AOT	Acute Oral Toxicity
ATCC	American Type Cell Culture Collection
BCNU	Bis Chloroethyl Nitrous Urea
bFGF	basic fibroblast growth factor
Bpm	Beats per minute
Bw	Body weight
CAM	Chick chorioallantoic membrane
CAMs	Complementary and Alternative Medicines
CCM	Chick Chorioallantoic Membrane
cdk	Cycline-dependent kinase
CDER	Center for Drug Evaluation and Research
CMC3	Chemistry Manufacturing Control Coordinating Committee
<i>C roseus</i>	<i>Catharanthus roseus</i>
CSO	Cage Side Observations
CTR	Catharanthine
CV	Coefficient of Variation
CVT	Cell Viability Test

CYP3A4	Cytochrome P450, family 3, subfamily A, Polypeptide 4
cyt. Arabin	Cytosine Arabinoside
DMSO	Dimethylsulphoxide
DNA	Deoxyribo Nucleic Acid
DNA	Deoxyribo Nucleic Acid
EDTA	Ethylene Diamine Tetra Acetic acid
EMA	European Medicines Agency
FBS	Foetal Bovine Serum
FDA	Federal Drug Administration
FDP	Fixed Dose Procedure
G	Gavages (feeding tube)
G1 Phase	Gap 1 phase
GC	Gibco, Canada
GCP	Good collection practice
GHS	Globaly Harmonized System for classification and labeling
GI	Gastro Intestinal
GIT	Gastro Intestinal Tract
GLP	Good Laboratory Practice
HCT-116	Colon carcinoma cells
HPLC	High Performance Liquid Chromatography
HT-29	Colon carcinoma cells
IACUC	Institutional Animal Care and Use Committee
IC ₅₀	The half maximal inhibitory concentration
ICH	International Conference of Harmonization

IPPT	Institut Perubatan dan Pergigian Termaju (Advanced Medical and Dental Institute)
IPS	Institut pengajian siswazah (Institute of Postgraduate Studies)
LD50	Lethal dose for 50% of population
LOD	Limit of Detection
LOQ	Limit of Quantification
<i>m</i>	Slope of the calibration curve
MAKNA	Majlis Kanser Nasional
MDR	Multi drug resistance
MELD ₁₀	Mouse equivalent LD ₁₀
M phase	Mitotic phase
MRP	Multi-drug Resistance-associated Proteins
MIAs	Monoterpenoid indole alkaloids
MTAs	Microtubules targeting agents
MTD	Maximum tolerated dose
MTT	Methyl thiazoldiphenyl tetrazolium
NCI	National Cancer Institute
NY	New York
N mustard	Nitrogen mustard
OD	Optical density
OCDE	Organisation de coopération et de développement économiques
OECD	Organization of economic cooperation and development
ORAC	Oxygen radical absorbance capacity
PBS	Phosphate buffer saline

Pgp	P-glycoprotein
PDA	Photo diode array
PMS	Phenazine methosulphate
RAM	Rat aorta model
R ²	Correlation coefficient
RO	Reverse osmoses
RPC	Reversed phase chromatography
RSD	RSD relative standard deviation
Rt	Retention time
SD	Starting dose
SD	Standard deviation
S phase	Synthetic phase
3T3	Normal cells (colon)
TSGs	Tumor suppressor genes
T-47D	Breast carcinoma cells
Tc	Test animal (control)
Tm	Test animal (male)
Tf	Test animal (female)
TSGs	Tumor suppressor genes
UDP	Up and down procedure
USFDA	United States Food and Drug Authority
USM	University Sains Malaysia
UWL	Unstirred water layer
VEGF	Vascular endothelial growth factors
VRLP	<i>Vinca rosea</i> leaves extract

VBL	Vinblastine
VCR	Vincristine
VDL	Vindoline
WHO	World Health Organization

LIST OF SYMBOLS

Symbol	Description
α	Alpha (unclassified)
m	Slope of the calibration curve
I	Index for the day
S	Slope
σ_D	Standard Deviation
σ_D^2	Day to day variability
σ_e^2	Within day or intraday variability.
σ	Sigma (Standard deviation of the response that is y- intercept)
$10 \sigma_D$	Relative standard deviation is 10 % exhibiting a signal to noise ratio 0.1
μ	Micro
μ	Mu
μ_z	Unknown parameter

PEMPIAWAIAN DAN PENILAIAN EKSTRAK *CATHARANTHUS ROSEUS*

ABSTRAK

Satu ekstrak piawai *Chantarantus roseus* telah disediakan dengan menggunakan kaedah pengekstrakan air dan ethanol. Kandungan bahan aktif ekstrak telah ditentukan dengan kaedah HPLC. Jumlah alkaloid *Vinca* yang digunakan sebagai bahan antikanser vinblastina (VBL) dan vincristina (VCR) telah ditentukan. Kandungan VBL dan VCR ditentukan untuk menilai sumbangan alkaloid-alkaloid ini terhadap aktiviti anti kanser ekstrak yang dihasilkan. Pengesahan kaedah HPLC dilakukan berdasarkan prosedur yang terdapat dalam garispanduan *USFDA* dan *ICH*, dan didapati efisien dan boleh dihasilkan semula. Had minimum yang boleh dikesan untuk VCR adalah 0.25 µg/ml dan VBL ialah 0.5 µg/ml. Kelinearan yang boleh diterima dipamerkan dengan pekali regresi (R^2), iaitu 0.9999 hingga 1. Alkaloid pertama yang dielut adalah vinkristina dengan masa retensinya (R_t) adalah 3.2 minit, diikuti dengan vindolina 3.56 minit, vinblastina 3.89 minit dan katarantina 4.14 minit. Analisis HPLC daripada ekstrak *Catharanthus roseus* menunjukkan bahawa kandungan VBL adalah sama dalam semua kelompok ekstrak, namun demikian kepelbagaian dalam kandungan VCR turut diperhatikan. Kandungan ekstrak *Catharanthus roseus* yang terkira adalah rendah. Keputusan ekstrak etanol E_1 , E_3 , E_4 , E_6 dijulat sebagai 11.9 µg/10mg, 9.0 µg/10mg, 12.3 µg/10mg dan 14.4 µg/10mg bagi vinblastina dan kandungan ekstrak air W_1 , W_3 , W_4 , W_6 dijulat sebagai 8.2 µg/10mg, 11.0 µg/10mg, 10.3 µg/10mg dan 7.5 µg/10mg bagi vinblastina. Keputusan ekstrak E_1 , E_3 , E_4 , E_6 , W_1 , W_3 , W_4 , W_6 dijulat sebagai 19.9 µg, 13.2 µg, 8.1 µg, 26.3 µg, 8.2 µg, 13.1 µg, 11.1 µg, and 11.6 µg/10mg bagi vincristina. Meskipun fakta bahawa kandungan alkaloid wujud dalam ekstrak *Catharanthus roseus*, namun ia adalah rendah apabila terdedah terhadap lini sel kanser yang berbeza, ekstrak W_1 , W_2 , W_6 dan E_6 mempamerkan aktiviti antikarsinogen. Kajian lini sel menunjukkan bahawa semua ekstrak yang terpilih untuk kajian ini adalah agen sitotoksik yang baik. Ekstrak etanol (E_6) menunjukkan aktiviti kesitotoksikan yang boleh diterima terhadap sel kanser payu dara T47D (IC_{50} 12 µg/ml). Ekstrak air (W_6) menunjukkan aktiviti yang boleh diterima terhadap kedua-dua lini sel kanser kolon

HCT116 (IC₅₀ 10µg) dan HT29 (IC₅₀ 1.98µg). Ekstrak WI (etanol/air) juga aktif terhadap kedua-dua lini sel kanser kolon HCT116 (IC₅₀ 12.28µg) dan HT29 (IC₅₀ 1.65µg). Ketoksikan oral akut adalah ketoksikan yang terhasil dalam tempoh tidak melebihi 24 jam, selepas sesuatu drug yang diberikan. Semasa ujian, pemerhatian tepi-sangkar (cage side observations, CSO) dilakukan secara berterusan untuk menentukan ketidaknormalan dan kemorbidan. Pembolehubah berat badan dan pengambilan makanan diawasi dengan teliti. Tiada perubahan dalam pembolehubah ini ditemui. Perubahan dalam pembolehubah ini diketahui sebagai tanda ketoksikan pertama. Kajian Ketoksikan Oral Akut Dos Tunggal menunjukkan bahawa ekstrak air dan etanol didapati tidak toksik jika diberi secara oral. Kajian angiogenesis menunjukkan bahawa larutan ekstrak *Catharanthus roseus* boleh merencat pertumbuhan saluran darah. Aktiviti antiangiogenesis dalam ekstrak diuji menggunakan Model Aorta Tikus (Rat Aorta Model RAM). Diperoleh bahawa, ekstrak ini merupakan agen antiangiogenesis yang baik pada kepekatan 10 µg per ml. Oleh itu, daripada kajian *in vitro*, *in vivo* dan *ex vivo* dapat disimpulkan bahawa ekstrak *Catharanthus roseus* adalah sitotoksik bagi lini sel kanser, tidak toksik apabila diberikan secara oral kepada haiwan dan juga mempunyai aktiviti antiangiogenesis.

STANDARDIZATION AND EVALUATION OF THE EXTRACT OF *CATHARANTHUS ROSEUS*

ABSTRACT

A standardized *Catharanthus roseus* extract has been prepared by water and ethanol method. The widely used anticancer *Vinca* alkaloids, vinblastine (VBL) and vincristine (VCR) present in the extract were determined by using HPLC. Contents of VBL and VCR were determined to assess the contribution of these alkaloids towards the anticancer activity of the extract. The method was validated in accordance to the procedure mentioned in the Guidelines of *USFDA* and *ICH*. The method was found efficient and reproducible. The minimum detectable limit of VCR was 0.25 µg/ml and VBL was 0.50 µg/ml. An acceptable linearity was exhibited with regression coefficient (R^2) of 0.9999 to 1. The first eluted alkaloid was vincristine its retention time (R_t) was 3.2 minutes followed by vindoline 3.56 minutes, vinblastine 3.89 minutes and catharanthine 4.14 minute. The content of VBL for the ethanol extracts E₁, E₃, E₄ and E₆ were 11.9µg/100mg, 9.0µg/100mg, 12.3µg/100mg and 14.4µg/100mg of the dried extract. The contents of VBL determined in water extracts W₁, W₃, W₄, W₆ were 8.2µg/100mg, 11.0µg/100mg, 10.3µg/100mg and 7.5µg/100mg. Amount of VCR present in *Catharanthus roseus* extracts E₁, E₃, E₄, E₆, W₁, W₃, W₄, W₆ was determined as 19.9µg, 13.2µg, 8.1µg, 26.3µg, 8.2µg, 13.1µg, 11.1µg, and 11.6 µg/100mg of the dried extract. The study of the extract on cancer cells demonstrated good cytotoxic activity. Ethanol extract E₆ showed acceptable cytotoxicity against breast cancer cells T47D (IC₅₀ 12µg/ml). Water extract W₆ showed pronounced anticancer activity against colon cancer cell lines HCT116 (IC₅₀ 10 µg/ml) and HT29 (IC₅₀ 1.98 µg). Ethanol/water extract WI was also active against both colon cancer cell lines HCT116 (IC₅₀ 12.28 µg) and HT29 (IC₅₀ 1.65 µg). In single dose acute oral toxicity test, variables (body weight and feed intake) were carefully checked and no significant changes in these variables were initiated. There was no difference in physical state, behavior and response of the control and the treated groups. This study indicated that water and ethanol extracts were less toxic and can be administered orally. Antiangiogenesis study in Rat Aorta Model (RAM) revealed that the 10µg/ml solution of the alcohol

water extract WI and Ethanol extract E₆ of *Catharanthus roseus* inhibited the growth of blood vessels. The extracts appeared to be good antiangiogenic agent. The current study indicated that the *Catharanthus roseus* extract is less toxic when administered orally, exhibited cytotoxicity on cell lines and demonstrated good anti-angiogenesis activity.

CHAPTER 1

GENERAL INTRODUCTION AND LITERATURE SURVEY

1.1 CANCER

Cancer is produced from a normal cell when it becomes abnormal. Loss of growth control in a single cell causes trillions of cells to arise from a single cell (Bissell, 1981). The body has an unknown mechanism to eliminate cells losing growth control, but sometimes due to a variety of chromosomal changes, the cell cycle is driven crazy into nonstop division. The transcription factors such as E2F and c-myc are responsible for the cell cycle or cell growth, and involve in forcing cells into apoptosis or cell death. Thus, cell birth and cell death are initiated by the same pathway (Evan and Littlewood, 1998; King and Cidlowski, 1998).

Cancer results from the genetic and epigenetic abnormalities in susceptible cells (Ponder, 2001; Houghton *et al.*, 2007). In general, somatic mutations are involved, but it can also be inherited. Linkages are found through linkage-analysis of families with inherited predisposition to cancer (Wooster *et al.*, 1995). Dietary and environmental factors also contribute in cancer formation (Doll and Peto, 1981; Houghton *et al.*, 2007). Cancer is proven to be a potentially fatal disorder that involves proliferation, transformation and deregulations of apoptosis. Cancer genes initiate extensive changes in cellular structure and cellular functions. These changes increase sensitivity to growth signals and insensitivity to growth-inhibitory signals,

decrease apoptosis and increase potential to replicate, generate angiogenesis and metastasis (Hanahan and Weinberg, 2000; Ichikawa *et al.*, 2007).

1.1.1 Oncogenes and tumor suppressor genes

Three types of genes are known to take part in development of cancer. The first type is oncogenes, is mutated or over expressed from normal proto-oncogenes. These are positive regulators of cell growth for example, *n-myc* in neuroblastoma, *c-erb-B2* in breast cancer and *k-ras* in pancreas cancer. The second type is known as tumor suppressor genes (TSGs). These are negative regulators of cell growth (Knudson, 1971). The third type is DNA repair genes (Cleaver, 1994). The tumor originates from the activation of oncogenes and the inactivation of tumor-suppressor genes.

Over hundred oncogenes have been identified and can be defined as gene-encoding products. The increased activity of oncogenes leads to increased proliferation. The deregulation of cancer genes results in a wide range of changes in cellular function and structure, all are contributing in various ways to malignancy (Hanahan and Weinberg, 2000). Some oncogenes, such as *ras*, *myc* and *erb* are identified as the human homologs of the viral transforming genes. Cellular genes become activated-oncogenes when these are incorporated into the viral genome (Harkin and Johnston, 2005).

1.1.2 Molecular natural cell cycle clock

A molecular natural clock instructs the cell when to replicate its DNA and when to divide. Some proteins regulate the timings of events occur in the cell cycle. The loss of control on timing leads towards cancer (Houghton *et al.*, 2007; Ichikawa *et al.*, 2007). A protein known as *cyclin* is the oscillator that runs the molecular clock. The cellular concentration of *cyclin* increases till the time of cell division and then suddenly decreases (Evan *et al.*, 1983). One specific enzyme known as *cyclin*-dependent kinase (cdk) also takes active part in the clock mechanism. Then another component protease (in the form of a proteasome) reset the clock (Glotzer *et al.*, 1991; King *et al.*, 1996). The cell cycle time for human tumors is around two days to several weeks (Wilson *et al.*, 1988; Basse *et al.*, 2002).

1.1.3 A brief history of treatment of cancer

In early days, the bacterial toxins were used in cancer treatment. It was based on host–tumor interactions, which opens the way to anti-angiogenesis and immune approaches. Modern research is developing a combination of complex strategies for growth and death control systems within the tumor cell. However, the development of low molecular-weight anticancer drugs and new strategies are likely to be continued. Currently the treatment for cancer ailment involves a combination of surgery, radiotherapy and chemotherapy. Natural products from plant origin play a very significant role in cancer therapy. One of the flowering plants, *Catharanthus roseus* is fundamentally cytotoxic and inhibits proliferation by exhibiting a very unique strategy, known as cell cycle specificity (Johnson *et al.*, 1963). A brief history of clinical cytotoxic drugs is summarized in the following Table 1.1

Table 1.1 A brief history of clinical cytotoxic drugs

No	Drug	Approximate Year of introduction	No	Drug	Approximate Year of introduction		
1	N Mustard	1948		WHO updated the list into three categories	1999		
2	Methotrexate	1953					
3	Chlorambucil	1956					
4	Thioguanine	1958					
5	Cyclophosphamide	1959					
6	Vinblastine	1960					
7	5-Fluorouracil	1961					
8	Vincristine	1962					
9	Melphalan	1964					
10	Daunorubicin	1965					
11	Cytarabine.	1970					
12	Bleomycin	1974	1			Bleomycin	1999
13	Doxorubicin	1976	2			Chlorambucil	
14	Mitomycin	1977	3			Cisplatin	
15	Dacarbazine	1978	4			Cyclophosphamide	
16	BCNU	1980	5			Cytarabine	
17	Cisplatin	1978	6			Dactinomycine	
18	Etoposide	1979	7			Daunorubicin	
19	Teniposide	1980	8			Doxorubicin	
20	Amsacrine	1981	9			Etoposide	
21	Carboplatin	1984	10			Fluorouracil	
22	Epirubicin	1987	11			Mercaptopurine	
23	Mitoxantrone	1989	12			Methotrexate	
24	Paclitaxel	1992	13			Prednisolone	
			14	Procarbazine			
			15	Tamoxifen			
			16	Vinblastine			
			17	Vincristine			
	WHO recognized this list of 24 essential drugs	1994		Category 2 12 drugs (See Martindale, 2007)	1999		
25	Gemcitabine	1995		Category 3 Non essential 13 drugs (See Martindale, 2007)	1999		
26	Docetaxel	1996					
27	Topotecan	1997					
28	Irinotecan	1998					

1.2 DRUG DISCOVERY OF NATURAL FLORA

Nature contributes a very significant role in medicinal therapy. It is likely to continue to be a source of new drugs in this modern era (Shah and Kaye, 2003). Natural compounds are complexes of well kept secrets of nature. Researchers are disclosing these wonders by using different techniques through available knowledge in the field of drug discovery to select out the compound of their interest. Current high throughput instrumental development had made drug discovery and development process more efficient (Drager, 2002; Houghton, 2002; McCally, 2002; Molyneux *et al.*, 2002; Stockigt *et al.*, 2002; Chen *et al.*, 2004; Balunas; King-Horn, 2006).

1.2.1 Lead structures of the synthetic compounds

Lead structures of most synthetic chemicals are based on structure of natural products (Mann, 2002). The supremacy of natural products over synthetic compounds is due to their colossal structural and chemical diversity. –About 40% of the chemical scaffolds found in natural products are not available in today’s medicinal chemistry” (Muller-Kuhrt, 2003). More than 50% of the drugs in clinical use are obtained from plant origin (Mann, 2002). According to the world health organization (WHO), up to 80% of people living in developing countries are following Traditional Medicine for their healthcare. People believe that majority of the natural products used are pharmacologically safer, more affordable and have in-built advantage that they hit multiple targets. (Kong and Liu, 2006)

1.2.2 “Natural Inhibitor of Carcinogenesis”

More than 10 countries have collaborated with the United States to conduct research under the project of “Natural Inhibitor of Carcinogenesis” and listed more than 250 compounds out of 5000 plant samples as potential cancer chemoprevention. The European Parliament passed a new legislation for European Union member nations to facilitate traditional medicine makers in terms of determination of efficacy. Canada opened a new Natural Health Products Directorate programme in January 2004. Pakistan, India, and Brazil are also developing botanical-drug research and testing centers (Kong and Liu, 2006). More than 2000 species grown in Malaysia are reported to have medicinal value (Jaganath and Ng, 2000). The climate of Malaysia is also suitable for the growth of *Catharanthus roseus* (*C. roseus*).

1.3 CATHARANTHUS ROSEUS (C. ROSEUS)

Catharanthus roseus is an ever blooming sub-tropical shrub. It was originally indigenous to Madagascar. It is now widely grown and used as indigenous medicine all over the world (Yarnell and Philhower, 2005).

1.3.1 Nomenclature of *Catharanthus roseus*

Its official name is *Catharanthus roseus* Linn G. Don, from family Apocynaceae (Leveque and Jehl, 2007; Magnotta *et al.*, 2007). Its synonym is *Vinca rosea* Linn, other included names are *Lochnera rosea* Reichb (Daniel, 2000), Periwinkle, Baramasi and Rattan-jot (Chopra *et al.*, 1956 and 1986). In Pakistan this plant is locally named as *Sadaa bahaar* (ever blooming). In Malaysia it is locally

known as *Kemuning cina* (Siddiqui, 2010). It produces potent anticancer *Vinca* alkaloids (Cutts *et al.*, 1960; Johnson, *et al.*, 1963).



Figure 1.1 *Catharanthus roseus*

1.3.2 Ethno-medical uses of *Catharanthus roseus*.

Most of the useful drugs derived from plants have been discovered by follow-up of the ethno medical uses (Farnsworth *et al.*, 1985; Dutcher *et al.*, 2000). In 1910, *Catharanthus roseus* was reported to be useful in Brazil as infusion of the leaves for mouth washes and toothaches, for the cleansing and healing of chronic wounds and also used in hemorrhage and scurvy (Synold, 2005; Johnson *et al.*, 1960). In British West Indies, it has been used to treat diabetic ulcers (Johnson *et al.*, 1960) In the Philippines; it has been used orally in the treatment of hyperglycemia and hypertension (Garcia, 1953).

In the forests of Madagascar, local people have been using decoction of the roots of *Catharanthus roseus* for the treatment of parasitic worms (Norscia and Borgognini-Tarli, 2006). Water extract of *Catharanthus roseus* was used for bleeding arrest, diabetes and fever or rheumatism (Ross, 2003). The leaves of the plant were chewed to suppress the sensations of hunger and fatigue (Ross, 2003; van der Heijden *et al.*, 2004; Ferreres *et al.*, 2008). The herbal and other preparations of *C. roseus* have been used for cancer and hypertensive treatment since ancient times (Hardman and Limbird, 2001).

1.4 VINCA ALKALOIDS AND OTHER COMPOUNDS PRESENT IN *CATHARANTHUS ROSEUS* (*C. ROSEUS*)

Antihypertensive/antiarrhythmic alkaloid ajmalicine and tranquillizer alkaloid serpentine and anticancer alkaloids vincristine and vinblastine isolated from *Catharanthus roseus* (*C. roseus*) were introduced as *Vinca* alkaloids (Hardman and Limbird, 2001). Developments in cancer treatment started in the 20th century (Baguley, 2002).

More than hundred indole *Vinca* alkaloids are reported from *Catharanthus roseus* (Heijden *et al.*, 2004; Hisiger and Jolicoeur, 2007). Twenty Indole-indoline dimeric alkaloids are more important. They are listed in Table 1.2. Only serpentine as tranquillizer (Iwase *et al.*, 2005), ajmalicine as anti-arrhythmic (Srivastava *et al.*, 2006) and vinblastine, vincristine and 3, 4- anhydrovinblastine as anti-cancer were marketed (Heijden, *et al.* 2004). Vindoline and catharanthine are the precursors in the biosynthetic pathways of dimeric indole alkaloids vinblastine and vincristine (Noble, 1990; Jolicoeur, 2007). *Catharanthus roseus* plant gains more interest due to its anticancer alkaloids (Sousa *et al.*, 2008). A list of anticancer alkaloids is mentioned bellow in Table 1.3.

Table 1.2 Important alkaloids reported from *Catharanthus roseus*

Alkaloids	Old name	Source	Empirical Formula
Vinblastine	Vincalokoblastine	Whole plant	$C_{46}H_{58}N_4O_9$
Vincristine	Leurocristine and Vincalurocristine	Arial parts	$C_{46}H_{56}N_4O_{10}$
Vindesine	Vindesine	Arial parts	$C_{43}H_{57}N_5O_{11}S$
Leurosine	Leurosine	Arial parts	$C_{46}H_{58}N_4O_9$
Ajmalicine	Raubasine	Flowers	$C_{21}H_{24}N_2O_2$
Lochnerine	Lochnerine	Flowers	$C_{20}H_{24}N_2O_2$
Lochnericine	Lochnericine	Flowers	$C_{21}H_{24}N_2O_2$
Catharanthine	Catharanthine	Flowers	$C_{21}H_{24}N_2O_2$
Serpentine	Serpentine	Flowers	$C_{21}H_{22}N_2O_3$
Vindoline	Vindoline	Arial parts	$C_{25}H_{32}N_2O_6$
Vindolinine	Vindolinine	Arial parts	$C_{21}H_{24-6}N_2O_2$
Reserpine	Reserpine	Arial parts	$C_{23}H_{40}N_2O_9$

(Johnson, *et al.*, 1960; Daniel, 2000; British Pharmacopoeia, 2000; Dutta *et al.*, 2005)

Table 1.3 Frequently used salts of anticancer alkaloids from *Catharanthus*

roseus

Salts of anticancer alkaloids	Brand name	Molecular weight of salt	Empirical formula
Vinblastine sulfate	Exal, Velban, Velbe, Velsar	909.1 (sulfate)	$C_{46}H_{58}N_4O_9 \cdot H_2SO_4$
Vincristine sulfate	Oncovine, Vincasar, Kyocristine, Vincosid, Vincrex	923 (sulfate)	$C_{46}H_{56}N_4O_{10} \cdot H_2SO_4$
Vindesine sulfate	Eldistine, Fildesine	852 (sulfate)	$C_{43}H_{57}N_5O_{11}S$
Vinorelbine ditartrate	Navelbine	1079 (di tartrate)	$C_{45}H_{54}N_4O_8 \cdot 2C_4H_6O_6$

(Johnson, *et al.*, 1960; Daniel, 2000; British Pharmacopoeia, 2000; Dutta *et al.*, 2005)

The anticancer activity of the *C. roseus* is mainly dependent on alkaloids such as vincristine, vinblastine, vindesine and vinorelbine, but the presence of phenols (Piovan and Filippini, 2007; Mustafa and Verpoorte, 2007; Ferreres *et al.*, 2008) and antioxidants (Misra and Gupta, 2006; Abduljaleel and Gopi, 2008) also contribute highly towards anticancer activity. The known compounds other than alkaloids and their mechanism of action are discussed in Section 1.10.2.

1.4.1 Discovery of anticancer alkaloids from *C. roseus*

Vinca alkaloids were isolated from *C. roseus* leaves in the late 1950s independently by two groups; Professor Noble working at the Western University of Ontario, Canada, and the other researchers at the Eli Lilly research laboratories in Indianapolis, USA (Noble, 1958 and 1990; van der Heijden *et al.*, 2004). *Vinca* alkaloids were scientifically studied initially as oral hypoglycemic agents (Johnson *et al.*, 1963).

Robert Noble was working in Canada on some hypoglycemic oral compound found in *Catharanthus roseus*, (*C. roseus*) not grown in Canada (Tailor and Farnsworth, 1975). *C. roseus* extract solution when injected into mice caused immune suppression (Noble *et al.*, 1958; Duffin, 2002). It indicated the extract can induce pronounced activity without any toxic effect (Johnson *et al.*, 1963; Yarnell and Philhower, 2005).

1.4.2 Earliest chemotherapeutic agents

Catharanthus roseus is the only source for vinblastine and vincristine. *Catharanthus roseus* produced around 130 indole alkaloids, but a full picture of its biosynthetic pathway has yet to be elucidated (Zeffrin, 1984; El-Sayed and Verpoorte, 2007). Before 1961, vinblastine was the only anticancer drug commercially available for cancer treatment from *C. roseus*. In March 1961, Eli Lilly announced the commercial availability of a new anticancer compound called leurocristine, which was the former name of vincristine (Johnson *et al.*, 1960). Vinblastine and vincristine are among the earliest agents developed for cancer therapy. They are approved for cancer treatment for more than 30 years and remain widely used in anticancer therapy.

1.4.3 Cancers treated by *Vinca* alkaloids

Vinca alkaloids and their derivatives act as mitotic inhibitors (Rowinsky and Donhower, 1996). Vinblastine and vincristine have been used as chemotherapeutic agents in the treatment of a wide range of tumors, particularly childhood tumors, leukemia and lymphomas (Riyaz and Stanley 2003; Synold, 2005; Levêque and Jehl, 2007), Hodgkin's disease (Svoboda, *et al.*, 1959; Levêque and Jehl, 2007) and testicular teratoma (Mann, 2002; Synold, 2005). It is also effective in treating breast carcinoma, nephroblastoma, brain tumors, lung cancer, leiomyosarcoma, cervix cancer, alimentary tract tumor (Pui and Evans, 1998) and Kaposi's sarcoma (Kaplan *et.al.*, 1986; Gill *et al.*, 1991; Tulpule and Matheny, 1998). Different cancers treated by *Vinca* alkaloids are listed in Table 1.4

Table 1.4 Cancers treated by *Vinca* alkaloids.

Alkaloid	Uses
Vinblastine Mitotic inhibitor	Leukemia, Hodgkin's lymphoma and germ cell cancer, Kaposi's sarcoma, bladder, breast and lung cancer.
Vincristine Topoisomerase inhibitors	Ovarian cancer, leukemia, Hodgkin's lymphoma, Wilm's tumor, multiple myeloma, breast and lung cancer.
Vinorelbine (Navalbine)	Breast, non-small cell lung cancer, gastrointestinal cancer
Vinorelbine Phase-II clinical trials	Phase-II clinical trials bladder, non-small cell lung carcinoma and breast cancers.

(Mac Lenna and Cusack, 1985; Boccardo *et al.*, 1989; Daniel, 2000; British Pharmacopoeia, 2002; Extra Pharmacopoeia, 2002; Ichikawa *et al.*, 2007)


1.4.4 Sulfate and ditartrate salts of *Vinca* alkaloids

In treatment of the cancer, either of the *Vinca* alkaloids is prescribed in combination with the other cytotoxic drugs. A collection of alkaloids from *Catharanthus roseus* (*C. roseus*) is presented in Table 1.2. *C. roseus* produced various monoterpenoid indole alkaloids (MIAs) (Ichikawa *et al.*, 2007). They are utilized in salt form as listed in Table 1.3. The sulfate salts of vinblastine, vincristine and vindesine are administered via intravenous route. Vinorelbine ditartrate salt (Novelbine) is administered orally (Degardin *et al.*, 1994; Extra Pharmacopoeia, 2002). A semi synthetic fluorinated *Vinca* alkaloid is produced through change in the catharanthine moiety of vinblastine (Kruczynski and Hill, 2001; Jacquesy, 2006).

1.5 STRUCTURE OF VINCA ALKALOIDS

A group of researchers at Eli Lilly discovered the structures and revealed the clinical activity of vincristine and vinblastine (Edwards, 1994). *Vinca* alkaloids are identical in structure with difference in the group attached to the nitrogen at position 1 at which vincristine possesses a labile N- formyl group and vinblastine has a stable methyl group Figure 1.2. D ring is a small portion of *Vinca* molecule which is extremely sensitive in term of overall biological properties and tubulin interactions (Jacques, 2001).

Fig 1.2 Chemical structure of *Vinca* alkaloids



Alkaloids	R1	R2	R3
Vinblastine	-CH ₃	-CO ₂ CH ₃	-OCOCH ₃
Vincristine	-CHO	-CO ₂ CH ₃	-OCOCH ₃
Vindesine	-CH ₃	-CONH ₂	-OH
Vinorelbine	-CH ₃	-CO ₂ CH ₃	-OCOCH ₃

1.5.1 Structural difference and clinical activity

There are minor structural differences present between anticancer *Vinca* alkaloids and they behave in the same way with the drug-tubulin interaction but their toxicity and spectrum of clinical activity is significantly different from each other. (Himes *et al.*, 1976; Owellen *et al.*, 1977)

1.5.2 Derivatives of *Vinca* alkaloids

In combination with other anticancer therapies, vinblastine and vincristine are given for a curative purpose (Levêque & Jehl, 2007). From vinblastine (Oncovin) and vincristine (Velbe), a large number of derivatives have been formed. They are modified in the vindoline moiety (mido derivative). Vindesine (deacetylvinblastine amide) (Barnett *et al.*, 1978) was registered in Europe in 1980. Vindesine is not approved in the United States, and in France. It is used in the treatment of aggressive forms of non-Hodgkin lymphomas in combination with other anticancer drugs.

In 1980s a semi synthetic derivative is introduced as vinorelbine nor-5'-anhydrovinblastine (Navelbine). It is invented by the pharmacist Pierre Potier and his team in France, produced by changing the catharanthine moiety of vinblastine. Vinorelbine is given in palliative treatment of advanced nonsmall-cell lung cancer, refractory lymphoid malignancies and advanced breast cancer. Vinorelbine gives greater therapeutic activity when given as adjuvant (or postoperative) treatment associated with cisplatin in patients with resected non-small-cell lung cancer (Winton *et al.*, 2005; Douillard *et al.*, 2006).

Vinflunine is a fluorinated derivative of vinorelbine. It is in the process of clinical development (Kruczynski & Hill, 2001; Bennouma *et al.*, 2005). The substitution of fluorine in the structure of natural products has been proven beneficial. The fluorination increases the lipophilic profile of a molecule, which can be more effective (Thomas, 2006). Novel vinflunine (Javlor) is under clinical trial (Kruczynski and Hill, 2001; Jacquesy, 2006) and advanced to the phase III clinical trials (Shnyder, 2004). Vinflunine hold a potent antiangiogenic effect (Holwell *et al.*, 2001).

1.6 MODE OF ACTION OF VINCA ALKALOIDS

The mechanism of antineoplastic activity is worked during cell mitosis (George *et al.*, 1965). *Vinca* alkaloids are mitotic inhibitors (Himes, 1991; da Rocha *et al.*, 2001; Honore *et al.*, 2003). These agents target microtubules. They are the structural backbone of both normal and abnormal cells. Microtubules are polymers of long tube shaped dynamic structures. They are constantly growing and shortening (Mitchison and Kirachner, 1984).

Microtubules contribute essential roles in construction and function of the mitotic spindle. They actively take part in many cellular events, including cellular organization, cell division, intracellular transport, intracellular transfer of signals, neurotransmission, and the transmission of signals from the cell-surface-receptors to the nucleus (Dutcher and Novik, 2000).

1.6.1 Target of *Vinca* alkaloids for anticancer activity

Tubulin is one of the essential proteins for chromosomal segregation. The tubulin- microtubule system is an important target for anticancer therapy (Jordan and Hadfield, 1998; Houghton, 2002). Hydrophobic *Vinca* alkaloids bind to tubulin in a reversible manner (Zhou and Rahmani, 1992). The *Vinca* alkaloids inhibit the microtubule polymerization at high non-physiologically relevant concentrations (Jordan *et al.*, 1991; Dhamodharan, *et al.*, 1995). *Vinca* alkaloids act by binding to the micro-tubular proteins of the spindle and arresting mitosis at the metaphase/anaphase transitions leading to apoptosis (Nagan, 2000; Kruczynski *et al.* 2002).


The mechanism of the mitotic blockage is not well known (Okouneva *et al.*, 2005; Weaver and Cleveland, 2005), the *Vinca* alkaloids interfere with the function of microtubules in axons, which mediates the neuronal vesicle transport (Dumontet *et al.*, 1999; Bacher *et al.*, 2001). The research is performed on an array of chemically diverse antimitotic and microtubule targeting agents. They depolymerize microtubules of newly developed vascular system and stop the blood supply to tumor (Okuneva *et al.*, 2003; Honore *et al.*, 2003; Jordan and Wilson, 2004). The function of *Vinca* alkaloids as angiogenesis agent is discussed in chapter 6.

1.6.2 *Vinca* alkaloids as mitotic inhibitors during cell cycle

There are two prominent phases occur in mammalian cell division. One is known as synthetic or S phase and the other is mitosis or M phase. There are two gaps G1 and G2 occur between S phase and M phase. *Vinca* alkaloids function as mitotic inhibitors between G2 and M phase of the cell cycle. (Musunru and Hinds,

1997) The transitions from *G1*-phase to *S*-phase, and from *G2*-phase to *M*-phase, are controlled by a system of cyclin-dependent kinases (cdk), cyclins and phosphatases (Houghton *et al.*, 2007; Ichikawa *et al.*, 2007). The cell cycle clock is demonstrated in Figure 3.

Figure 1.3 The cell cycle clock



S = synthetic

G = gap

M = mitosis

The known mechanism of drug action is to arrest cells at the metaphase by interfering with the assembly or disassembly of α - and β - tubulin into microtubules and inhibits tubulin polymerization (Rowinsky and Donhower, 1996). These drugs target β -tubulin subunit of α -/ β - tubulin heterodimers, inhibiting the addition of heterodimers onto growing microtubules, hence, preventing polymerization of microtubules (Musunru and Hinds, 1997; Jordan and Wilson, 1998).

1.6.3 Activity of *Vinca* alkaloids in high and low concentration

At high concentrations, the *Vinca* alkaloids cause complete microtubule de-polymerization. At low concentrations de-polymerization does not occur but there is sufficient alteration in the dynamics of tubulin loss or addition at the ends of mitotic spindle. It prevents the spindle from carrying out its function of attaching to and segregating the chromosomes, and cause cell arrest during mitosis (Jordan *et al.*, 1992; Jordan, 2002; Jordan and Wilson, 2004). Prolonged arrest finally leads to cell death, either in mitosis or after an ultimate escape from mitotic arrest (Jordan and Wilson, 1998).

Lower concentrations of microtubule-targeted drugs can suppress dynamics of microtubule without changing mass of microtubule (Okouneva *et al.*, 2003). Tubulin binding and disruption of the cell membranes simultaneously occur due to interference with the lipid bi-layer at the same concentration ((Rowinsky and Donhower, 1996).

1.6.4 Special *Vinca*-specific high-affinity and low-affinity sites

Each heterodimer of tubulin possesses special *Vinca*-specific high-affinity and low-affinity sites. These binding sites are different from the interacting sites of other drugs such as taxanes (Rao *et al.*, 1992; Rao and Krauss *et al.*, 1994). Binding to the high affinity sites decreases the rate of dissociation and association of tubulin dimers of the microtubules (Jordan *et al.* 1986). Binding to the low-affinity sites appears to be responsible for disruption of the microtubule configuration, leading to disintegration (Jordan *et al.*, 1986).

Despite the above mentioned promising anticancer activity, these alkaloids have many problems during their uses. The drug resistance, toxicity, and low specificity are major difficulties in the treatment of cancer (de Mesquita *et al.*, 2009). A brief account on these problems is discussed as follows.

1.7 CELL RESISTANCE AGAINST *VINCA* ALKALOIDS

The efficacy or activity of *Vinca* alkaloids decreases when cancer cells develop resistance against anticancer drugs (Dumontet and Sikic, 1999). When cancer cells develop clinical resistance to one drug, the simultaneous resistance to several other structurally and mechanistically unrelated drugs is also induced. Clinical resistance reduces the concentration of the drug in the target cell, which in consequence reduces activity and decreases the clinical effectiveness of drug (Husain and Wozniak, 1993; Hunter, 1997; ACPE, 2002).

1.7.1 Classical multiple drug resistance (MDR) of cancer cells to natural hydrophobic drug

Vinca alkaloids are comparatively more hydrophobic in nature. Resistance development of cancer cells to natural hydrophobic drugs are known as classical multi drug resistance. Study of cancer cells in culture with vinblastine, paclitaxel or doxorubicin frequently results in multi drug resistance due to expression of ATP-dependent efflux pumps with broad drug specificity. Family of ATP-binding cassette (ABC) transporters is responsible for efflux pumps. It is further divided into seven subfamilies (ABCA-ABCG). The ABC transporter P-glycoprotein (P-gp) is the product of the ABCB1 or *mdr1*-gene (Tsuruo *et al.*, 1972; Kolars *et al.*, 1992; Gottesman *et al.*, 2002).

There are different mechanisms of resistance including P-glycoprotein efflux pump and multi drug resistance protein (MRP). These proteins are transmembrane transporters which are also responsible for the rapid efflux of intracellular chemotherapeutic agents. Over expression of these proteins indicate the less concentration of intracellular *Vinca* alkaloids and reduction in cytotoxicity. In brief, the development of multiple drug resistance (MDR) is common for *Vinca* alkaloids due to the following reasons:

1. Over expression of transmembrane efflux pump system known as P-glycoprotein (P-gp) (Cole and Deeley, 1998; Bardelmeijer *et al.*, 2000; Bacher and Nickel, 2001).
2. The action of multi-drug resistance-associated proteins (MRP) (Fardel *et al.*, 1996; Cole and Deeley, 1998; Bardelmeijer *et al.*, 2000; Bacher and Nickel, 2001).