













16. Halazonetis DJ. Morphometrics for cephalometric diagnosis. *Am J Orthod Dentofacial Orthop* 125: 571-581, 2004
17. Silverman FN. Roentgen standards for size of the pituitary fossa from infancy through adolescence. *Am J Roentgenol Radium Ther Nucl Med* 78: 45-60, 1957
18. Axelsson S, Storhaug K and Kjaer I. Post-natal size and morphology of the sella turcica-longitudinal cephalometric standards for Norwegians between 6 and 21 years of age. *Eur J Orthod* 26: 597-604, 2004
19. Alkofide EA. The shape and size of the sella turcica in skeletal Class I, Class II, and Class III Saudi subjects. *Eur J Orthod* 29: 457-463, 2007
20. Tetradis S and Kantor ML. Prevalence of skeletal and dental anomalies and normal variants seen in cephalometric and other radiographs of orthodontic patients. *Am J Orthod Dentofacial Orthop* 116: 572-577, 1999
21. Argyropoulou M, Perignon F, Brunelle F, Brauner R and Rappaport R. Height of normal pituitary gland as a function of age evaluated by magnetic resonance imaging in children. *Pediatr Radiol* 21: 247-249, 1991
22. Chilton LA, Dorst JP and Garn SM. The volume of the sella turcica in children: new standards. *Am J Roentgenology* 140:797 – 801, 1983
23. Shah AM, Bashir U and Ilyas T. The shape and size of the sella turcica in skeletal class I, II & III in patients presenting at Islamic International Dental Hospital, Islamabad. *Pakistan Oral Dent J* 31: 104-110, 2011
24. Ruiz CR, Wafae N and Wafae GC. Sella turcica morphometry using computed tomography. *Eur J Anat* 12: 47-50, 2008
25. Valizadeh S, Shahbeig S, Mohseni S, Azimi F and Bakhshandeh H. Correlation of shape and size of sella turcica with the type of facial skeletal class in an iranian group. *Iran J Radiol* 12: 1-7, 2015
26. Sathyanarayana HP, Kailasam V and Chitharanjan AB. The size and morphology of sella turcica in different skeletal pattern among South Indian population. *J Indian Orthod Soc* 47: 266-271, 2013
27. Osunwoke EA, Mokwe CR and Amah-Tariah FS. Radiologic measurements of the sella turcica in an adult Nigerian population. *Int J Clin Pharmacol Res* 4: 115-117, 2014
28. Marsan G and Öztaş E. Incidence of bridging and dimensions of sella turcica in class I and III Turkish adult female patients. *World J Orthod* 10: 99-103, 2009

