



# Asian Pacific Journal of Tropical Biomedicine

## Original Article



doi: 10.4103/2221-1691.260397

www.apjtb.org

## Methanolic extract of *Abrus precatorius* promotes breast cancer MDA-MB-231 cell death by inducing cell cycle arrest at G<sub>0</sub>/G<sub>1</sub> and upregulating Bax

Wan Suriyani Wan-Ibrahim<sup>1</sup>, Norzila Ismail<sup>1</sup>✉, Siti Farhanah Mohd-Salleh<sup>1</sup>, Aidy Irman Yajid<sup>2</sup>, Michael Pak-Kai Wong<sup>3</sup>, Mohd Nizam Md Hashim<sup>3</sup>

<sup>1</sup>Department of Pharmacology, School of Medical Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

<sup>2</sup>Department of Pathology, School of Medical Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

<sup>3</sup>Department of Surgery, School of Medical Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia

### ARTICLE INFO

#### Article history:

Received 5 March 2019

Revision 18 April 2019

Accepted 10 June 2019

Available online 17 June 2019

#### Keywords:

*Abrus precatorius*

MDA-MB-231

Apoptosis

Cell cycle

Breast cancer

### ABSTRACT

**Objective:** To determine the anti-proliferative activity of *Abrus precatorius* (*A. precatorius*) leaf extracts and their effect on cell death.

**Methods:** *A. precatorius* leaves were extracted successively with hexane, ethyl acetate and methanol by Soxhlet extraction. Aqueous extract was prepared by decoction at 50 °C. Extracts of *A. precatorius* leaves were used to treat selected cancer and normal cell lines for 72 h. Furthermore, 3-(4,5-dimethyl thiazol-2-yl) 2,5-diphenyl tetrazolium bromide assay was performed to determine cell viability. Analysis of cell cycle arrest, apoptosis assay and apoptosis protein expressions were determined by flow cytometry.

**Results:** Methanolic extract of *A. precatorius* leaves showed the lowest IC<sub>50</sub> on MDA-MB-231 cells at (26.40±5.40) µg/mL. Flow cytometry analysis revealed that cell arrest occurred at G<sub>0</sub>/G<sub>1</sub> phase and the apoptosis assay showed the occurrence of early apoptosis at 48 h in MDA-MB-231 cells treated with methanolic extract of *A. precatorius* leaves. Methanolic extract of *A. precatorius* leaves induced apoptosis by upregulation of Bax, p53 and caspase-3 and downregulation of Bcl-2.

**Conclusions:** Methanolic extract of *A. precatorius* leaves promotes MDA-MB-231 cell death by inducing cell cycle arrest and apoptosis possibly *via* the mitochondrial-related pathway.

## 1. Introduction

Apoptosis is a characterized form of cell death and is mostly studied. Known commonly as programmed cell death, apoptosis is the packaging of dying cells into fragments that are easily consumed and eliminated by phagocytes without disturbing the normal function of surrounding tissues[1]. Equilibrium between cell death and cell proliferation is important to avoid disruption of the cellular balance.

Excessive apoptosis or deficient apoptosis is the cause of many clinical diseases including cancer[2]. Apoptosis can be initiated through two separate pathways, the intrinsic or mitochondrial pathway and the extrinsic or death receptor pathway. In cancer management, apoptosis has become an important tool as a target by potent apoptosis-inducing agents, including both chemical and biological[3].

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** reprints@medknow.com

©2019 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow. All rights reserved.

**How to cite this article:** Wan-Ibrahim WS, Ismail N, Mohd-Salleh SF, Yajid AI, Wong MPK, Md Hashim MN. Methanolic extract of *Abrus precatorius* promotes breast cancer MDA-MB-231 cell death by inducing cell cycle arrest at G<sub>0</sub>/G<sub>1</sub> and upregulating Bax. Asian Pac J Trop Biomed 2019; 9(6): 249-256.

✉Corresponding author: Dr. Norzila Ismail, Department of Pharmacology, School of Medical Sciences, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia.

Tel: +6097676135

Fax: +6097653370

E-mail: norzila\_ismail@usm.my

Funding: This study was funded by the Universiti Sains Malaysia Short Term Grant (304/PPSP/61313046).

Cancer prevalence is increasing and has been one of the main causes of death. World Health Organization[4] reported that cancer is the second leading cause of death worldwide with an estimated 9.6 million of deaths in 2018, in which 1 in 6 deaths is caused by cancer. The leading cancers causing deaths include lung, colorectal, stomach and breast cancer. About 627 000 death cases were reported due to breast cancer globally. In Malaysia, 43 837 new cases were reported in 2018 with breast cancer amounting to 7 593 cases[5]. Advances in the medical field have proven to be the preferable choice to combat cancer. However, these advances are used with concern due to their side effects and limitations. This scenario has generated increasing demand of traditional medicines with medicinal plants either as complementary to the allopathy treatments or as a complete alternative. Many issues were raised with this development especially concerning the toxicity and efficacy of the medicinal plants extract. Therefore, it is noted that more studies are needed in order to provide better understanding of the biological activities and underlying mechanism of the medicinal plants.

Medicinal plants are widely sought as an alternative in various treatments including cancer. These plants are extensively studied to screen for potential phytochemical as anticancer agents. *Abrus precatorius* (*A. precatorius*) is a flowering plant that belongs to the legume family, Fabaceae. This plant is native to India, however, it is also found in other parts of tropical areas, like Malaysia[6]. This plant is phenotypically recognized by its slender, perennial, climbing twigs with pinnate leaves arranged in pairs. *A. precatorius* seeds are red with black dot. In Malaysia, the leaves of *A. precatorius* are traditionally used to treat simple illness such as mouth ulcer and fever. Moreover, *A. precatorius* is also reported to be used as an anticancer agent[7–13].

Gul et al[7,12] reported anti-proliferative activities of *A. precatorius* against human acute monocytic leukemia cell line (THP-1), while Sofi et al[8,13] reported its anti-proliferative activities against MDA-MB-231 by using aqueous extract and fractions from gradient elution of ethyl acetate extract. Our previous study[6] firstly reported the phytochemical analysis of this plant in Malaysia. Therefore, this study aimed to further elucidate the anticancer properties of *A. precatorius* leaves extracts on cancer cell lines and investigate their effect on cancer cell death.

## 2. Materials and methods

### 2.1. Plants collection and extract preparation

*A. precatorius* leaves were collected from Kampung Sabak, Pengkalan Chepa Kelantan and given a specimen voucher USM 11730 by the Herbarium Unit, School of Biological Sciences, Universiti Sains Malaysia for future references. About 22 g of ground *A. precatorius* were subjected to successive Soxhlet extraction with hexane, ethyl acetate and methanol. Aqueous extract was

prepared by decoction of the ground leaves at 50 °C in 500 mL of distilled water until the water was reduced to a third of the initial volume. All extracts were kept at -20 °C until use.

### 2.2. Cell culture

Human breast cancer cell lines, MDA-MB-231 and MCF-7; human liver cancer cell lines, HepG2; human colon cancer cell lines, SW480; human cervical cancer cell line, SiHa and HeLa; human normal breast cell, MCF-10a and mouse normal fibroblast cell, NIH were obtained from American Type Cell Culture Collection, Maryland, USA. Cells were seeded in 25 cm<sup>2</sup> tissue culture and grown at 37 °C under humidified 5% CO<sub>2</sub> in Dulbecco's Modified Eagle's medium supplemented with 5% fetal bovine serum and 1% penicillin-streptomycin. Confluent cells were harvested by trypsinization (0.25%).

### 2.3. Anti-proliferative activity by 3-(4,5-dimethyl thiazol-2-yl) 2,5-diphenyl tetrazolium bromide (MTT) assay

To determine the anti-proliferative activity of *A. precatorius*, human breast cancer cell lines, MDA-MB-231 and MCF-7; human liver cancer cell lines, HepG2; human colon cancer cell lines, SW480; human cervical cancer cell line, SiHa and HeLa; human normal breast cell, MCF-10a and mouse normal fibroblast cell, NIH were treated with all extracts. The anti-proliferative activities were investigated by MTT assay. IC<sub>50</sub> values were determined, with lower IC<sub>50</sub> values indicating higher anti-proliferative activity. Cytotoxicity of plant extracts against cancer cell line was categorized based on US National Cancer Institute and Geran Protocol as follows: highly cytotoxic (IC<sub>50</sub> ≤ 20 µg/mL), moderately cytotoxic (21 µg/mL ≤ IC<sub>50</sub> ≤ 200 µg/mL), weakly cytotoxic (201 µg/mL ≤ IC<sub>50</sub> ≤ 500 µg/mL), and, no cytotoxicity (IC<sub>50</sub> ≥ 501 µg/mL)[14].

Cells were seeded into 60 wells at the centre of a 96-wells plate with the concentration of 5 × 10<sup>4</sup> cells/mL per well. Extracts of *A. precatorius* leaves were added following a serial dilution starting from 99 µg/mL until 0.39 µg/mL in each well. Anti-proliferative activity of *A. precatorius* extracts was measured by the MTT assay, which was performed after a 72-h incubation post treatment with the extracts, tamoxifen (positive control) and dimethyl sulfoxide (negative control). Absorbance was read at OD of 570 nm. The absorption value at this wavelength directly represents the relative cell numbers in comparison with the control group[15]. The percentage of cell viability was determined according to the following equation: Percentage of cell viability (%) = Absorbance of treated cells (extracts or tamoxifen)/Absorbance of treated cells (dimethyl sulfoxide) × 100

Anti-proliferative activity of *A. precatorius* leaves extract was screened on selected cancer and normal cells. Extract that exhibited the lowest IC<sub>50</sub> value with its corresponding cell was used for subsequent analysis in this study.

**Table 1.** IC<sub>50</sub> values of *A. precatorius* leaf extracts against selected normal and cancer cell lines (µg/mL).

Types of cancer cell lines	Hexane extract	Ethyl acetate extract	Methanol extract	Aqueous extract	Tamoxifen
HeLa (cervix)	>99	>99	73.60±6.17	>99	4.32±0.40
SiHa (cervix)	>99	>99	>99	>99	3.27±0.93
MCF7 (breast)	52.65±7.14	99.00±11.86	59.03±9.40	>99	1.81±1.78
MDA-MB-231 (breast)	45.60±11.60	54.50±9.05	26.40±5.40	>99	2.27±0.38
SW 480 (colon)	>99	>99	77.23±6.39	>99	2.31±0.59
HepG2 (liver)	>99	99.00±10.32	67.72±6.21	>99	4.10±0.88
MCF-10a (normal breast)	>99	>99	>99	>99	3.78±7.79
NIH (normal fibroblast)	>99	>99	>99	>99	3.78±1.78

## 2.4. Cell cycle assays

MDA-MB-231 cells were treated with IC<sub>50</sub> of the methanolic extract of *A. precatorius* leaves (APME) and incubated for 24, 48 and 72 h. Cells were harvested by trypsinization and the cell cycle assay was performed according to the manufacturer protocol, BD Cycletest™. All samples readings were acquired with FACSCANTO II (BD Bioscience). Data obtained was analysed with ModFit LT 5.0 software.

## 2.5. Apoptosis assays

### 2.5.1. Annexin-V and propidium iodide (PI) staining

MDA-MB-231 cells were treated with IC<sub>50</sub> of the APME and incubated for 24, 48 and 72 h. Cells were harvested by trypsinization and the apoptosis assay was performed according to the manufacturer protocol, Annexin V-FITC detection kit I (BD Bioscience). All samples readings were acquired with FACSCANTO II (BD Bioscience). Data was analysed with FlowJo software.

### 2.5.2. Bax, Bcl-2, caspase-3 and p53 activity

MDA-MB-231 cells were treated with IC<sub>50</sub> of the APME and incubated for 24, 48 and 72 h. Cells were harvested by trypsinisation following each incubation time and washed twice with phosphate buffer saline (PBS). Ethanol (70%) was used to fix the cells at 4 °C for 1 h. Cells were washed twice with PBS and then blocked with 2% bovine serum albumin for 10 min at room temperature. Another cell wash was performed, and cells were resuspended in PBS.

About 100 µL of the cell suspension (1 × 10<sup>6</sup> cells) were mixed independently in different tubes, with antibodies (SantaCruz); Bax-PE (sc-7480), Bcl-2 – Alexa Fluor 647 (sc-7382), p53 – Alexa Fluor 488 (sc-126) and Caspase-3 – Alexa Fluor 488 (sc-7272). These cells-antibodies mixtures were incubated for 20 min at room temperature, then washed once and resuspended in 500 µL PBS. All samples readings were acquired with FACSCANTO II (BD Bioscience). Data was analysed with FlowJo software.

## 2.6. Statistical analysis

The data were expressed as mean ± SD of three repeated experiments. The level of statistical significance was tested using repeated measure one-way ANOVA, followed by Dunnett's multiple comparison test. The difference was considered significant if  $P < 0.05$ .

Analyses were all done using GraphPad Prism7.


## 3. Results

### 3.1. Screening of *A. precatorius* extracts on various cancer and normal cells

The IC<sub>50</sub> values are summarized in Table 1. Methanol extract had the lowest IC<sub>50</sub> value on MDA-MB-231 cells at (26.40±5.40) µg/mL, which could be categorized to moderate toxicity. IC<sub>50</sub> values of all extracts were determined by plotting the graph of concentration of the extract or tamoxifen versus percentage of cell viability as depicted by Figure 1 which represents the data for MDA-MB-231 cells.

### 3.2. Effects of APME on cell cycle progression in MDA-MB-231 cells


The ability of the APME to reduce cell viability could be due to cell death mediated by cell cycle arrest. Therefore, cell cycle arrest was investigated using flow cytometric analysis of PI stained DNA. Representative profiles of the cell cycle progression are presented in Figure 2(a) and percentage of each phase of cell cycle is presented in Figure 2(b). Significant increment of the arrest was found at G<sub>0</sub>/G<sub>1</sub> phase of the cell cycle, suggestive of cell death consistent with the reduction of cell growth as the percentage of cell population decreased in both S-phase and G<sub>2</sub>/M phase.



**Figure 1.** Antiproliferative effects of all extracts on MDA-MB-231 cells. Data were expressed as mean±SD of three repeated experiments with three replicates.

### 3.3. Effects of APME on apoptosis in MDA-MB-231 cells

To determine the ability of the APME to induce apoptosis, an apoptosis assay was performed using the Annexin V-FITC detection kit I (BD Bioscience). Representative profiles of the apoptosis assays are presented in Figure 3(a) and percentage of each phase of cell death progression is presented in Figure 3(b). Live cells did not uptake any stain and represented at Q4 (Figure 3a). Annexin V bound to the phosphatidylserine of the plasma membrane which was exposed in early apoptosis (Q3; annexin V positive, PI negative). Late apoptotic cells lost their cell integrity thus allowing the penetration of PI (Q2; annexin V positive, PI positive), while necrotic cells were stained with PI only (Q1 PI positive). APME induced early apoptosis in MDA-MB-231 cells at 48 h and late apoptosis at 72 h after treatment.



**Figure 2.** Effects of methanolic extract of *A. precatorius* leaves on cell cycle progression in MDA-MB-231 cells. (a) Cell cycle arrest analysis of MDA-MB-231 cells; (b) percentage of each phase of cell cycle. Data are expressed as mean±SD of three repeated experiments. \* $P < 0.05$  is considered significant when comparing treated cells vs untreated cells.

### 3.4. Bax, Bcl-2, caspase-3 and p53 protein expression in MDA-MB-231 cells treated with APME


To determine whether Bax, Bcl-2, caspase-3 and p53 proteins are involved in APME-induced apoptosis in MDA-MB-231 cells, the proteins expressions were measured by flow cytometry at 24, 48, and 72 h (Figure 4a). The MDA-MB-231 cells treated with the APME increased pro-apoptotic protein, Bax and reduced anti-apoptotic protein, Bcl-2 expressions in a time dependent manner (Figure 4b). Significant increase was shown in the p53 protein expressions in the treated cells compared with untreated ones. Caspase-3 protein expressions were also increased. These findings indicated that APME induced apoptosis in MDA-MB-231 cells by upregulating Bax, p53 and caspase-3 protein and downregulating Bcl-2 protein. These findings suggest that the apoptosis in MDA-MB-231 cells was possibly triggered in the intrinsic pathway.

## 4. Discussion

Medicinal plants have been actively studied over the years in research labs and used traditionally worldwide. *A. precatorius* is one of the medicinal plants that belongs to the legume family Fabaceae. The legume family has many plants that exhibit excellent anticancer properties[16–19].

Anti-proliferative study gives an insight on the possibility of an occurrence of cell death of selected cancer cells induced by the extract. Our study demonstrated that the methanol leaves extract had the best anti-proliferative activity at 26.4 µg/mL against the human breast cancer cells, MDA-MB-231, an androgen-independent human breast cancer cells that express wild-type p53. However, these values are not comparable to the control, tamoxifen, which demonstrated the IC<sub>50</sub> values of 2.27 µg/mL in MDA-MB-231. Based on the National Cancer Institute criteria, these results indicated that APME has a moderate cytotoxicity on MDA-MB-231 cell lines. Sofi *et al*[13] demonstrated that the IC<sub>50</sub> of the aqueous extract on MDA-MB-231 cells was 98 µg/mL. Our results showed better inhibitory activity on MDA-MB-231 cells by treatment with the APME. The aqueous leaves extract of our experiment did not show any significant activity at maximum concentration of 99 µg/mL. Another study by Gul *et al*[12] claimed the hexanoic and ethanolic extracts of *A. precatorius* leaves had stronger anti-proliferative activity both in human colon adenocarcinoma cells (Colo-205) and human retinoblastoma cancer cells (Y79), while milder anti-proliferative activities were observed in human hepatocellular carcinoma cells (HepG2) and leukemia cells (SupT1).

Cell growth and proliferation of mammalian cells occurs through cell cycle; thus, the inhibition of the cell cycle progression is the ideal target for anticancer agents[20,21]. APME exhibited growth inhibitory effects on the MDA-MB-231 cells, inducing cell cycle




**Figure 3.** Effects of methanolic extract of *A. precatarius* leaves on apoptosis in MDA-MB-231 cells. (a) Apoptosis assay in a time-dependent manner. Quadrants represent the percentage of cell populations; Q1- Necrosis, Q2- Late apoptosis, Q3-Early apoptosis, Q4-Live cells; (b) Percentage of each phase of cell death. Data are expressed as mean $\pm$ SD of three repeated experiments. \* $P < 0.05$  is considered significant when comparing treated cells *vs* untreated cells.

arrest at  $G_0/G_1$  phase. Increased percentage of cell population in  $G_0/G_1$  phase and reduction of the population in S-phase proved this claim. In S-phase, genetic information is transferred from one cell generation to another. Genome replication in S-phase is important to segregate two daughter cells during mitosis or the M-phase. Mitosis only occurs when S-phase is completed. Two gaps separate between M- and S-phase. Between M- and S-phase, there is the  $G_1$ , and between S- and M-phase, there is  $G_2$ . DNA damage activates these checkpoints. When growth arrest occurs at any checkpoints, cells will repair the damage. If the damage is repaired, cell progression will successfully resume, otherwise the cell will be eliminated

through apoptosis[21]. DNA arrest occurred during  $G_0/G_1$  phase in this current study, which indicated that the cell proliferation was inhibited, thus showing reduction of the cell percentage in S-phase and  $G_2/M$  phase. At this point, it is clear that cell proliferation was halted by DNA arrest at  $G_0/G_1$  phase. Furthermore, it is important to find out if the cell inhibition was caused by apoptosis.

Apoptosis induction is regarded as the best strategy in cancer treatment. It is an important programmed cell death to eliminate unnecessary cells and thus became the common mode of action for most chemotherapeutic agents[2]. Induction of apoptosis signifies the success of plant products as anticancer agents and it is the





**Figure 4.** Bax, Bcl-2, caspase-3 and p53 protein expression in MDA-MB-231 cells treated with methanolic extract of *A. precatarius* leaves. (a) Apoptosis protein expression by flow cytometry (Bax, Bcl-2, p53, caspase-3). (b) Proteins expressions of Bax, Bcl-2, p53, and caspase-3. Data are expressed as mean±SD of three repeated experiments. \* $P < 0.05$  is considered significant when comparing treated cells vs untreated cells.

optimal way in cancer treatment. In order to confirm whether the inhibition of cell proliferation induced by APME is due to apoptosis, rather than necrosis, apoptosis assay using Annexin V-FITC and PI staining was performed following the treatment with the extract. Annexin V stains the phosphatidylserine of the inner cell membrane which is exposed during the early stage of apoptosis. Our results demonstrated that the APME promotes cell death *via* apoptosis. Early apoptosis occurred after 48 h and eventually led to late apoptosis following 72 h of treatment.

Decrease of mitochondrial outer membrane permeability (MOMP) indicated the irreversible events of early apoptosis. MOMP is highly regulated by anti-apoptotic and pro-apoptotic proteins[22]. Bax is a pro-apoptotic protein and Bcl-2 is an anti-apoptotic protein[23]. Upon stimuli in the intrinsic pathway, Bax is activated which induces MOMP. In this pathway, Bcl-2 prevented MOMP by inhibiting the activity of the BH3-only proteins that are responsible to activate Bax. MOMP allows the release of intermembrane space proteins, such as cytochrome c and SMAC. Cytochrome c forms an apoptosome complex after binding to the APAF-1 protein. This complex is responsible to activate caspase-3 and caspase-7 which eventually lead to apoptosis. Our results indicated that the expression of Bax proteins increased while Bcl-2 proteins decreased. Chien *et al*[24] demonstrated that quercetin induced apoptosis in MDA-MB-231 cells also by reducing Bcl-2 and increasing Bax protein levels. Overexpression of Bcl-2 like proteins such as Bcl-2, Bcl-xL and MCL-1 was detected in various cancers including lymphoma[25], lung cancer[26], neuroblastoma[27] and breast cancer[28]. MOMP is an important event of apoptosis because once it is activated, cell will face death sentence regardless of caspase activation[29].

Increased expression of caspase-3 protein signified the apoptosis events in the MDA-MB-231 cells treated with the APME. Caspase-3 activation can occur both in the extrinsic or intrinsic pathway. Upon activated, caspase-3 will cleave hundreds of other proteins, which subsequently leads to the biochemical and morphological signals of apoptosis. These hallmark events include DNA fragmentation, plasma membrane blebbing and phosphatidylserine exposure[30]. As shown in the apoptosis assay, exposure of phosphatidylserine was detected by Annexin V-FITC and this indicates the early apoptosis event.

The results of this current study exhibited that APME inhibited MDA-MB-231 cell proliferation by DNA arrest at G<sub>0</sub>/G<sub>1</sub> and induced apoptosis through Bax/Bcl-2 and caspase-3 proteins regulation. These findings suggest that APME successfully promotes cell death in breast cancer cell, MDA-MB-231 *via* apoptosis.

### Conflict of interest statement

The authors declare that there is no conflict of interest.

### Funding

This study was funded by the Universiti Sains Malaysia Short Term Grant (304/ PPSP/61313046).

### References

- [1] Green DR. *Means to an end: Apoptosis and other cell death mechanisms*. America: Cold Spring Harbor laboratory press; 2011, p. 5-14.
- [2] Cao K, Tait SW. Apoptosis and cancer: Force awakens, phantom menace, or both?. In: Galluzzi L (ed.) *International review of cell and molecular biology*. Elsevier; 2018, p. 135-152.
- [3] Sreelatha SA, Jeyachitra, Padma PR. Antiproliferation and induction of apoptosis by *Moringa oleifera* leaf extract on human cancer cells. *Food Chem Toxicol* 2011; **49**(6): 1270-1275.
- [4] World Health Organization. *Cancer*. [Online] Available from: <https://www.who.int/news-room/fact-sheets/detail/cancer>. [Accessed on 5th February 2019].
- [5] World Health Organization. *Cancer today: Data visualization tools for exploring the global cancer burden in 2018*. [Online] Available from: <https://gco.iarc.fr/today/home>. [Accessed on 5th February 2019].
- [6] Wan-Ibrahim WS, Ismail TNT, Mohd-Salleh SF, Ismail N. GC-MS analysis of phytochemical compounds in aqueous leaf extract of *Abrus precatorius*. *Pertanika J Trop Agric Sci* 2018; **41**(1): 241-250.
- [7] Gul MZ, Manjulatha K, Bhat MY, Maurya R, Qureshi IA, Ghazi IA. Antiproliferative and apoptosis-inducing effects of *Abrus precatorius* against human monocytic leukaemia (THP-1) cell line. *Indian J Pharm Sci* 2018; **80**(2): 307-317.
- [8] Sofi MS, Sateesh MK, Bashir M, Ganie MA, Nabi S. Chemopreventive and anti-breast cancer activity of compounds isolated from leaves of *Abrus precatorius* L. *3 Biotech* 2018; **8**(8): 371.
- [9] Leбри M, Tilaoui M, Bahi C, Achibat H, Akhramez S, Fofie YBN, et al. Phytochemical analysis and *in vitro* anticancer effect of aqueous extract of *Abrus precatorius* Linn. *Der Pharma Chemica* 2015; **7**(8): 112-117.
- [10] Ghosh T, Mitra P, Jha DK, Mitra PK. *Abrus precatorius* Linnaeus and its biological activities-a review. *World J Pharm Pharm Sci* 2017; **6**(8): 352-368.
- [11] Oladimeji AO, Babatunde O, Musa RT, M'civer FA, Lawal AT, Oguwande IA. GC-MS analysis and cytotoxic activity of essential oils from the leaves of *Abrus precatorius* L. Gaertn. *Asian Pac J Trop Dis* 2016; **6**(5): 372-375.
- [12] Gul MZ, Ahmad F, Kondapi AK, Qureshi IA, Ghazi IA. Antioxidant and antiproliferative activities of *Abrus precatorius* leaf extracts-An *in vitro* study. *BMC Complement Altern Med* 2013; **13**(53): 1-12.
- [13] Sofi MS, Sateesh MK, Bashir M, Harish G, Lakshmeesha TR, Vedashree S, et al. Cytotoxic and pro-apoptotic effects of *Abrus precatorius* L. on human metastatic breast cancer cell line, MDA-MB-231. *Cytotechnology* 2013; **65**(3): 407-417.
- [14] Geran R, Greenberg N, MacDonald M, Schumacher A, Abbott

- B. National Cancer Institute protocols for screening of anticancer compounds. *Cancer Chemother Rep* 1972; **3**: 1-103.
- [15]Igarashi M, Miyazawa T. The growth inhibitory effect of conjugated linoleic acid on a human hepatoma cell line, HepG2, is induced by a change in fatty acid metabolism, but not the facilitation of lipid peroxidation in the cells. *Biochim Biophys Acta* 2001; **1530**(2-3): 162-171.
- [16]Sharma A, Kaur R, Katnoria JK, Kaur R, Nagpal AK. Family Fabaceae: A boon for cancer therapy. In: Malik S (ed.) *Biotechnology and production of anti-cancer compounds*. Springer; 2017, p. 157-175.
- [17]Rungruangmaitree R, Jiraungkoorskul W. Pea, *Pisum sativum*, and its anticancer activity. *Phcog Rev* 2017; **11**(21): 39-42.
- [18]Lee JH, Kim KS, Lee YC, Cho CW, Kim KM, Chung YS. *Anticancer composition containing herbal extract*. US 2011/0293754 A1 (Patent) 2017.
- [19]Ahmad F, Anwar F, Hira S. Review on medicinal importance of fabaceae family. *PhOL* 2016; **3**: 151-156.
- [20]Kim SJ, Min HY, Chung HJ, Park EJ, Hong JY, Kang YJ, et al. Inhibition of cell proliferation through cell cycle arrest and apoptosis by thio-Cl-IB-MECA, a novel A<sub>3</sub> adenosine receptor agonist, in human lung cancer cells. *Cancer Lett* 2008; **264**(2): 309-315.
- [21]Li A, Blow JJ. The origin of CDK regulation. *Nat Cell Biol* 2001; **3**(8): E182-184.
- [22]Czabotar PE, Lessene G, Strasser A, Adams JM. Control of apoptosis by the BCL-2 protein family: Implications for physiology and therapy. *Nat Rev Mol Cell Biol* 2014; **15**(1): 49-63.
- [23]Bai L, Wang S. Targeting apoptosis pathways for new cancer therapeutics. *Annu Rev Med* 2014; **65**: 139-155.
- [24]Chien SY, Wu YC, Chung JG, Yang JS, Lu HF, Tsao MF, et al. Quercetin-induced apoptosis acts through mitochondrial-and caspase-3-dependent pathways in human breast cancer MDA-MB-231 cells. *Hum Exp Toxicol* 2009; **28**(8): 493-503.
- [25]Pepper C, Hoy T, Bentley P. Elevated Bcl-2/Bax are a consistent feature of apoptosis resistance in B-cell chronic lymphocytic leukaemia and are correlated with *in vivo* chemoresistance. *Leuk Lymphoma* 1998; **28**(3-4): 355-361.
- [26]Beroukhim R, Mermei CH, Porter D, Wei G, Raychaudhuri S, Donovan J, et al. The landscape of somatic copy-number alteration across human cancers. *Nature* 2010; **463**(7283): 899-905.
- [27]Castle VP, Heidelberger KP, Bromberg J, Ou X, Dole M, Nuñez G. Expression of the apoptosis-suppressing protein bcl-2, in neuroblastoma is associated with unfavorable histology and N-myc amplification. *Am J Pathol* 1993; **143**(6): 1543-1550.
- [28]Olopade OI, Adeyanju MO, Safa AR, Hagos F, Mick R, Thompson CB, et al. Overexpression of BCL-x protein in primary breast cancer is associated with high tumor grade and nodal metastases. *Cancer J Sci Am* 1997; **3**(4): 230-237.
- [29]Tait SW, Green DR. Mitochondria and cell death: Outer membrane permeabilization and beyond. *Nature Rev Mol Cell Biol* 2010; **11**(9): 621-632.
- [30]McIlwain DR, Berger T, Mak TW. Caspase functions in cell death and disease. *Cold Spring Harb Perspect Biol* 2013; **5**(4): a008656. Doi: 10.1101/cshperspect.a008656.



ISSN 2221-1691(Print)  
ISSN 2588-9222(Online)  
CODEN APJTC7

# Asian Pacific Journal of Tropical Biomedicine



**6**  
**2019**



Vol 9 No 6 Jun