

PHYTOCHEMICAL AND BIOLOGICAL ACTIVITY STUDIES OF
GARCINIA HOMBRONIANA

by

NARGIS JAMILA

Thesis submitted in fulfillment of the requirements

for the degree of

Doctor of Philosophy

JANUARY 2014

ACKNOWLEDGMENT

It would not have been possible to finish my doctoral research without the help and immense support of some obliging and doting people around me. First of all, I would like to deeply acknowledge my supervisor Dr. Melati Khairuddean for her good advice, support and her brooking attitude towards me. Her friendly nature helped me in compiling this thesis very early. I am also thankful to my ex-supervisor, Dr. Wong Keng Chong for providing me a refreshing insight and vitalizing suggestions if I ever needed during the entire research period. A thank is also addressed to my Co-supervisor, Dr. Hasnah Osman for her helpful suggestions during my research phase and thesis compilation.

I would like to thank TWAS (Third World Academy of Sciences) and Institute of Postgraduate Studies, Universiti Sains Malaysia for providing me with a TWAS-USM fellowship. Special thanks to the Dean of the School of Chemical Sciences, Prof. Dr. Wan Ahmad Kamil Mahmood for providing the research facilities.

I thank the technical and laboratory staff, Mr. Chow Cheng Por, Mr. Megat Hasnul Zamani and Mr. Clement D'Silva who have helped me in understanding the operation of equipment. I am also very grateful to Mr. Zahari Othman who has been very supportive and helpful in running the NMR samples for me.

I would also like to thank Dr. Lai Choon Sheen from Centre for Drug Research, Dr. Vikneswaran a/l Murugaiyah from School of Pharmaceutical Sciences and Prof. Nik

Soriani Yaacob from Department of Chemical Pathology, USM, Kebung Kerian for providing me continuous help in biological activity studies.

I value my friends and lab fellows: Naeem Khan (Chosun University, South Korea), Mohamed Ali Abdunnabi Alwahsh, Tan Wen Nee, Foong Shin Yee, Irfan Shah, Alice Wong, Chan Yee Chen, Aboubakar Abdunabi, Samina Khan Yusafzai and Francis who always found time for me on their calendar and have assisted me readily whenever I was stringed to any problem or when I was shattered of being home sick.

Last but not least, I would like to acknowledge my loved ones, my lovable family members for their prayers and devotion. It is an honour to acknowledge that my mother has been an ideal for me. Her encouraging words are always indispensable and superfluous. The overflowing and warmest wishes of my brothers and sisters to see me as a doctoral degree holder were priceless and exemplary.

TABLE OF CONTENTS

	Page
Acknowledgement	ii
Table of Contents	iv
List of Tables	ix
List of Figures	xii
List of Abbreviations/Symbols	xviii
List of Appendices	xxi
Abstrak	xxv
Abstract	xxvii
CHAPTER 1 INTRODUCTION	1
1.1 Natural Products	1
1.2 The Clusiaceae (Guttiferae) Family	2
1.3 The Genus <i>Garcinia</i>	2
1.4 Botanical Characterization of <i>Garcinia</i>	2
1.5 <i>Garcinia hombroniana</i> Pierre	3
1.6 Botanical Characterization of <i>G. hombroniana</i>	3
1.7 Uses of <i>G. hombroniana</i>	4
CHAPTER 2 LITERATURE REVIEW	6
2.1 Phytochemical Studies of <i>Garcinia</i>	6
2.2 Phytochemical Investigation of <i>G. hombroniana</i>	6
2.2.1 Chemical Constituents Isolated from the Stem Wood of <i>G. hombroniana</i>	6
2.2.2 Chemical Constituents Isolated from the Pericarp of <i>G. hombroniana</i>	7
2.2.3 Chemical Constituents Isolated from the Leaves of <i>G. hombroniana</i>	8

2.2.4	Chemical Constituents Isolated from the Twigs of <i>G. hombroniana</i>	9
2.3	Pharmacological Investigation of <i>G. hombroniana</i>	12
2.4	Problem Statement	13
2.5	Objectives of the Research	14
CHAPTER 3 MATERIALS AND METHODS		15
3.1	Plant Material	15
3.2	Chemicals	15
3.2.1	Chemicals for Extraction and Isolation	15
3.2.2	Chemicals for Biological Activities Evaluation	16
3.3	Instruments Used for Structures Elucidation and Biological Activities	17
3.3.1	Melting Point	17
3.3.2	Infrared (IR) Spectroscopy	17
3.3.3	Nuclear Magnetic Resonance (NMR) Spectroscopy	17
3.3.4	Direct Probe Mass Spectrometry	17
3.3.5	Ultraviolet (UV) Spectroscopy	18
3.3.6	Circular Dichroism (CD) Spectroscopy	19
3.3.7	X-Ray Crystallography Analysis	19
3.3.8	Microplate Reader	19
3.4	Extraction Procedure	19
3.5	Phytochemical Screening	21
3.5.1	Detection of Carbohydrates and Glycosides	21
3.5.2	Detection of Proteins and Amino Acids	21
3.5.3	Detection of Terpenoids (Salkowski's Test)	22
3.5.4	Detection of Phytosterols (Liebermann-Burchard's Test)	22
3.5.5	Detection of Phenolic Compounds	22

3.5.6	Detection of Flavonoids	22
3.5.7	Detection of Tannins	23
3.5.8	Detection of Phlobatannins	23
3.5.9	Detection of Alkaloids	23
3.6	Separation Techniques	23
3.6.1	Thin Layer Chromatography (TLC)	23
3.6.2	Column Chromatography (CC)	24
3.7	Isolation and Purification	24
3.7.1	Dichloromethane (DCM) Extract	24
3.7.2	Ethyl Acetate (EtOAc) Extract	28
3.8	Biological Activity Studies of the Bark of <i>G. hombroniana</i>	31
3.8.1	Total Phenolic Contents (Folin-Ciocalteu Assay)	31
3.8.2	Total Flavonoid Contents (Colorimetric Method)	32
3.8.3	<i>In vitro</i> Antioxidant Activities	33
3.8.3.1	DPPH Radical Scavenging Assay	33
3.8.3.2	ABTS Radical Scavenging Assay	34
3.8.3.3	FRAP Assay	34
3.8.4	Antibacterial Activities	35
3.8.4.1	Preparation of Bacteria Culture	35
3.8.4.2	Disc Diffusion Method	35
3.8.4.3	Dilution Method	36
3.8.5	Cholinesterase Enzymes Inhibitory Activities	36
3.8.5.1	AChE Inhibitory Assay	37
3.8.5.2	BChE Inhibitory Assay	37
3.8.6	Cytotoxic Activities	37
3.9	Statistical Analyses	38

CHAPTER 4	RESULTS AND DISCUSSION	39
4.1	Phytochemical Screening of Extracts	39
4.2	Phytochemical Study	39
4.2.1	Constituents of the Dichloromethane Extract	41
4.2.1.1	Compound DN1	41
4.2.1.2	Compound DN2	49
4.2.1.3	Compound DN3	56
4.2.1.4	Compound DN4	63
4.2.1.5	Compound DN5	70
4.2.1.6	Compound DN6	78
4.2.1.7	Compound DN7	86
4.2.1.8	Compound DN8	93
4.2.1.9	Compound DN9	108
4.2.1.10	Compound DN10	124
4.2.2	Constituents of the Ethyl Acetate Extract	132
4.2.2.1	Compound EN1	132
4.2.2.2	Compound EN2	139
4.2.2.3	Compound EN3	152
4.2.2.4	Compound EN4	163
4.2.2.5	Compound EN5	173
4.2.2.6	Compound EN6	194
4.2.2.7	Compound EN7	200
4.2.2.8	Compound EN8	207
CHAPTER 5	BIOLOGICAL ACTIVITY STUDIES	212
5.1	Total Phenolic Contents of Extracts	212
5.2	Total Flavonoid Contents of Extracts	213
5.3	<i>In vitro</i> Antioxidant Activities of Extracts and Pure Compounds	215
5.4	Antibacterial Activities of Extracts and Pure Compounds	218

5.5	Cholinesterase Inhibitory Activities of Extracts and Pure Compounds	220
5.6	Cytotoxic Activities of Some Selected Pure Compounds	222
	CHAPTER 6 CONCLUSION	224
	REFERENCES	226
	APPENDICES	234
	PUBLICATIONS	310

LIST OF TABLES

		Page
Table 4.1	Phytochemical Screening of the Bark Extracts of <i>G. hombroniana</i>	40
Table 4.2	Comparison of $^1\text{H-NMR}$ Data of DN1 and those of Friedelin (Oliviera et al., 2012)*	42
Table 4.3	Comparison of $^{13}\text{C-NMR}$ Data of DN1 and those of Friedelin (Oliveira et al., 2012 ^a ; Klass et al., 1992 ^b)	46
Table 4.4	Comparison of $^1\text{H-NMR}$ Data of DN2 and those of Stigmasterol (Koay et al., 2013)*	50
Table 4.5	Comparison of $^{13}\text{C-NMR}$ Data of DN2 and those of Stigmasterol (Koay et al., 2013 ^a ; Xie et al., 2000 ^b)	52
Table 4.6	Comparison of $^1\text{H-NMR}$ Data of DN3 and those of Lupeol (Burns et al., 2000)*	57
Table 4.7	Comparison of $^{13}\text{C-NMR}$ Data of DN3 and those of Lupeol (Burns et al., 2000 ^a ; Mouffok et al., 2012 ^b)	59
Table 4.8	Comparison of $^1\text{H-NMR}$ Data of DN4 and those of Taraxerol (Mejin, 2009)*	64
Table 4.9	Comparison of $^{13}\text{C-NMR}$ Data of DN4 and those of Taraxerol (Mejin, 2009 ^a ; Koay et al., 2013 ^b)	66
Table 4.10	Comparison of $^1\text{H-NMR}$ Data of DN5 and those of Betulin (Gherraf et al., 2010)*	71
Table 4.11	Comparison of $^{13}\text{C-NMR}$ Data of DN5 and those of Betulin (Gherraf et al., 2010 ^a ; Tijjani et al., 2012 ^b)	73
Table 4.12	Comparison of $^1\text{H-NMR}$ Data of DN6 and those of Garcihombronane D (Rukachaisirikul et al., 2000)*	79
Table 4.13	Comparison of $^{13}\text{C-NMR}$ Data of DN6 and those of Garcihombronane D (Rukachaisirikul et al., 2000) ^a	81

Table 4.14	Comparison of ^1H -NMR Data of DN7 and those of Garcihombronane B (Rukachaisirikul et al., 2000)*	87
Table 4.15	Comparison of ^{13}C -NMR Data of DN7 and those of Garcihombronane B (Rukachaisirikul et al., 2000) ^a	89
Table 4.16	Comparison of ^{13}C -NMR Data of DN8 and those of Garcihombronane I Monoacetate (Rukachaisirikul et al., 2005a) ^a	97
Table 4.17	^1H , ^1H - ^1H COSY, ^1H - ^1H NOESY and ^1H - ^{13}C HMBC NMR Data of DN8 (MeOD- d_4)	99
Table 4.18	^1H (500 MHz) and ^{13}C -NMR (125 MHz) Data of DN9 (MeOD- d_4)	114
Table 4.19	^1H - ^1H COSY, ^1H - ^1H ROESY and ^1H - ^{13}C HMBC NMR Data of DN9 (MeOD- d_4)	120
Table 4.20	Comparison of ^1H -NMR Data of DN10 (mixture), Betulinic acid (Peng et al., 1998) ^a and Betulin (Gherraf et al., 2010) ^b	126
Table 4.21	Comparison of ^{13}C -NMR Data of DN10 (mixture), Betulinic acid (Peng et al., 1998) ^a and Betulin (Gherraf et al., 2010) ^b	128
Table 4.22	^1H and ^{13}C -NMR Data of DN10 (Pure)	129
Table 4.23	Comparison of ^1H -NMR Data of EN1 and those of Stigmasterol Glucoside (Alam et al., 1996)*	133
Table 4.24	Comparison of ^{13}C -NMR Data of EN1 and those of Stigmasterol Glucoside (Alam et al., 1996) ^a	138
Table 4.25	^1H , ^1H - ^1H ROESY, ^{13}C and ^1H - ^{13}C HMBC NMR Data of EN2 (MeOD- d_4 , 500/125 MHz)	147
Table 4.26	Comparison of ^1H -NMR Data of EN3 and those of Morelloflavone (Li et al., 2002)* at 25 °C	154
Table 4.27	Comparison of ^{13}C -NMR Data of EN3 and those of Morelloflavone (Li et al., 2002) ^a at 25 °C	157
Table 4.28	Comparison of ^1H -NMR Data of EN4 and those of Volkensiflavone (Li et al., 2002)* at 25 °C	165

Table 4.29	Comparison of ^{13}C -NMR Data of EN4 and those of Volkensiflavone (Compagnone et al., 2008) ^a at 25 °C	167
Table 4.30	^1H -NMR Data of EN5 (DMSO- <i>d</i> ₆ and MeOD- <i>d</i> ₄ , 500 MHz) at 25 °C	178
Table 4.31	^{13}C -NMR Data of EN5 at 25 °C	181
Table 4.32	^1H - ^1H COSY, ^1H - ^1H NOESY and ^1H - ^{13}C HMBC NMR Data of the Major Conformer of EN5 at 25 °C	186
Table 4.33	Comparison of ^1H -NMR Data of EN6 and those of Volkensiflavone (Compagnone et al., 2008)* at 25 °C	195
Table 4.34	^{13}C -NMR Data of EN6 at 25 °C	197
Table 4.35	Comparison of ^1H -NMR Data of EN7 and those of 4',5,7-Trihydroxyflavanone-7-rutinoside (Kim et al., 2007)*	201
Table 4.36	Comparison of ^{13}C -NMR Data of EN7 and those of 4',5,7-Trihydroxyflavanone-7-rutinoside (Kim et al., 2007) ^a	203
Table 4.37	Comparison of ^1H and ^{13}C -NMR Data of EN8 and those of Norathyriol (Noro et al., 1984)*	208
Table 5.1	Antioxidant Activities, Total Phenolic and Flavonoid Contents of the Extracts and Some Isolated Pure Compounds of <i>G. hombroniana</i>	216
Table 5.2	Antibacterial Activities of the Extracts and Some Isolated Pure Compounds of <i>G. hombroniana</i>	219
Table 5.3	Cholinesterase Inhibitory Activities of the Extracts and Some Isolated Pure Compounds of <i>G. hombroniana</i>	221

LIST OF FIGURES

		Page
Figure 1.1	<i>Garcinia hombroniana</i> Pierre	4
Figure 1.2	Parts of <i>G. hombroniana</i>	5
Figure 3.1	Extraction procedure of <i>G. hombroniana</i> bark	20
Figure 3.2	Isolation and purification procedure of the dichloromethane bark extract of <i>G. hombroniana</i>	27
Figure 3.3	Isolation and purification procedure of the ethyl acetate bark extract of <i>G. hombroniana</i>	30
Figure 4.1	¹ H-NMR (a) full and (b) expanded spectra of DN1 (CDCl ₃ , 500 MHz)	43
Figure 4.2	¹³ C-NMR (a) full and (b) expanded (δ_C 5-45 ppm) spectra of DN1 (CDCl ₃ , 125 MHz)	47
Figure 4.3	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN1 (CDCl ₃ , 125 MHz)	48
Figure 4.4	¹ H-NMR (a) full, and (b and c) expanded spectra of DN2 (CDCl ₃ , 500 MHz)	51
Figure 4.5	¹³ C-NMR (a) full and (b) expanded (δ_C 10-60 ppm) spectra of DN2 (CDCl ₃ , 125 MHz)	53
Figure 4.6	DEPT (a) 135 and (b) 90 NMR spectra of DN2 (CDCl ₃ , 125 MHz)	54
Figure 4.7	¹ H-NMR (a) full, and (b and c) expanded spectra of DN3 (CDCl ₃ , 500 MHz)	58
Figure 4.8	¹³ C-NMR (a) full and (b) expanded (δ_C 14-56 ppm) spectra of DN3 (CDCl ₃ , 125 MHz)	60
Figure 4.9	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN3 (CDCl ₃ , 125 MHz)	61

Figure 4.10	¹ H-NMR (a) full, and (b and c) expanded spectra of DN4 (CDCl ₃ , 500 MHz)	65
Figure 4.11	¹³ C-NMR (a) full and (b) expanded (δ_C 15-50 ppm) spectra of DN4 (CDCl ₃ , 125 MHz)	67
Figure 4.12	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN4 (CDCl ₃ , 125 MHz)	68
Figure 4.13	¹ H-NMR (a) full, and (b and c) expanded spectra of DN5 (CDCl ₃ , 500 MHz)	72
Figure 4.14	¹³ C-NMR (a) full and (b) expanded (δ_C 14-62 ppm) spectra of DN5 (CDCl ₃ , 125 MHz)	74
Figure 4.15	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN5 (CDCl ₃ , 125 MHz)	75
Figure 4.16	¹ H-NMR (a) full, and (b and c) expanded spectra of DN6 (DMSO- <i>d</i> ₆ , 500 MHz)	80
Figure 4.17	¹³ C-NMR (a) full and (b) expanded (δ_C 15-53 ppm) spectra of DN6 (DMSO- <i>d</i> ₆ , 125 MHz)	82
Figure 4.18	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN6 (DMSO- <i>d</i> ₆ , 125 MHz)	83
Figure 4.19	¹ H-NMR (a) full, and (b-d) expanded spectra of DN7 (MeOD- <i>d</i> ₄ , 500 MHz)	88
Figure 4.20	¹³ C-NMR (a) full and (b) expanded (δ_C 12-56 ppm) spectra of DN7 (MeOD- <i>d</i> ₄ , 125 MHz)	90
Figure 4.21	DEPT (a) 135 and (b) 90 NMR spectra of DN7 (MeOD- <i>d</i> ₄ , 125 MHz)	91
Figure 4.22	EI and +ESI-MS of DN8	94
Figure 4.23	IR spectrum of DN8	95
Figure 4.24	¹ H-NMR (a) full, and (b and c) expanded spectra of DN8 (MeOD- <i>d</i> ₄ , 500 MHz)	96

Figure 4.25	^{13}C -NMR (a) full and (b) expanded (δ_{C} 12-54 ppm) spectra of DN8 (MeOD- d_4 , 125 MHz)	100
Figure 4.26	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN8 (MeOD- d_4 , DMSO- d_6 , 125 MHz)	101
Figure 4.27	^1H - ^1H COSY NMR spectrum of DN8 (MeOD- d_4)	102
Figure 4.28	^1H - ^{13}C HSQC NMR spectrum of DN8 (MeOD- d_4)	103
Figure 4.29	^1H - ^{13}C HMBC NMR spectrum of DN8 (MeOD- d_4)	104
Figure 4.30	^1H - ^1H NOESY NMR spectrum of DN8 (MeOD- d_4)	106
Figure 4.31	(a) + and (b) –ESI-MS of DN9	109
Figure 4.32	UV spectrum of DN9	110
Figure 4.33	IR spectrum of DN9	111
Figure 4.34	^1H -NMR (a) full and (b-e) expanded spectra of DN9 (MeOD- d_4 , 500 MHz)	112
Figure 4.35	^{13}C -NMR (a) full and (b) expanded (δ_{C} 16-53) spectra of DN9 (MeOD- d_4 , 125 MHz)	115
Figure 4.36	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN9 (MeOD- d_4 , 125 MHz)	116
Figure 4.37	^1H - ^1H COSY NMR spectrum of DN9 (MeOD- d_4)	117
Figure 4.38	^1H - ^{13}C HSQC NMR spectrum of DN9 (MeOD- d_4)	118
Figure 4.39	^1H - ^{13}C HMBC NMR spectrum of DN9 (MeOD- d_4)	122
Figure 4.40	^1H - ^1H ROESY NMR spectrum of DN9 (MeOD- d_4)	123
Figure 4.41	^1H -NMR (a) full and (b) expanded spectra of DN10 (mixture) ($\text{C}_5\text{D}_5\text{N}$, 500 MHz)	127
Figure 4.42	^{13}C -NMR (a) full and (b) expanded (δ_{C} 15-58 ppm) spectra of DN10 (mixture) ($\text{C}_5\text{D}_5\text{N}$, 125 MHz)	130

Figure 4.43	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of DN10 (mixture) (C_5D_5N , 125 MHz)	131
Figure 4.44	1H -NMR (a) full and (b-d) expanded spectra of EN1 ($DMSO-d_6$, 500 MHz)	134
Figure 4.45	^{13}C -NMR (a) full and (b) expanded (δ_C 10-40 ppm) spectra of EN1 ($DMSO-d_6$, 125 MHz)	135
Figure 4.46	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN1 ($DMSO-d_6$, 125 MHz)	136
Figure 4.47	EI-MS of EN2	140
Figure 4.48	UV spectrum of EN2	141
Figure 4.49	IR spectrum of EN2	142
Figure 4.50	1H -NMR (a) full and (b) expanded spectra of EN2 ($MeOD-d_4$, 500 MHz)	143
Figure 4.51	^{13}C -NMR spectrum of EN2 ($MeOD-d_4$, 125 MHz)	144
Figure 4.52	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN2 ($MeOD-d_4$, 125 MHz)	145
Figure 4.53	1H - ^{13}C HMQC NMR spectrum of EN2 ($MeOD-d_4$)	148
Figure 4.54	1H - ^{13}C HMBC NMR spectrum of EN2 ($MeOD-d_4$)	149
Figure 4.55	1H - 1H ROESY NMR spectrum of EN2	150
Figure 4.56	Single crystal X-Ray structure of EN2	151
Figure 4.57	1H -NMR (a) full and (b) expanded spectra of EN3 at 25 °C ($DMSO-d_6$, 500 MHz)	155
Figure 4.58	^{13}C -NMR (a) full and (b) expanded (δ_C 95-168 ppm) spectra of EN3 at 25 °C ($DMSO-d_6$, 125 MHz)	158
Figure 4.59	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN3 at 25 °C ($DMSO-d_6$, 125 MHz)	159

Figure 4.60	^1H -NMR spectrum of EN3 at 90 °C (DMSO- d_6 , 500 MHz)	160
Figure 4.61	^{13}C -NMR spectrum of EN3 at 90 °C (DMSO- d_6 , 125 MHz)	161
Figure 4.62	^1H -NMR (a) full and (b) expanded spectra of EN4 at 25 °C (DMSO- d_6 , 500 MHz)	166
Figure 4.63	^{13}C -NMR (a) full and (b) expanded (δ_{C} 68-168 ppm) spectra of EN4 at 25 °C (DMSO- d_6 , 125 MHz)	168
Figure 4.64	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN4 at 25 °C (DMSO- d_6 , 125 MHz)	169
Figure 4.65	^1H -NMR spectrum of EN4 at 90 °C (DMSO- d_6 , 500 MHz)	170
Figure 4.66	^{13}C -NMR spectrum of EN4 at 90 °C (DMSO- d_6 , 125 MHz)	171
Figure 4.67	LR (+) and HR (-) ESI-MS of EN5	175
Figure 4.68	UV spectra of EN5 with MeOH and shift reagents (NaOMe, NaOAc, H_3BO_3 , AlCl_3 and HCl)	176
Figure 4.69	IR spectrum of EN5	177
Figure 4.70	^1H -NMR (a) full and (b) expanded spectra of EN5 at 25 °C (DMSO- d_6 , 500 MHz)	179
Figure 4.71	^{13}C -NMR (a) full and (b) expanded (δ_{C} 68-168 ppm) spectra of EN5 at 25 °C (DMSO- d_6 , 125 MHz)	182
Figure 4.72	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN5 at 25 °C (DMSO- d_6 , 125 MHz)	183
Figure 4.73	^1H -NMR spectrum of EN5 at 90 °C (DMSO- d_6 , 500 MHz)	184
Figure 4.74	^{13}C -NMR spectrum of EN5 at 90 °C (DMSO- d_6 , 125 MHz)	185
Figure 4.75	^1H - ^1H COSY NMR spectrum of EN5 (DMSO- d_6)	187
Figure 4.76	^1H - ^{13}C HSQC NMR spectrum of EN5 (DMSO- d_6)	188
Figure 4.77	^1H - ^{13}C HMBC NMR spectrum of EN5 (DMSO- d_6)	191
Figure 4.78	^1H - ^1H NOESY NMR spectrum of EN5 (DMSO- d_6)	192

Figure 4.79	CD spectrum of EN5	193
Figure 4.80	¹ H-NMR (a) full and (b) expanded spectra of EN6 at 25 °C (DMSO- <i>d</i> ₆ , 500 MHz)	196
Figure 4.81	¹³ C-NMR (a) full and (b) expanded (δ_C 68-168 ppm) spectra of EN6 at 25 °C (DMSO- <i>d</i> ₆ , 125 MHz)	198
Figure 4.82	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN6 at 25 °C (DMSO- <i>d</i> ₆ , 125 MHz)	199
Figure 4.83	¹ H-NMR (a) full, and (b and c) expanded spectra of EN7 (DMSO- <i>d</i> ₆ , 500 MHz)	202
Figure 4.84	¹³ C-NMR (a) full and (b) expanded (δ_C 68-104 ppm) spectra of EN7 (DMSO- <i>d</i> ₆ , 125 MHz)	204
Figure 4.85	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN7 (DMSO- <i>d</i> ₆ , 125 MHz)	205
Figure 4.86	¹ H-NMR (a) full and (b) expanded spectra of EN8 (MeOD- <i>d</i> ₄ , 500 MHz)	209
Figure 4.87	¹³ C-NMR spectrum of EN8 (MeOD- <i>d</i> ₄ , 125 MHz)	210
Figure 4.88	DEPT (a) 135, (b) 90 and (c) Q NMR spectra of EN8 (MeOD- <i>d</i> ₄ , 125 MHz)	211
Figure 5.1	Calibration curve of gallic acid in total phenolic contents determination	213
Figure 5.2	Calibration curves of standards (a) rutin and (b) quercetin in flavonoid contents determination	214
Figure 5.3	Cytotoxic effects of DN6 and EN2 on DBTRG human cancer cell lines	223
Figure 5.4	(A) Dose- and time-dependent cytotoxic effects of DN6 and EN2 on DBTRG cells and (B) EC ₅₀ determination	223

LIST OF ABBREVIATIONS/SYMBOLS

Abbreviations

MeOD- <i>d</i> ₄	Deuterated Methanol
CDCl ₃	Deuterated Chloroform
DMSO- <i>d</i> ₆	Deuterated Dimethyl Sulfoxide
Acetone- <i>d</i> ₆	Deuterated Acetone
C ₅ D ₅ N	Deuterated Pyridine
AR	Analytical Reagent
HPLC	High Performance Liquid Chromatography
EI-MS	Electron Ionization Mass Spectrometry
ESI-MS	Electrospray Ionization Mass Spectrometry
HR-ESI-MS	High Resolution Electrospray Ionization Mass Spectrometry
LR-ESI-MS	Low Resolution Electrospray Ionization Mass Spectrometry
FT-IR	Fourier Transformer Infrared
1D-NMR	One Dimensional Nuclear Magnetic Resonance
2D-NMR	Two Dimensional Nuclear Magnetic Resonance
COSY	Correlation Spectroscopy
DQF COSY	Double Quantum Filtered Correlation Spectroscopy
DEPT	Distortionless Enhancement by Polarization Transfer
HMBC	Heteronuclear Multiple Bond Correlation
HMQC	Heteronuclear Multiple Quantum Correlation
HSQC	Heteronuclear Single Quantum Correlation
NOESY	Nuclear Overhauser Enhancement Spectroscopy

NOE	Nuclear Overhauser Effect
ROESY	Rotating Frame Overhauser Enhancement Spectroscopy
LDL	Low Density Lipoproteins
MRSA	Methicillin Resistant <i>Staphylococcus aureus</i>
MIC	Minimum Inhibitory Concentration
AChE	Acetylcholinesterase Enzyme
BChE	Butyrylcholinesterase Enzyme

Symbols

m/z	mass/charge
w/v	weight/volume
v/v	volume/volume
R_f	retention factor
nm	nano meter
MHz	mega Hertz
s	singlet
br. s	broad singlet
d	doublet
br. d	broad doublet
dd	doublet of doublets
ddd	doublet of doublets of doublets
t	triplet
dt	doublet of triplet
q	quartet
dq	doublet of quartet

<i>m</i>	multiplet
α	alpha
β	beta
μg	microgram
ppm	parts per million
SD	standard deviation
cm^{-1}	per centimeter
$^{\circ}\text{C}$	degree Celsius
δ	chemical shift
g	gram
$\mu\text{g/mL}$	microgram per millilitre
μL	microlitre
μM	micromole
IC_{50}	concentration required inhibiting 50% of effect
EC_{50}	effective concentration required to induce a 50% effect
λ_{max}	Lambda (Maximum Wavelength)

LIST OF APPENDICES

	Page
Appendix A	EI-MS of DN1 234
Appendix B	IR spectrum of DN1 235
Appendix C	^1H - ^1H DQF-COSY NMR spectrum of DN1 (CDCl_3) 236
Appendix D	^1H - ^{13}C HSQC NMR spectrum of DN1 (CDCl_3) 237
Appendix E	^1H - ^{13}C HMBC NMR spectrum of DN1 (CDCl_3) 238
Appendix F	^1H - ^1H NOESY NMR spectrum of DN1 (CDCl_3) 239
Appendix G	EI-MS of DN2 240
Appendix H	IR spectrum of DN2 241
Appendix I	^1H - ^1H COSY NMR spectrum of DN2 (CDCl_3) 242
Appendix J	^1H - ^{13}C HMQC NMR spectrum of DN2 (CDCl_3) 243
Appendix K	^1H - ^{13}C HMBC NMR spectrum of DN2 (CDCl_3) 244
Appendix L	^1H - ^1H ROESY NMR spectrum of DN2 (CDCl_3) 245
Appendix M	EI-MS of DN3 246
Appendix N	IR spectrum of DN3 247
Appendix O	^1H - ^1H COSY NMR spectrum of DN3 (CDCl_3) 248
Appendix P	^1H - ^{13}C HMQC NMR spectrum of DN3 (CDCl_3) 249
Appendix Q	^1H - ^{13}C HMBC NMR spectrum of DN3 (CDCl_3) 250
Appendix R	^1H - ^1H NOESY NMR spectrum of DN3 (CDCl_3) 251
Appendix S	EI-MS of DN4 252

Appendix T	IR spectrum of DN4	253
Appendix U	^1H - ^1H COSY NMR spectrum of DN4 (CDCl_3)	254
Appendix V	^1H - ^{13}C HSQC NMR spectrum of DN4 (CDCl_3)	255
Appendix W	^1H - ^{13}C HMBC NMR spectrum of DN4 (CDCl_3)	256
Appendix X	^1H - ^1H NOESY NMR spectrum of DN4 (CDCl_3)	257
Appendix Y	EI-MS of DN5	258
Appendix Z	IR spectrum of DN5	259
Appendix AA	^1H - ^1H DQF-COSY NMR spectrum of DN5 (CDCl_3)	260
Appendix AB	^1H - ^{13}C HSQC NMR spectrum of DN5 (CDCl_3)	261
Appendix AC	^1H - ^{13}C HMBC NMR spectrum of DN5 (CDCl_3)	262
Appendix AD	^1H - ^1H NOESY NMR spectrum of DN5 (CDCl_3)	263
Appendix AE	EI-MS of DN6	264
Appendix AF	IR spectrum of DN6	265
Appendix AG	^1H - ^1H COSY NMR spectrum of DN6 ($\text{DMSO-}d_6$)	266
Appendix AH	^1H - ^{13}C HSQC NMR spectrum of DN6 ($\text{DMSO-}d_6$)	267
Appendix AI	^1H - ^{13}C HMBC NMR spectrum of DN6 ($\text{DMSO-}d_6$)	268
Appendix AJ	^1H - ^1H ROESY NMR spectrum of DN6 ($\text{DMSO-}d_6$)	269
Appendix AK	EI-MS of DN7	270
Appendix AL	IR spectrum of DN7	271
Appendix AM	^1H - ^1H COSY NMR spectrum of DN7 ($\text{MeOD-}d_4$)	272
Appendix AN	^1H - ^{13}C HMQC NMR spectrum of DN7 ($\text{MeOD-}d_4$)	273
Appendix AO	^1H - ^{13}C HMBC NMR spectrum of DN7 ($\text{MeOD-}d_4$)	274
Appendix AP	EI-MS of DN10 (pure)	275

Appendix AQ	IR spectrum of DN10 (pure)	276
Appendix AR	¹ H-NMR spectrum of DN10 (pure) (CDCl ₃ , 500 MHz)	277
Appendix AS	¹³ C-NMR (a) full (b) expanded (14-68 ppm) spectrum of DN10 (pure) (CDCl ₃ , 125 MHz)	278
Appendix AT	EI-MS of EN1	279
Appendix AU	IR spectrum of EN1	280
Appendix AV	¹ H- ¹ H DQF-COSY NMR spectrum of EN1 (DMSO- <i>d</i> ₆)	281
Appendix AW	¹ H- ¹³ C HSQC NMR spectrum of EN1 (DMSO- <i>d</i> ₆)	282
Appendix AX	¹ H- ¹³ C HMBC NMR spectrum of EN1 (DMSO- <i>d</i> ₆)	283
Appendix AY	EI-MS of EN3	284
Appendix AZ	UV spectra of EN3 with MeOH and shift reagents (NaOMe, NaOAc, H ₃ BO ₃ , AlCl ₃ and HCl)	285
Appendix BA	IR spectrum of EN3	286
Appendix BB	¹ H- ¹ H COSY NMR spectrum of EN3 (DMSO- <i>d</i> ₆)	287
Appendix BC	¹ H- ¹³ C HMQC NMR spectrum of EN3 (DMSO- <i>d</i> ₆)	288
Appendix BD	¹ H- ¹³ C HMBC NMR spectrum of EN3 (DMSO- <i>d</i> ₆)	289
Appendix BE	¹ H- ¹ H NOESY NMR spectrum of EN3 (DMSO- <i>d</i> ₆)	290
Appendix BF	EI-MS of EN4	291
Appendix BG	UV spectra of EN4 with MeOH and shift reagents (NaOMe, NaOAc, H ₃ BO ₃ , AlCl ₃ and HCl)	292
Appendix BH	IR spectrum of EN4	293
Appendix BI	¹ H- ¹ H COSY NMR spectrum of EN4 (DMSO- <i>d</i> ₆)	294
Appendix BJ	¹ H- ¹³ C HSQC NMR spectrum of EN4 (DMSO- <i>d</i> ₆)	295

Appendix BK	^1H - ^{13}C HMBC NMR spectrum of EN4 (DMSO- d_6)	296
Appendix BL	^1H - ^1H NOESY NMR spectrum of EN4 (DMSO- d_6)	297
Appendix BM	EI-MS of EN6	298
Appendix BN	^1H - ^1H COSY NMR spectrum of EN6 (DMSO- d_6)	299
Appendix BO	^1H - ^{13}C HMQC NMR spectrum of EN6 (DMSO- d_6)	300
Appendix BP	^1H - ^{13}C HMBC NMR spectrum of EN6 (DMSO- d_6)	301
Appendix BQ	EI-MS of EN7	302
Appendix BR	IR spectrum of EN7	303
Appendix BS	^1H - ^1H COSY NMR spectrum of EN7 (DMSO- d_6)	304
Appendix BT	^1H - ^{13}C HMQC NMR spectrum of EN7 (DMSO- d_6)	305
Appendix BU	^1H - ^{13}C HMBC NMR spectrum of EN7 (DMSO- d_6)	306
Appendix BV	^1H - ^1H NOESY NMR spectrum of EN7 (DMSO- d_6)	307
Appendix BW	^1H - ^1H COSY NMR spectrum of EN8 (MeOD- d_4)	308
Appendix BX	^1H - ^{13}C HMQC NMR spectrum of EN8 (MeOD- d_4)	309

KAJIAN FITOKIMIA DAN AKTIVITI BIOLOGI BAGI *GARCINIA* *HOMBRONIANA*

ABSTRAK

Kajian ini ialah laporan pertama yang menggambarkan kajian fitokimia dan aktiviti biologi bagi kulit batang *G. hombroniana*. Ekstrak diklorometana dan etil asetat bagi kulit pokok *Garcinia hombroniana* yang dipisahkan secara kromatografi turus telah memberikan 4 sebatian baru (**DN8, DN9, EN2 dan EN5**) dan 14 sebatian yang telah diketahui (**DN1-DN7, DN10, EN1, EN3, EN4, EN6-EN8**). Struktur bagi semua sebatian baru telah dicirikan secara kaedah spektroskopi UV, IR dan 1D dan 2D NMR, EI-MS dan ESI-MS. Kesemua sebatian yang diketahui telah dikenalpasti dan ditentukan secara perbandingan data spektroskopi sebatian ini dengan data yang telah dilaporkan.

Ekstrak mentah telah dikaji bagi menentukan jumlah kandungan fenolik dan jumlah kandungan flavonoid. Antioksidan, antibakteria dan aktiviti perencat kolinesterase bagi ekstrak tersebut dan beberapa sebatian tulen turut dikaji. Sebatian **DN6** dan **EN2** juga telah dikaji bagi aktiviti sitotoksik. Ekstrak etil asetat dan metanol telah menunjukkan kandungan fenolik yang sama manakala kandungan flavonoid didapati lebih tinggi dalam ekstrak etil asetat. Kesemua ekstrak polar telah menunjukkan aktiviti antioksidan yang ketara dalam ujian DPPH, ABTS dan FRAP. Bagi sebatian tulen yang dikaji **EN2, EN3, EN4, EN8** dan **DN9** telah menunjukkan aktiviti memerangkap radikal yang paling tinggi dalam cerakin DPPH. Aktiviti tersebut adalah lebih tinggi berbanding aktiviti memerangkap radical bagi rujukan piawai, Trolox. Sebatian **EN2, EN8** dan **EN3** juga menunjukkan aktiviti

pemerangkapan yang paling tinggi dalam ujian ABTS, yaitu tiga kali lebih tinggi berbanding aktiviti Trolox tetapi hampir sama dengan aktiviti pememerangkapan sebatian rujukan, asid gallik.

Dalam asai aktiviti antibakteria terhadap dua bakteria Gram positif, *Staphylococcus aureus*, *Bacillus subtilis* dan dua bakteria Gram negatif, *Pseudomonas aeruginosa* dan *Escherichia coli*, ekstrak diklorometana telah merencatkan pertumbuhan *B. subtilis* dan *P. aeruginosa*, dengan nilai MIC 62.5 µg/mL. Akan tetapi, ekstrak metanol dan air didapati tidak aktif terhadap kesemua strain bakteria yang dikaji. Aktiviti antibakteria bagi sebatian tulen **DN1-DN10**, **EN1** dan **EN2** juga telah dikaji terhadap strain bakteria yang disebutkan di atas tetapi tiada yang menunjukkan aktiviti antibakteria.

Dalam asai aktiviti perencatan kolinesterase, ekstrak etil asetat telah menunjukkan perencatan yang paling tinggi terhadap AChE dan BChE tetapi ekstrak metanol dan air telah menunjukkan perencatan yang sangat lemah. Antara sebatian tulen yang diuji (**DN1-DN10** dan **EN1-EN7**), tulen **DN4-DN7**, **DN9**, **DN10** dan **EN2** yang menunjukkan aktiviti perencatan terhadap AChE dan BChE.

Sebatian tulen **DN6** dan **EN2** telah menunjukkan aktiviti sitotoksik yang lebih tinggi terhadap sel DBTRG berbanding sel PC-3, U2OS dan MCF-7. Aktiviti sitotoksik bagi sebatian tulen ini didapati bergantung kepada faktor leluhur sel, masa dan kepekatan.

Keputusan kajian ini telah menonjolkan potensi kegunaan *G. hombroniana* dalam bidang perubatan tradisional dan penemuan ubat-ubatan.

**PHYTOCHEMICAL AND BIOLOGICAL ACTIVITY STUDIES OF
*GARCINIA HOMBRONIANA***

ABSTRACT

This study is the first report describing the phytochemical investigation and biological activities of the bark of *G. hombroniana*. The dichloromethane and ethyl acetate extracts of the bark of *Garcinia hombroniana* were separated by column chromatography to yield 4 new (**DN8**, **DN9**, **EN2** and **EN5**) and 14 known (**DN1-DN7**, **DN10**, **EN1**, **EN3**, **EN4**, **EN6-EN8**) compounds. The structures of the new compounds were characterized by spectroscopic techniques such as UV, IR, and 1D and 2D NMR, together with EI and ESI-MS spectrometric methods. The known compounds were identified and confirmed by comparison of their spectroscopic data with the reported literature.

Crude extracts were examined for total phenolic and total flavonoids contents. Antioxidant, antibacterial and cholinesterase inhibitory activities of these extracts together with some selected pure compounds were also examined. Compounds **DN6** and **EN2** were further tested for cytotoxic activities. The ethyl acetate and methanol extracts showed equal phenolic contents, while flavonoids content was found to be higher in the ethyl acetate extract. All the polar extracts showed significant antioxidant activities by DPPH, ABTS and FRAP assays. For pure compounds **EN2**, **EN3**, **EN4**, **EN8** and **DN9**, they showed the strongest scavenging activities in DPPH assay, all of which were higher than that of the referenced standard, Trolox. Compounds **EN2**, **EN3** and **EN8** also showed the

strongest scavenging activities in ABTS assay which is three times higher than that of Trolox and almost equal to that of the referenced compound, gallic acid.

In the antibacterial activity assays which were carried out against two Gram positive bacteria, *Staphylococcus aureus*, *Bacillus subtilis*, and two Gram negative bacteria, *Pseudomonas aeruginosa* and *Escherichia coli*, the dichloromethane extract inhibited the growth of *B. subtilis* and *P. aeruginosa*, with a MIC value of 62.5 µg/mL. However, the methanol and water extracts were found to be inactive towards all the tested bacterial strains. The pure compounds **DN1-DN10**, **EN1** and **EN2** were also tested for antibacterial activities against the aforementioned bacteria strain but none of them showed antibacterial activity.

In cholinesterase inhibitory activity assays, ethyl acetate extract showed the strongest inhibition against AChE and BChE but methanol and water extracts showed very weak inhibition. Among the tested pure compounds (**DN1-DN10** and **EN1-EN7**), compounds **DN4-DN7**, **DN9**, **DN10** and **EN2** showed inhibition against AChE and BChE.

Pure compounds **DN6** and **EN2** possessed significant cytotoxic activities against DBTRG cells as compared with PC-3, U2OS and MCF-7 cells. The cytotoxic activities of these compounds were found to be cell-line, time and concentration dependent.

Results of this study concluded the potential uses of *G. hombroniana* in traditional medicine and drug discovery.

CHAPTER 1

INTRODUCTION

1.1 Natural Products

The name “natural products” is given to compounds derived from natural sources such as plants, animals and microorganisms, which are used for self-defense and to enhance survival period (Williams et al., 1989). More than 80,000 of 250,000 species of plants are used as traditional medicine for healing purposes (Newman, 2007). They can be either in crude forms or as pure compounds, derived from a whole plant or certain parts of a plant, or from exudates of plants (Joy et al., 2001). Today, about three-quarter of the world’s population used herbal medicine in their health care needs (Moerman, 1996). However, only a small percentage of plants have been evaluated for their phytochemical and pharmacological potential (Tringali et al., 2001) while a large portion of which remains unknown. Interests in the use of traditional medicine are increasing rapidly due to their easy accessibility, a relatively low cost and low side effects to be used against various diseases including cancer, HIV/AIDS, and etcetera (Liang & Fang, 2006; WHO, 2002). A rapid development of natural products chemistry has resulted in a remarkable effect on the discoveries of bioactive components in natural sources. The present research work focuses on a plant named *Garcinia hombroniana* (from the Clusiaceae family). The bark of *G. hombroniana* was used in the extraction and isolation of some pure compounds. Characterization of their chemical structures was carried out using various spectroscopic techniques. These compounds were also assayed for their antioxidant, antibacterial, anticholinesterase and cytotoxic activities.

1.2 The Clusiaceae (Guttiferae) Family

The Clusiaceae family in Peninsular Malaysia is composed of approximately 37 genera and 1610 species. The plants are trees, shrubs or herbs which can grow from small to medium sized woody plants producing resinous white to yellow viscous exudates. Several plants of the Clusiaceae family produce important commercial products, edible fruits and more than 15% of oil from the seeds, and the timbers of many of its species are used for building houses and making furniture (Xiwen & Yan-hui, 1990).

1.3 The Genus *Garcinia*

Garcinia is the largest genus of the Clusiaceae family with about 450 species, indigenous to tropical Asia, tropical and southern Africa, Madagascar, North East Australia, West Polynesia and tropical America (Xiwen & Yan-hui 1990). It is a genus of small to medium sized trees which can be found from seashore to the lowland and up to the mountain forests. It is traditionally used in the treatment of abdominal pain, dysentery, diarrhea, infected wound, leucorrhoea, chronic ulcer and gonorrhoea (Balemba et al., 2010; Braide, 1993; Moongkarndi et al., 2004a).

1.4 Botanical Characterization of *Garcinia*

Trees in this genus are either dioecious or polygamous. The flowers may be solitary, fascicled and umbelled or paniced, usually with four to five decussate or imbricate and free sepals, and four to five imbricate petals. The male flowers have two to four lobed free or joined stamens. Anthers are straight with a horse shoe shape. Female flowers have 2-12 celled ovaries. The stigma is visible and sessile.

The berry is covered by tough peel and contains numerous large seeds suspended in a pulpy inner part (Xiwen & Yan-hui, 1990).

1.5 *Garcinia hombroniana* Pierre

G. hombroniana is widely distributed and is native to Peninsular Malaysia where it can be found in the coastal regions, from the lowland forests near the sea to the lower mountain forests and the highlands, upper parts of Borneo, Cambodia, Thailand, Andaman, Nicobar Islands and Vietnam. The name *Garcinia hombroniana* was given by Pierre (1882-1885), after a French physician J. B. Hombron, who had collected this plant while travelling from Singapore to Malacca in the Peninsular Malaysia (Nazre, 2010).

1.6 Botanical Characterization of *G. hombroniana*

G. hombroniana, a seashore mangosteen (English) or manggis hutan (Malay) is an evergreen fast growing medium sized dioecious tree, reaching to a height of 4.6-6.0 m and 1.8 m girth (Figure 1.1). It is the closest species to mangosteen (*G. mangostana*), and is thought to be one of its ancestors but interestingly the flowering time is much more different from that of mangosteen (Hammer, 2001). Due to the physical similarities of these two species, sometimes the nurseries mistakenly sell the seedlings as those of mangosteen. *G. hombroniana* has straight stem with horizontal and opposite dense branches. Young twigs are smooth and green (Figure 1.2a) but the older bark is dark brown and rough with white latex (Figure 1.2b). Leaves are bright yellowish green which become glossy green on maturity (Figure 1.2c).



Figure 1.1. Garcinia hombroniana Pierre

The fruit (Figure 1.2d) is smooth and spherical with a pinkish red peel. The internal part is segmented, with yellowish pulp and has a good flavour. The flowers (Figure 1.2e) are whitish, with four sepals and petals, and 2.5 cm in diameter. The berry (Figure 1.2f) is ash-green in the beginning but turns crimson red on ripening. The pericarp (Figure 1.2d) is relatively thin. The seeds are about 6-8 in number and the ripened berry is full with soft, sour, juicy and pale creamish arillus of a peach flavour (John et al., 2008).

1.7 Uses of *G. hombroniana*

The fruit of *G. hombroniana* is aromatic with thin and sour flesh. It is used in making juices and jellies. The timber is used to make houses and oars. The roots and leaves are used medicinally to relieve itching and as a protective medicine after childbirth (Pierre, 1882-1885).



(a) Young twig




(b) Bark



(c) Leaves



(d) Fruit and pericarp



(e) Flower



(f) Unripened berry

Figure 1.2. Parts of G. hombroniana

CHAPTER 2

LITERATURE REVIEW

2.1 Phytochemical Studies of *Garcinia*


Garcinia is a rich source of bioactive secondary metabolites including oxygenated prenylated xanthenes (Peres et al., 2000), benzophenones (Magadulla et al., 2008), flavonoids and biflavonoids (Harati et al., 2007), and triterpenes (Rukachaisirikul et al., 2005a). These phenolic compounds possess various pharmacological properties such as antioxidative, anti-inflammatory, antiallergic, antimicrobial, antimalarial, antihepatotoxic and HIV inhibitory activities (Gustafson et al., 1992; Hay et al., 2004).

2.2 Phytochemical Investigation of *G. hombroniana*

Previous phytochemical investigation of the different parts of *G. hombroniana* such as stem wood, pericarp, leaves and twigs had resulted in the isolation of different classes of compounds including xanthenes, flavonoids, biflavonoids and triterpenes. The chemical constituents isolated from *G. hombroniana* up to the year 2013 are reported in this thesis using Scifinder database.

2.2.1 Chemical Constituents Isolated from the Stem Wood of *G. hombroniana*


Bronianone (**1**) is the only compound isolated by Ollis et al. (1969) from the stem wood of *G. hombroniana*.




(1)

2.2.2 Chemical Constituents Isolated from the Pericarp of *G. hombroniana*


Rukachaisirikul et al. (2000) reported the isolation of triterpenes as the major compounds from dichloromethane extract of the pericarp of *G. hombroniana*. The compounds include (24*E*)-3 α -hydroxy-17,14-friedolanostan-8,14,24-dien-26-oic acid (2), garcihombronane B (3), garcihombronane C (4), garcihombronane D (5) and garcihombronane E (6).




(2)




(3)



(4)




(5)




(6)

2.2.3 Chemical Constituents Isolated from the Leaves of *G. hombroniana*


Rukachaisirikul et al. (2005a) further reported the isolation of triterpenoids along with some other classes of compounds from the leaves of *G. hombroniana*. The compounds include garcihombronane F (7), garcihombronane G (8), garcihombronane H (9), monoacetate of garcihombronane H (10), garcihombronane I (11), monoacetate of garcihombronane I (12), garcihombronane J (13), methyl (25*R*)-3β-hydroxy-23-ox-9,15-lanostadien-26-oate (14), vitexin (15), isovitexin (16), bluminol-C-9-*O*-β-D-apiofuranosyl-(1→6)-β-D-glucopyranoside (17), vomifoliol-9-*O*-β-D-apiofuranosyl (1→6)-β-D-glucopyranoside (18) and garcihombronanes B-E.




(7)




(8)




(9)




(10)




(11)




(12)




(13)




(14)




(15)



(16)



(17)



















(18)







2.2.4 Chemical Constituents Isolated from the Twigs of *G. hombroniana*

Klaiklay et al. (2013) reported triterpenes and other compounds from the twigs of *G. hombroniana*. The compounds include (22*Z*, 24*E*)-3β,9α-dihydroxy-

17,14-friedolanostan-14,22,24-trien-26-oic acid (**19**), garcihombronane K (**20**), Monoacetate of garcihombronane K (**21**) garcihombronane L (**22**), garcihombronone A (**23**), garcihombronone B (**24**), garcihombronone C (**25**), garcihombronone D (**26**), norathyriol (**27**), gentisein (**28**), 1,3,6,7-tetrahydroxy-8-prenylxanthone (**29**), cheffouxanthone (**30**), bangangxanthone A (**31**), toxyloxanthone B (**32**), 1,3,5,7-tetrahydroxy-2-(3,7-dimethyl-6-hydroxyocta-2-7-dien)xanthone (**33**)*, volkensiflavone (**34**), naringenin 7-*O*- β -D-glucuronide (**35**)*, eriodictyol 7-*O*- β -D-glucuronide (**36**)*, 4-hydroxybenzoic acid (**37**) and protocatechuic methyl ester (**38**). Saputri and Jantan (2012) also reported 3,5,3',5'-tetrahydroxy-4-methoxybenzophenone (**39**) and 1,7-dihydroxyxanthone (**40**) from the methanol extract of the twigs of *G. hombroniana*.







*Klaiklay PhD Thesis, 2009

2.3 Pharmacological Investigation of *G. hombroniana*

The *Garcinia* species are well known tropical trees used in the folk medicine for the treatment of various diseases and have been reported for their significant pharmacological properties. *G. hombroniana* is reported to have antiplatelet aggregation activities. Saputri and Jantan (2012) reported that the crude methanol extract of the twigs of *G. hombroniana* and the two compounds: (39) and (40) isolated from it exhibited strong copper mediated LDL antioxidation with IC_{50} of 6.6 and 1.7 μ M, respectively. It was further reported that compound (39) showed archidonic acid (AA, IC_{50} = 53.6 μ M), adenosine diphosphate (ADP, IC_{50} = 125.7

μM) and collagen-induced ($\text{IC}_{50} = 178.6 \mu\text{M}$) antiplatelet aggregation activities while compound (40) displayed a selective inhibition against ADP induced aggregation with IC_{50} of $5.7 \mu\text{M}$. The crude methanol extract of the twigs of *G. hombroniana* showed antibacterial activities against *S. aureus* and MRSA with the MIC values of 128 and $200 \mu\text{g/mL}$, respectively. Compounds (30) and (31) displayed moderate activities against MRSA with MIC values of 64.0 and $32.0 \mu\text{g/mL}$, respectively. In addition, Klaiklay et al. (2013) also reported that compound (30) has a good activity against *S. aureus* with a MIC value of $16.0 \mu\text{g/mL}$.

2.4 Problem Statement

The biodiversity source in Malaysia is one of the biggest in the world but its potential uses need to be identified and exploited. During the compilation of the economic products of the Malay Peninsula, Burkill (1966) has reported that 1,300 plants were used as folk medicine. The rich biodiversity of Malaysia, the public interest in herbal medicine and the contribution of industries and multinationals in herbal medicine have led to new emphasis of the medicinal plant research. Extensive work on the isolation and identification of chemical compounds from the *Garcinia* species, as well as their biological activities have been reported as many species from this genus are easily available and are known for their medicinal values. Previously the stem wood, twigs, pericarp and leaves of *G. hombroniana* have been investigated for their chemical constituents and biological activities (Klaiklay et al., 2013; Rukachaisirikul et al., 2005a; Saputri & Jantan, 2012). However, to the best of our knowledge, there is no authentic previous study on the bark of *G. hombroniana* either in terms of phytochemical investigation or bioactivities. Taking this into

account, this research work was aimed to isolate, purify and identify biologically active compounds from the bark of *G. hombroniana*.

2.5 Objectives of the Research

The objectives of this research are:

1. To isolate the chemical constituents from dichloromethane and ethyl acetate extracts of the bark of *G. hombroniana*.
2. To characterize and identify the isolated chemical constituents using different spectroscopic techniques such as UV, IR, NMR (1D and 2D) and mass spectrometry (EI-MS and HR/LR-ESI-MS).
3. To evaluate the phenolic and flavonoid contents and to correlate these contents to the antioxidant activities of extracts.
4. To evaluate the antibacterial, anticholinesterase and anticancer activities of the extracts and some selected pure compounds.

CHAPTER 3

MATERIALS AND METHODS

3.1 Plant Material

The plant (*G. hombroniana*) was collected in April 2010 from Penang Botanic Garden, Malaysia. A voucher specimen (PBGK12) has been deposited at the herbarium of this garden.

3.2 Chemicals

All the chemicals and solvents used for extraction, isolation and biological activities were either of analytical or HPLC grade.

3.2.1 Chemicals for Extraction and Isolation

Acetone, AR grade (QRĕC, Malaysia); acetone-*d*₆ for NMR 99.5 atom% D Nitrogen flushed (Merck, Germany); aluminium chloride AlCl₃.6H₂O (Merck, Germany); chloroform CHCl₃, AR grade (QRĕC, Malaysia); chloroform-*d*₁, with 0.03% TMS (v/v), 99.8 atom% D, stabilized with silver foil (Merck, Germany); DMSO-*d*₆ 99.8 atom% D (Merck, Germany); ethyl acetate EtOAc, AR grade, (QRĕC, Malaysia); ethanol 95% and 99% (QRĕC, Malaysia); methanol MeOH, AR grade (QRĕC, Malaysia); MeOD-*d*₄ 99.8 atom% D, (ACROS, USA); pyridine C₅D₅N (Merck, Germany); sephadex LH-20 (Sigma-Aldrich, USA); silica gel 60 for column chromatography (0.040-0.063 mm; 230-400 mesh, Merck, Germany); sodium carbonate Na₂CO₃, (System, Malaysia); sodium hydroxide NaOH,

(System, Malaysia); sulfuric acid H₂SO₄, (R & M Chemicals Essex, U.K); TLC aluminium sheets, pre-coated silica gel 60 F₂₅₄, 20 cm x 20 cm (Merck, Germany).

3.2.2 Chemicals for Biological Activities Evaluation

Acetylthiocholine iodide (ATCI), (Sigma-Aldrich, USA); 2,2'-azino-bis-3-ethyl benzthiazoline-6-sulphonic acid (ABTS), (Sigma Aldrich, Germany); buffer pH 3.6 (Sigma-Aldrich, USA); *S*-butyrylthiocholine chloride (BTCCl) (Sigma-Aldrich, USA); cytotoxicity detection kit (Roche Diagnostics, Germany); deionized water; dimethyl sulfoxide DMSO, (Sigma-Aldrich, USA); 5,5-dinitrobenzoic acid (DTNB), (Sigma- Aldrich, USA); 2,2-diphenyl-1-picrylhydrazyl (DPPH), (Sigma Aldrich, Germany); ferric chloride (FeCl₃.6H₂O), (Bendosen Laboratory Chemicals); 10% fetal bovine serum (Gibco, USA); Folin-Ciocalteu Reagent (Sigma Aldrich, Germany); gallic acid (Sigma Aldrich, Germany); glioma DBTRG cell lines [American Type Culture Collection (ATCC) Rockville, USA]; human breast cancer MCF-7 cell lines [American Type Culture Collection (ATCC), Rockville, USA]; hydrochloric acid HCl, (R & M Chemicals Essex, U.K); 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), (Sigma Aldrich, Germany); Muller-Hinton agar and broth MHA and MHB, (Oxoid, England), *p*-nitrotetrazolium violet (INT), (Sigma-Aldrich, USA); osteosarcoma U2OS cell lines, [American Type Culture Collection (ATCC), Rockville, USA]; potassium persulfate (K₂S₂O₈), (Merck, Germany); prostate PC3 cell lines, [American Type Culture Collection (ATCC), Rockville, USA]; physostigmine (Sigma-Aldrich, USA); quercetin (Aldrich Milwaukee, WI); rutin (Aldrich Milwaukee, WI); 2,4,6-tripyridyl-*s*-triazine TPTZ, (Merck, Germany).

3.3 Instruments Used for Structures Elucidation and Biological Activities

3.3.1 Melting Point

Melting points of the isolated pure compounds were determined by a Stuart Scientific SMP-1 (UK) melting point apparatus.

3.3.2 Infrared (IR) Spectroscopy

IR spectra of the isolated compounds were recorded on a Perkin-Elmer System 2000 FT-IR spectrometer (England, UK). The samples were prepared as potassium bromide (KBr) discs and scanned in a measurement range of 400-4000 cm^{-1} .

3.3.3 Nuclear Magnetic Resonance (NMR) Spectroscopy

NMR spectra of the isolated pure compounds were recorded on a Bruker Avance 500 spectrometer operated at 500 MHz for ^1H -NMR, and 125 MHz for ^{13}C -NMR, respectively. The measurements of the samples were performed in deuterated solvents (chloroform- d_1 , dimethyl sulfoxide- d_6 , methanol- d_4 , acetone- d_6 and pyridine- d_5). The coupling constants (J) were reported in Hertz (Hz).

3.3.4 Direct Probe Mass Spectrometry

Direct probe mass spectrometry through EI-MS and LR/HR-ESI-MS spectra were respectively recorded on an Agilent 5975C MSD and ThermoFinnigan MAT95XL mass spectrometers at the National University of Singapore (NUS), Singapore.

3.3.5 Ultraviolet (UV) Spectroscopy

UV spectra were recorded using a Perkin Elmer, Lambda 25 UV/Vis spectrometer in the region of 200-500 nm. The UV spectra for biflavonoids were measured using different shift reagents according to the method described by Bloor (2001). Stock solutions of the biflavonoids (**EN3-EN5**) were prepared by dissolving about 0.1 mg of compounds in 10 mL of methanol and divided each into three parts. The UV spectra were then measured using the following procedure.

- i. The UV spectra of the samples prepared in methanol were measured.
- ii. The spectra were recorded immediately after the addition of few drops of NaOMe (2.5 g metallic Na/100 mL MeOH) to first part of the stock solutions.
- iii. After 5 minutes the NaOMe spectra were recorded again.
- iv. The spectra were recorded immediately after the addition of 3 drops of 5% AlCl₃ reagent to second part of the stock solutions.
- v. The spectra were recorded immediately after the addition of 6 drops of 50% HCl reagent to the solutions of step (iv).
- vi. The spectra were measured after the addition and mixing of anhydrous sodium acetate (NaOAc) powder to third part of the stock solutions. The quantity of NaOAc added was such that about 2 mm layer settled to the bottom of the cuvette.
- vii. The spectra were recorded immediately after the addition of anhydrous boric acid powder to the solutions used in step (vi).

3.3.6 Circular Dichroism (CD) Spectroscopy

The absolute stereochemistry of **EN5** was determined using Jasco-815 spectrometer at 25 °C (Scan range $\lambda = 200\text{-}350$ nm, cell length = 10 mm).

3.3.7 X-Ray Crystallography Analysis

Single crystal X-Ray structure analysis of compound **EN2** was performed on a Bruker SMART APEXII CCD area detector diffractometer at the School of Physics, USM. The X-Ray crystallographic structure was solved and refined using SHELXTL software.

3.3.8 Microplate Reader

A Tecan Infinite 200 Pro Microplate spectrometer was used for the evaluation of the biological assays. Absorbance at different concentrations was measured with the microplate reader.

3.4 Extraction Procedure

The air dried ground bark (5.2 kg) of *G. hombroniana* was sequentially extracted using Soxhlet extractor with *n*-hexane, dichloromethane, chloroform, ethyl acetate and methanol at 40 °C. The filtered extracts were evaporated to dryness using a rotary evaporator at 40 °C to give a 30.1 g of *n*-hexane extract, 50.0 g dichloromethane extract, 5.0 g of a chloroform extract, 18.2 g of ethyl acetate extract and 30.1 g of a methanol extract. Another 100 g was extracted by the direct maceration into distilled hot water for 24 h to give aqueous-di extract (Figure 3.1).




Figure 3.1. Extraction procedure of *G. hombroniana* bark

3.5 Phytochemical Screening

The crude *n*-hexane, dichloromethane, chloroform, ethyl acetate, methanol and aqueous extracts, and the extract (aqueous-di) were screened for the detection of some secondary metabolites such as terpenoids, phytosterols, phenolics, flavonoids, carbohydrates, reducing sugars, alkaloids, phlobatannins and tannins, following the procedures of Indian Pharmacopoeia (1985).

3.5.1 Detection of Carbohydrates and Glycosides

Molish's Test: Few drops of α -naphthol were added to 2 mL filtrate of the crude extracts. After mixing and shaking, 1 mL of sulfuric acid was gradually added along the sides of the test tube and allowed to stand. A violet ring indicated the presence of carbohydrates.

Fehling Solution's Test: 1 mL of Fehling's solution was added to 2 mL filtrate of crude extracts and boiled at 100 °C. The formation of red precipitate indicated the presence of reducing sugars.

3.5.2 Detection of Proteins and Amino Acids

Biuret Test: A Few drops of 2% CuSO₄ solution were added to 5 mL filtrate of the crude extracts. Then 2 mL of 95% ethanol and excess of potassium hydroxide pellets were added. An absence of pink colour indicated an absence of proteins.

Ninhydrin Test: Few drops of ninhydrin solution (5% w/v in acetone) were added to 2 mL filtrate of the crude extracts. Appearance of no purple colour revealed the absence of amino acids.

3.5.3 Detection of Terpenoids (Salkowski's Test)

2 mL chloroform was added to 5 mL filtrate of the crude extracts, and then 3 mL of conc. H_2SO_4 was added slowly along the sides of the test tube. A reddish interface showed the presence of terpenoids.

3.5.4 Detection of Phytosterols (Liebermann-Burchard's Test)

2 mL of acetic anhydride was added to 5 mL filtrate of the crude extracts. Afterwards, 1-3 drops of conc. H_2SO_4 were added by the sides of the test tube. Array of the colour change showed the presence of phytosterols.

3.5.5 Detection of Phenolic Compounds

Ferric Chloride Test: A few drops of 5% ferric chloride solution were added to 2 mL filtrate of the crude extracts. Formation of dark green colour indicated the presence of phenolic compounds.

Lead Acetate Test: 3 mL of 10% lead acetate solution was added to 5 mL filtrate of the crude extracts. A white precipitate indicated the presence of phenolic compounds.

3.5.6 Detection of Flavonoids

NH_4OH Solution Test: 1 mL of 10% NH_4OH solution was added to 5 mL filtrate of the crude extracts. A formation of yellow florescent colour indicated the presence of flavonoids. Furthermore, 5 mL of DMSO was added to it, heated it, and

magnesium chloride and 1 mL of conc. HCl was added. Appearance of red colour showed the presence of flavonoids.

NaOH Solution Test: 4 mL of dil. NaOH solution was added to 2 mL filtrate of the crude extracts. The formation of golden yellow colour revealed the presence of flavonoids.

3.5.7 Detection of Tannins

A few drops of 5% ferric chloride were added to 5 mL filtrate of the crude extracts. A brown colour indicated the presence of tannins.

3.5.8 Detection of Phlobatannins

5 mL of deionized water was added to 5 mL filtrate of the crude extracts. Then a few drops of 1% HCl were added and boiled it at 100 °C for 10 min. The absence of formation of red precipitate indicated the absence of phlobatannins.

3.5.9 Detection of Alkaloids

A few drops of Wagner's reagent were added to 5 mL filtrate of the crude extracts. Orange precipitate showed the presence of alkaloids.

3.6 Separation Techniques

3.6.1 Thin Layer Chromatography (TLC)

The crude extracts were preliminary examined by thin layer chromatography, followed by the determination of the appropriate solvent systems. The selected

solvent systems were used for the column chromatographic separation of compounds in the crude extracts. The purity of the isolated compounds was also checked with TLC. In this phytochemical investigation, pre-coated TLC plates of silica gel 60 F₂₅₄ (20 x 20) on aluminium sheets were used. The developed plates were visualized with a UV light (Vilber Lourmet, multiband UV-254/365 nm) and/or by spraying them with 95% methanolic H₂SO₄ or the Libermann-Burchard's reagent (Horborne, 1998) followed by heating with a heat gun at 100-105 °C.

3.6.2 Column Chromatography (CC)

Column chromatography was carried out using silica gel (230-240 mesh, 0.040-0.060 mm) and Sephadex LH-20 (bead size 25-100 μ). Elution was carried out with a gradient solvent system of increasing polarity (*n*-hexane, chloroform, ethyl acetate, acetone and methanol) when silica gel was used as an absorbent. However, 100% methanol was used as an eluting solvent with Sephadex LH-20.

3.7 Isolation and Purification

3.7.1 Dichloromethane (DCM) Extract

The DCM extract (20.0 g) was subjected to silica gel CC. Elution was carried out using *n*-hexane, *n*-hexane-ethyl acetate, ethyl acetate-methanol in a polarity gradient of 10:0, 9:1, 8:2, 7:3, 6:4, 1:1, 4:6, 3:7, 2:8 and 0:10. Eluents were collected in 42 fractions DFA1 to DFA42. Fractions having similar TLC profiles were combined to give nine fractions (DFB1 to DFB9).

Fraction DFB1 (3.8 g) with a greenish white crystalline solid was rechromatographed on silica gel CC, eluted with *n*-hexane and ethyl acetate. Sub-