
UNIVERSITI SAINS MALAYSIA

Peperiksaan Kursus Semasa Cuti Panjang
Sidang Akademik 2002/2003

April 2003

ZCT 304E/3 - Keelektrikan dan Kemagnetan II

Masa : 3 jam

Sila pastikan bahawa kertas peperiksaan ini mengandungi **DUAABELAS** muka surat yang bercetak sebelum anda memulakan peperiksaan ini.


Jawab kesemua **LIMA** soalan. Pelajar dibenarkan menjawab semua soalan dalam bahasa Inggeris ATAU bahasa Malaysia ATAU kombinasi kedua-duanya.

1. (a) Suatu cakera bulat berjejari R mempunyai ketumpatan cas permukaan yang seragam σ . Carikan medan elektrik pada satu titik pada paksi cakera yang berjarak z dari satahnya. (8/20)
(b) Suatu silinder bulat yang tegak berjejari R dan panjang L diletakkan di sepanjang paksi z . Silinder tersebut mempunyai ketumpatan isipadu tak seragam yang diberikan dengan persamaan $\rho(z) = \rho_0 + \beta z$ merujuk kepada titik asalan pada pusat silinder. Carikan daya keatas satu titik cas q yang diletakkan pada pusat silinder tersebut. (*gunakan jawapan yang diperoleh dari bahagian (a)) (12/20)
2. Suatu cas q ditaburkan secara seragam pada keseluruhan suatu isipadu sferaan bukan pengkonduksi yang mempunyai jejari R .
(a) Tunjukkan bahawa keupayaan pada titik yang berjarak r dari pusat, di mana $r < R$, diberikan dengan persamaan.
$$V = \frac{q(3R^2 - r^2)}{8\pi\epsilon_0 R^3} \quad (15/20)$$

(b) Apakah keupayaan pada titik $r > R$? (5/20)

- 2 -

3. Dua petala konduktor sfera sepusat berjejari r_1 dan r_2 ditetapkan pada keupayaan φ_1 and φ_2 tiap-tiap satunya. Kawasan antara petala sfera tersebut di penuhi dengan suatu bahan dielektrik.
- (a) Dengan pengiraan terus, tunjukkan tenaga yang tersimpan dalam dielektrik adalah bersamaan dengan $\frac{C(\varphi_1 - \varphi_2)^2}{2}$. (5/20)
- (b) Tentukan C , kapasitans sistem tersebut diatas. (15/20)
-
4. Suatu kabel sepaksi yang panjang terdiri daripada dua konduktor sepusat dengan jejarinya seperti ditunjukkan dalam rajah dibawah. Kabel konduktor-konduktor tersebut membawa arus i yang magnitudnya adalah sama tetapi arahnya bertentangan. Tentukan medan magnet B di r jika
- (a) $r < a$, (5/20)
- (b) $a < r < b$, (5/20)
- (c) $b < r < c$ dan (5/20)
- (d) $r > c$ (di luar kabel) (5/20)



5. Tunjukkan keupayaan vektor kemagnetan untuk dua dawai panjang, lurus dan selari yang membawa arus I yang sama tetapi bertentangan arah diberikan oleh

$$\vec{A} = \frac{\mu_0 I}{2\pi} \ln\left(\frac{r_2}{r_1}\right) \hat{n},$$

dimana r_2 dan r_1 adalah jarak-jarak dari titik medan ke dawai-dawai berkenaan dan \hat{n} ialah vektor unit selari dengan dawai-dawai tersebut.

(20/20)

TERJEMAHAN

UNIVERSITI SAINS MALAYSIA

Third Semester Examination
2002/2003 Academic Session

April 2003

ZCT 304E/3 - Electricity and Magnetism II

Time : 3 hours

Please check that the examination paper consists of TWELVE printed pages before you commence this examination.

Answer all FIVE questions. Students are allowed to answer all questions in English OR bahasa Malaysia OR combinations of both.

1. (a) A circular disk of radius R has a uniform surface charge density σ . Find the electrical field at a point on the axis of the disk at a distance z from the plane of the disk. (8/20)

 (b) A right circular cylinder of radius R and height L is oriented along the z -axis. It has a nonuniform volume density of charge given by $\rho(z)=\rho_0 + \beta z$ with reference to an origin at the center of the cylinder. Find the force on a point charge q placed at the center of the cylinder. (*hint: use the answer obtained from part (a)) (12/20)

2. A charge q is distributed uniformly throughout a non-conducting spherical volume of radius R .

 (a) Show that the potential a distance r from the center, where $r < R$ is given by

$$V = \frac{q(3R^2 - r^2)}{8\pi\epsilon_0 R^3}$$
 (15/20)

 (b) What is the potential at a point $r > R$? (5/20)

3. Two concentric, spherical, conducting shells of radii r_1 and r_2 are maintained at potentials ϕ_1 and ϕ_2 respectively. The region between the shells is filled with a dielectric medium.

(a) Show by direct calculation that the energy stored in the dielectric is equal to $\frac{C(\phi_1 - \phi_2)^2}{2}$. (5/20)

(b) Determine C , the capacitance of the system. (15/20)


4. A long coaxial cable consists of two concentric conductors with the dimensions shown below. There are equal and opposite currents i in the conductors. Determine the magnetic field B at r if

(a) $r < a$, (5/20)

(b) $a < r < b$, (5/20)

(c) $b < r < c$ and (5/20)

(d) $r > c$ (outside the cable) (5/20)



5. Show that the magnetic vector potential for two long, straight, parallel wires carrying the same current, I in opposite directions is given by

$$\tilde{A} = \frac{\mu_0 I}{2\pi} \ln\left(\frac{r_2}{r_1}\right) \hat{n},$$

where r_2 and r_1 are the distances from the field point to the wires, and \hat{n} is a unit vector parallel to the wires.

(20/20)

Mathematical Guidance

Possibly Useful Integrals:

$$\int_{-1}^1 \frac{(z-r\mu)d\mu}{(r^2+z^2-2zr\mu)^{3/2}} = \frac{1}{z^2} \left(\frac{z-r}{|z-r|} + \frac{z+r}{|z+r|} \right)$$

$$\int_{-1}^1 \frac{d\mu}{(r^2+z^2-2zr\mu)^{1/2}} = \frac{1}{zr} (\lfloor z+r \rfloor - \lfloor z-r \rfloor)$$

$$\int \frac{dx}{(x^2+a^2)^{3/2}} = \frac{1}{a^2} \cdot \frac{x}{(x^2+a^2)^{1/2}}$$

$$\int x e^{ax} dx = e^{ax} \left[\frac{x}{a} - \frac{1}{a^2} \right]$$

Useful Constants

$$k = \frac{1}{4\pi\epsilon_0} = 8.99 \times 10^9 \frac{N \cdot m^2}{C^2} \quad e = 1.60 \times 10^{-19} C$$

$$\epsilon_0 = 8.85 \times 10^{-12} \frac{C^2}{N \cdot m^2} \quad \mu_0 = 4\pi \times 10^{-7} \frac{T \cdot m}{A}$$

LAMPIRAN

Vector Calculus**Cartesian Coordinates**

$$\vec{\nabla} u = \hat{x} \frac{\partial u}{\partial x} + \hat{y} \frac{\partial u}{\partial y} + \hat{z} \frac{\partial u}{\partial z}$$

$$\vec{\nabla} \cdot \vec{A} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}$$

$$\vec{\nabla} \times \vec{A} = \hat{x} \left(\frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + \hat{y} \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) + \hat{z} \left(\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right)$$

$$d\tau = dx dy dz \quad da_x = \pm dy dz \quad da_y = \pm dx dz \quad da_z = \pm dx dy$$

Cylindrical Coordinates

$$\vec{\nabla} u = \hat{\rho} \frac{\partial u}{\partial \rho} + \hat{\phi} \frac{1}{\rho} \frac{\partial u}{\partial \phi} + \hat{z} \frac{\partial u}{\partial z}$$

$$\vec{\nabla} \cdot \vec{A} = \frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\rho) + \frac{1}{\rho} \frac{\partial A_\phi}{\partial \phi} + \frac{\partial A_z}{\partial z}$$

$$\vec{\nabla} \times \vec{A} = \hat{\rho} \left(\frac{1}{\rho} \frac{\partial A_z}{\partial \phi} - \frac{\partial A_\phi}{\partial z} \right) + \hat{\phi} \left(\frac{\partial A_\rho}{\partial z} - \frac{\partial A_z}{\partial \rho} \right) + \hat{z} \left[\frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\phi) - \frac{1}{\rho} \frac{\partial A_\rho}{\partial \phi} \right]$$

$$d\tau = \rho d\rho d\phi dz \quad da_\rho = \pm \rho d\phi dz \quad da_\phi = \pm d\rho dz \quad da_z = \pm \rho d\rho d\phi$$

$$\hat{\rho} = \cos \phi \hat{x} + \sin \phi \hat{y} \quad \hat{\phi} = -\sin \phi \hat{x} + \cos \phi \hat{y}$$

Spherical Coordinates

$$\vec{\nabla} u = \hat{r} \frac{\partial u}{\partial r} + \hat{\theta} \frac{1}{r} \frac{\partial u}{\partial \theta} + \hat{\phi} \frac{1}{r \sin \theta} \frac{\partial u}{\partial \phi}$$

$$\vec{\nabla} \cdot \vec{A} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 A_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta A_\theta) + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi}$$

$$\vec{\nabla} \times \vec{A} = \frac{\hat{r}}{r \sin \theta} \left[\frac{\partial}{\partial \theta} (\sin \theta A_\phi) - \frac{\partial A_\theta}{\partial \phi} \right] + \frac{\hat{\theta}}{r} \left[\frac{1}{\sin \theta} \frac{\partial A_r}{\partial \phi} - \frac{\partial}{\partial r} (r A_\phi) \right] + \frac{\hat{\phi}}{r} \left[\frac{\partial}{\partial r} (r A_\theta) - \frac{\partial A_r}{\partial \theta} \right]$$

$$d\tau = r^2 \sin \theta dr d\theta d\phi \quad da_r = \pm r^2 \sin \theta d\theta d\phi \quad da_\theta = \pm r \sin \theta dr d\phi \quad da_\phi = \pm r dr d\theta$$

$$\hat{r} = \sin \theta \cos \phi \hat{x} + \sin \theta \sin \phi \hat{y} + \cos \theta \hat{z} \quad \hat{\theta} = \cos \theta \cos \phi \hat{x} + \cos \theta \sin \phi \hat{y} - \sin \theta \hat{z}$$

$$\hat{\phi} = -\sin \phi \hat{x} + \cos \phi \hat{y}$$

LAMPIRAN**Important Equations****Maxwell's Equations:**

$$\bar{\nabla} \cdot \bar{D} = \rho_f \quad \bar{\nabla} \cdot \bar{B} = 0 \quad \bar{\nabla} \times \bar{E} = -\frac{\partial \bar{B}}{\partial t} \quad \bar{\nabla} \times \bar{H} = J_f + \frac{\partial \bar{D}}{\partial t}$$

Lorentz Force: $\vec{F} = q(\vec{E} + \vec{v} \times \vec{B})$

Equation of Continuity: $\bar{\nabla} \cdot \bar{J}_f + \frac{\partial \rho_f}{\partial t} = 0$

Coulomb's Law: $\vec{F}_q = \sum_i \frac{qq_i \vec{R}_i}{4\pi\epsilon_0 R_i^3}$ (for a collection of point charges)

$$\vec{F}_q = \frac{q}{4\pi\epsilon_0} \int_L \frac{\lambda(\vec{r}') \vec{R} ds'}{R^3} \quad (\text{for a line charge distribution})$$

$$\vec{F}_q = \frac{q}{4\pi\epsilon_0} \int_S \frac{\sigma(\vec{r}') \vec{R} da'}{R^3} \quad (\text{for a surface charge distribution})$$

$$\vec{F}_q = \frac{q}{4\pi\epsilon_0} \int_V \frac{\rho(\vec{r}') \vec{R} d\tau'}{R^3} \quad (\text{for a volume charge distribution})$$

Electric Field: $\vec{E} = \frac{\vec{F}_q}{q}$

Electric Flux: $\Phi_e = \int \vec{E} \cdot d\vec{a}$

Gauss' Law: $\oint_S \vec{E} \cdot d\vec{a} = \frac{Q_t}{\epsilon_0}$ (integral form)

$$\bar{\nabla} \cdot \bar{E} = \frac{\rho(\vec{r})}{\epsilon_0} \quad (\text{differential form})$$

LAMPIRAN

Scalar Potential: $\phi(\vec{r}) = \sum_i \frac{q_i}{4\pi\epsilon_0 R_i}$ (for a collection of point charges)

$$\phi(\vec{r}) = \frac{1}{4\pi\epsilon_0} \int_L \frac{\lambda(\vec{r}')ds'}{R} \quad (\text{for a line charge distribution})$$

$$\phi(\vec{r}) = \frac{1}{4\pi\epsilon_0} \int_S \frac{\sigma(\vec{r}')da'}{R} \quad (\text{for a surface charge distribution})$$

$$\phi(\vec{r}) = \frac{1}{4\pi\epsilon_0} \int_V \frac{\rho(\vec{r}')d\tau'}{R} \quad (\text{for a volume charge distribution})$$

Potential Energy: $U_e(\vec{r}) = q\phi(\vec{r})$ (for an isolated point charge)

$$U_e = \frac{1}{2} \sum_i q_i \phi_i(\vec{r}_i) \quad (\text{for a collection of point charges})$$

$$U_e = \frac{1}{2} \int_L \lambda(\vec{r}) \phi(\vec{r}) ds \quad (\text{for a line charge distribution})$$

$$U_e = \frac{1}{2} \int_S \sigma(\vec{r}) \phi(\vec{r}) da \quad (\text{for a surface charge distribution})$$

$$U_e = \frac{1}{2} \int_V \rho(\vec{r}) \phi(\vec{r}) d\tau \quad (\text{for a volume charge distribution})$$

$$u_e = \frac{1}{2} \epsilon_0 E^2 \quad (\text{energy density in an electric field})$$

$$U_e = \int u_e d\tau \quad (\text{total energy})$$

Multipole Moments: $Q = \sum_i q_i$ or $Q = \int_L \lambda ds$ or $Q = \int_S \sigma da$ or $Q = \int_V \rho d\tau$ (monopole)

$$\bar{p} = \sum_i q_i \vec{r}_i \quad \text{or} \quad \bar{p} = \int_L \lambda \vec{r} ds \quad \text{or} \quad \bar{p} = \int_S \sigma \vec{r} da \quad \text{or} \quad \bar{p} = \int_V \rho \vec{r} d\tau \quad (\text{dipole})$$

Boundary Conditions: $E_{t2} - E_{t1} = 0$ and $E_{n2} - E_{n1} = \frac{\sigma}{\epsilon_0}$ (electric field)
 $\phi_2 = \phi_1$ (scalar potential)

$$B_{n2} - B_{n1} = 0 \quad \text{and} \quad \bar{B}_{t2} - \bar{B}_{t1} = \mu_0 \bar{K} \times \hat{n} \quad (\text{magnetic induction})$$

LAMPIRAN

Electricity in Matter: $\rho = \rho_f + \rho_b$ (free charge and bound charge)

$$\rho_b = -\vec{\nabla} \cdot \vec{P} \text{ and } \sigma_b = \vec{P} \cdot \hat{n} \text{ (bound charge densities)}$$

$$\vec{D} = \epsilon_0 \vec{E} + \vec{P} \text{ (definition of electric displacement)}$$

$$\vec{D} = \kappa_e \epsilon_0 \vec{E} = \epsilon \vec{E} \text{ (for an l.i.h. dielectric)}$$

$$u_e = \frac{1}{2} \vec{D} \cdot \vec{E} \text{ (energy density in matter)}$$

$$\oint \vec{D} \cdot d\vec{a} = Q_{f,in} \text{ and } \vec{\nabla} \cdot \vec{D} = \rho_f \text{ (Gauss' Laws for } \vec{D})$$

Electric Current: $I = \frac{dq}{dt} = \int \vec{J} \cdot d\vec{a} = \int \vec{K} \cdot d\vec{s}$

$$\vec{J} = \rho \vec{v} \quad \vec{K} = \sigma \vec{v} \quad \text{(current density)}$$

$$Id\vec{s} = \vec{K}da = \vec{J}d\tau \quad \text{(current elements)}$$

$$\vec{J}_f = \sigma \vec{E} \quad \text{(Ohm's Law)}$$

Magnetostatic Force: $\vec{F}_{C \rightarrow C'} = \frac{\mu_0}{4\pi} \oint_C \oint_{C'} \frac{Id\vec{s} \times (I'd\vec{s}' \times \hat{R})}{R^2}$

Magnetic Induction: $\vec{B} = \frac{\mu_0}{4\pi} \oint_C \frac{I'd\vec{s}' \times \hat{R}}{R^2} \quad \text{(for a filamentary current)}$

$$\vec{B} = \frac{\mu_0}{4\pi} \int_S \frac{\vec{K}' \times \hat{R}da'}{R^2} \quad \text{(for a surface current)}$$

$$\vec{B} = \frac{\mu_0}{4\pi} \int_V \frac{\vec{J}' \times \hat{R}d\tau'}{R^2} \quad \text{(for a volume current)}$$

Ampere's Law: $\oint_C \vec{B} \cdot d\vec{s} = \mu_0 I_i \quad \text{(integral form)}$

$$\vec{\nabla} \times \vec{B} = \mu_0 \vec{J} \quad \text{(differential form)}$$

LAMPIRAN

Vector Potential: $\vec{B} = \vec{\nabla} \times \vec{A}$

$$\vec{A} = \frac{\mu_0}{4\pi} \oint_C \frac{I' d\vec{s}'}{R} \quad (\text{for a filamentary current})$$

$$\vec{A} = \frac{\mu_0}{4\pi} \int_S \frac{\vec{K}' da'}{R} \quad (\text{for a surface current})$$

$$\vec{A} = \frac{\mu_0}{4\pi} \int_V \frac{\vec{J}' d\tau'}{R} \quad (\text{for a volume current})$$

Magnetic Flux: $\Phi_b = \int \vec{B} \cdot d\vec{a}$

Faraday's Law: $\varepsilon_t = \oint \vec{E}_t \cdot d\vec{s} = \frac{-d\Phi_b}{dt} \quad (\text{integral form})$

$$\vec{\nabla} \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \quad (\text{differential form})$$

Magnetism in Matter: $\vec{J} = \vec{J}_f + \vec{J}_m \quad (\text{free current plus magnetisation current})$

$$\vec{J}_m = \vec{\nabla} \times \vec{M} \quad (\text{magnetisation volume current density})$$

$$\vec{K}_m = \vec{M} \times \hat{n} \quad (\text{magnetisation surface current density})$$

$$\vec{B} = \mu_0(\vec{H} + \vec{M}) \quad (\text{definition of magnetic field})$$

$$\vec{B} = \mu_0(\vec{H} + \vec{M}) = \mu_0(1 + \chi_m)\vec{H} = \mu\vec{H} \quad (\text{for l.i.h. material})$$