DEVELOPMENT, IMPLEMENTATION AND EVALUATION OF NUTRITION MODULE FOR TRAINERS OF PERSONS WITH DISABILITIES IN COMMUNITY-BASED REHABILITATION CENTRES, KELANTAN

by

CHEN SEONG TING

Thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy

February 2014

ACKNOWLEDGEMENT

First of all, I would like to express my deepest appreciation and gratitude to my principal supervisor, Dr. Sakinah Harith, for her invaluable guidance, understanding and encouragement throughout the whole research process and this thesis writing. I also wish to express my sincere appreciation and thanks to my three co-supervisors, Professor Hans Van Rostenberghe, Dr. Azriani Ab Rahman and Dr. Soo Kah Leng, for their relentless effort, expert advice and support in my doctoral journey.

I acknowledge Universiti Sains Malaysia for providing me the financial support under USM Fellowship Scheme. Also, for funding my research project through the Research University Grant (1001/PPSP/812049), Postgraduate Research Grant Scheme (1001/PPSK/8144001) and grants from Division of Industry and Community Network (BJIM).

I offer special thanks to all persons with disabilities, their families as well as the CBR teachers from Kelantan, Terengganu, Perlis, Pulau Pinang and Perak who had involved directly or indirectly in this study. Many thanks for their cooperation and support throughout the study. I am deeply grateful to the Sustainable Optimal Services for The Disabled (S.O.S OKU) research team members, staffs of Dietetics and Nutrition Programmes, and the statisticians for their assistance and guidance.

I would express my heartfelt appreciation to all my good friends, in particular Miss Loy See Ling, Ms. Nur Nadia Mohamed and Miss Tan Sze Lin for their moral support and countless help offered to me in a myriad of ways. My utmost gratitude always goes to the most special person in my life, Ms. Seow Choy Ha, my beloved mother and best friend, for her endless love, encouragement and trust in me. Special thanks to my father (Mr. Chen Ker Yam) and sister (Ms. Chen Seong Voon) for their compassionate support. Last but not least to my dearest husband, Mr. Chung Weng Hong, who always provides me the strongest inspiration and motivation. Thank you for your understanding, support and love in this journey. I dedicate this thesis to all my loved ones.

TABLE OF CONTENTS

ACKNO	OWLE	DGEMENT	ii	
TABLE OF CONTENTS iv				
LIST O	F TAE	BLES	ix	
LIST O	F FIGI	URES	xi	
LIST O	F SYN	BOLS AND ABBREVIATIONS x	iii	
ABSTR	AK	x	vi	
ABSTR	ACT.	xv	iii	
СНАРТ	ER 1	INTRODUCTION	. 1	
1.1	Back	ground of the study	. 1	
1.2	State	ment of the problem	. 3	
1.3	Signi	ficance of the study	. 5	
1.4	Conc	eptual framework	. 6	
1.5	Meth	odological framework	. 6	
1.6	Resea	arch questions	. 9	
1.7	Obje	ctives of the study	10	
1.′	7.1	General objective	10	
1.1	7.2	Specific objectives	10	
1.8	Нуро	theses	11	
1.9	Defin	ition of terms	12	
CHAPT	ER 2	LITERATURE REVIEW	13	
2.1	Disat	bility- size of the problem	13	
2.2	Com	munity-based rehabilitation	14	
2.3	Disat	bilities and nutritional concerns	19	
2	3.1	Down's syndrome	20	
2	3.2	Cerebral palsy	22	
2.	3.3	Autism spectrum disorders	23	
2.	3.4	Intellectual disability	26	
2.4	Maln	utrition	28	
2.4	4.1	General population	28	
2.4	4.2	Persons with disabilities	29	
	2.4.2	.1 Disability-specific growth reference	35	
2.5	Feedi	ing and nutrition problems in persons with disabilities	40	
2.6	Nutri	tion module for persons with disabilities	45	

	2.7	Beha	vioural models in nutrition intervention	50
	2	.7.1	Social Cognitive Theory	51
	2.8	Knov	vledge, attitude and practice (KAP)	54
	2.9	Kirkŗ	patrick's Four Levels of Evaluation Model	57
	2.10	Nutri	tion intervention for caregivers	62
	2	.10.1	Nutrition intervention for primary caregivers	63
	2	.10.2	Nutrition intervention for paraprofessionals	66
	2.11	Facto	ors affecting nutrition practice of caregivers	69
	2	.11.1	Personal factors	69
	2	.11.2	Environmental factors	71
CI	HAP	TER 3	MATERIALS AND METHODS	74
	3.1	Kela	ntan: The geographical and population background	74
	3.2	Ethic	al consideration	76
	3.3	Phase	e 1: Nutrition assessment of persons with disabilities	76
	3	.3.1	Study design and setting	76
	3	.3.2	Sample size calculation	78
	3	.3.3	Study participants	80
	3	.3.4	Sampling method and recruitment	80
	3	.3.5	Research instruments	81
	3	.3.6	Data collection	83
	3	.3.7	Data analysis	90
	3.4	Phase with Ques	e 2: Development of Nutrition Module for Trainers of Persons Disabilities & Nutrition Knowledge, Attitude and Practice tionnaire about Persons with Disabilities (KAP-nOKU)	94
	3	.4.1	Nutrition Module for Trainers of Persons with Disabilities	95
		3.4.1	.1 Module development	95
		3.4.1	.2 Content validation	99
	3	.4.2	Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU) 1	00
		3.4.2	.1 Questionnaire development 1	00
		3.4.2	.2 Validation and reliability testing of KAP-nOKU 1	05
	3.5	Phase	e 3: Evaluation of nutrition module training 1	14
	3	.5.1	Study design and setting 1	15
	3	.5.2	Sample size calculation 1	18
	3	.5.3	Recruitment of participants 1	19
	3	.5.4	Study procedures 1	20

		3.5.4.	1	Research instruments	120
3.5.4.2		2	Intervention group	121	
3.5.4.3		3	Comparison group	123	
		3.5.4.	4	Data collection	127
	3.5	5.5	Dat	a analysis	127
	3.5	5.6	The	e qualitative approach	128
		3.5.6.	1	Qualitative research method	129
		3.5.6.	.2	Sampling method and recruitment	129
		3.5.6.	.3	Data collection method	131
		3.5.6.	.4	Research instrument- the interviewer and interview protocol	131
		3.5.6	.5	Study procedures and data collection	134
		3.5.6	.6	Data analysis	135
		3.5.6.	.7	Validity and reliability of study	138
CHA	N PT	ER 4	RE	SULTS	140
4.	.1	Phase	e 1:1	Nutrition assessment of persons with disabilities	140
	4.]	1.1	Soc	io-demographic characteristics	140
	4.1	1.2	Fee	ding characteristics	142
	4.1	.3	Die	tary intake	142
	4.1	.4	Ant	hropometric characteristics	149
	4.1	.5	Prev	valence of malnutrition	157
4.2	2	Phase with Quest	2: Dis tionr	Development of Nutrition Module for Trainers of Persons abilities & Nutrition Knowledge, Attitude and Practice naire about Persons with Disabilities (KAP-nOKU)	161
	4.2	2.1	Nut	rition Module for Trainers of Persons with Disabilities	161
		4.2.1.	1	Module content	161
		4.2.1.	2	Content validation	161
	4.2	2.2	Nut Pers	rition Knowledge, Attitude and Practice Questionnaire about sons with Disabilities (KAP-nOKU)	164
		4.2.2.	1	Content validity	164
		4.2.2.	2	Face validity	166
		4.2.2.	3	Item analysis, construct validity and reliability testing	166
4.	3	Phase	: 3: E	Evaluation of nutrition module training	178
	4.3	3.1	Lev	el 1- Reaction	178
	4.3	3.2	Lev	el 2- Learning and Level 3- Behaviour	180
		4.3.2.	1	Baseline data	180
		4.3.2.	2	Effectiveness of nutrition module training	181

4.3.2.3 Correlations among knowledge, attitude and practice			
4.3.3 Qualitative findings 189			
4.3.3.1 Socio-demographic characteristics			
4.3.3.2 Thematic building: Cohen's Kappa agreement			
4.3.3.3 Description on the qualitative findings			
CHAPTER 5 DISCUSSION			
5.1 Nutrition assessment of persons with disabilities			
5.1.1 General characteristics of participants			
5.1.2 Feeding and nutrition problems			
5.1.3 Dietary intake 218			
5.1.4 Anthropometry 221			
5.1.5 Prevalence of malnutrition			
5.2 Nutrition Module for Trainers of Persons with Disabilities			
5.3 Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU)			
5.3.1 Item analysis for knowledge items: Difficulty and discrimination indices			
5.3.2 Construct validity for attitude and practice items			
5.3.3 Reliability testing: Internal consistency and test-retest reliability 232			
5.4 Effectiveness of nutrition module training			
5.5 Perceived motivations and barriers to nutrition practice			
5.5.1 Personal factors			
5.5.2 Environmental factors			
5.6 Strengths of the study			
5.7 Limitations of the study 248			
5.8 Recommendation for future research			
CHAPTER 6 SUMMARY AND CONCLUSION			
REFERENCES			

APPENDICES

Appendix A	Ethical approval letters from Human Research Ethics Committee of Universiti Sains Malaysia				
Appendix B	Ethical approval letters from Malaysia Department of Social Welfare				
Appendix C	Written informed consents				
Appendix D	Questionnaire on nutrition assessment of persons with disabilities				
Appendix E	Content validation form for Nutrition Module for Trainers of Persons with Disabilities				
Appendix F	Content validation form for Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU)				
Appendix G	Validated Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU)				
Appendix H	Training evaluation rating sheet				
Appendix I	Qualitative semi-structured interview protocol				
Appendix J	Content validation form for qualitative semi-structured interview protocol				

LISTS OF PUBLICATIONS

LIST OF TABLES

Table 2.1	Number of CBR centres, trainers and PWD registered in Malaysia16
Table 2.2	Disability categories 17
Table 2.3	Prevalence of malnutrition for children and adolescents with disabilities in different countries
Table 2.4	Prevalence of malnutrition for adults with disabilities in different countries
Table 2.5	Definition on the five groups of functional ability for population with cerebral palsy
Table 3.1	List of study locations for Phase 1 study78
Table 3.2	Age- and sex-specific body density equation
Table 3.3	Body fatness level classification for adults with disabilities
Table 3.4	BMI-for-age category for children and adolescent with disabilities
Table 3.5	List of study locations for Phase 2 validation study 109
Table 3.6	Guideline for the discrimination index 113
Table 3.7	List of CBR centres that participated in Phase 3 study 117
Table 3.8	Summary of nutrition module training 124
Table 3.9	Timetable for nutrition module training (intervention group) 125
Table 3.10	Timetable for general health training (comparison group) 126
Table 3.11	Participants for the qualitative approach
Table 4.1	Socio-demographic characteristics of PWD141
Table 4.2	Feeding characteristics of PWD143
Table 4.3	Energy intake and percentage contribution of energy from macronutrients for children with disabilities ^a
Table 4.4	Energy intake and percentage contribution of energy from macronutrients for adolescents with disabilities ^a
Table 4.5	Energy intake and percentage contribution of energy from macronutrients for adults with disabilities ^a
Table 4.6	Achievement of mean energy and protein intake of PWD to RNI ^a 149
Table 4.7	Achievement of mean energy intake of PWD to RNI according to sex, age category and body weight status
Table 4.8	Anthropometric characteristics of PWD according to sex and age category154
Table 4.9	Anthropometric status of PWD 156

Table 4.10	Topics, contents and learning outcomes of Nutrition Module for Trainers of Persons with Disabilities162
Table 4.11	Amendment on the Malay terms based on experts' suggestions 164
Table 4.12	Content validity index of KAP-nOKU 166
Table 4.13	Amendment of questionnaire based on experts' comments 167
Table 4.14	Face validation respondents' comments and modifications on the general format of questionnaire
Table 4.15	Descriptive statistics for knowledge, attitude and practice domains of KAP-nOKU
Table 4.16	Difficulty index, discrimination index and their interpretations for knowledge items of KAP-nOKU (n=165)
Table 4.17	Amendment on knowledge items based on item analysis 173
Table 4.18	Factor loading, corrected items-to-total score correlation and internal consistency for attitude domain (n=165)
Table 4.19	Factor loading, corrected items-to-total score correlation and internal consistency for practice domain (n=165)
Table 4.20	Summary of construct validity and reliability analysis for KAP- nOKU (n=165)
Table 4.21	Evaluation of teachers' acceptance towards the nutrition module training (n=45)
Table 4.22	Socio-demographic characteristics of participants (n=87) 181
Table 4.23	Comparisons of baseline knowledge, attitude and practice scores between intervention and comparison groups (n=87) 181
Table 4.24	Comparisons of knowledge, attitude and practice scores within each group based on time
Table 4.25	Mean differences in knowledge, attitude and practice scores between intervention and comparison groups
Table 4.26	Comparisons of knowledge, attitude and practice scores between groups based on time
Table 4.27	Correlations among knowledge, attitude and practice scores of intervention group at 6-month after training
Table 4.28	Experts' level of agreement on thematic building 190
Table 4.29	Summary of main themes for qualitative findings 191

LIST OF FIGURES

Figure 1.1	Conceptual framework of the study7					
Figure 1.2	Methodological framework of the study 8					
Figure 2.1	Social Cognitive Theory Model 52					
Figure 2.2	Knowledge-attitude-practice and Social Cognitive Theory models					
Figure 2.3	Kirkpatrick's Four Levels of Evaluation Model 58					
Figure 3.1	Maps of Peninsular Malaysia and Kelantan75					
Figure 3.2	Flow chart for Phase 1 study77					
Figure 3.3	Anthropometric equipments					
Figure 3.4	Flow chart for Phase 2 study					
Figure 3.5	Constructs for knowledge, attitude and practice domains 106					
Figure 3.6	Flow chart for Phase 3 study 116					
Figure 3.7	Number of participants for the intervention study 120					
Figure 3.8	Theme formulation 137					
Figure 4.1	Achievement of mean energy intake to RNI (%) according to age category and body weight status for males					
Figure 4.2	Achievement of mean energy intake to RNI (%) according to age category and body weight status for females					
Figure 4.3	Prevalence of malnutrition among PWD158					
Figure 4.4	Prevalence of malnutrition among children and adolescents with disabilities according to type of disability (n=276) 158					
Figure 4.5	Prevalence of malnutrition among adults with disabilities according to type of disability (n=186)					
Figure 4.6	Prevalence of malnutrition among children and adolescents with disabilities according to ambulatory status (n=276) 159					
Figure 4.7	Prevalence of malnutrition among adults with disabilities according to ambulatory status (n=186)					
Figure 4.8	Nutrition Module					
Figure 4.9	Adjusted mean knowledge scores at baseline, 1-month and 6- month after training					
Figure 4.10	Adjusted mean attitude scores at baseline, 1-month and 6-month after training					
Figure 4.11	Adjusted mean practice scores at baseline, 1-month and 6-month after training					
Figure 4.12	Healthy and variety food preparation for PWD 195					
Figure 4.13	Food modifications for children with feeding difficulties 196					

Figure 4.14	Persuasive method to tackle picky eater	198
Figure 4.15	Nutrition teaching towards PWD	200
Figure 4.16	Increased physical activity for the PWD	202
Figure 4.17	Food sources from growing produce of PWD	205
Figure 4.18	Positive food behavioural changes in PWD	207
Figure 4.19	Inadequate portion size of fruit due to lack of budget	209
Figure 4.20	Parents' unreceptive attitude towards nutrition advice	211

LIST OF SYMBOLS AND ABBREVIATIONS

ADA	American Dietetic Association
ADHD	Attention deficit hypersensitivity disorder
AIFO	Italian Association Amici di Raoul Follereau
α	Cronbach's alpha
ANCOVA	Analysis of covariance
ASD	Autism spectrum disorders
BAKTI	Badan Amal dan Kebajikan Tenaga Isteri-Isteri
BMI	Body mass index
BMR	Basal metabolic rate
CBR	Community-based rehabilitation
CDC	Centres for Disease Control
CI	Confidence interval
CITC	Corrected items-to-total score correlation
СР	Cerebral palsy
D	Discrimination index
dB	Decibel
EI	Energy intake
F	Female
FAO	Food and Agriculture Organization
Fr	Factor
GDP	Gross domestic product
GERD	Gastro-esophageal reflux disease
HSG	High scorer group
ICC	Intraclass correlation coefficient
ID	Intellectual disability
IDF	International Diabetes Federation

IPH	Institute for Public Health					
JIFSAN	Joint Institute of Food Safety and Applied Nutrition					
ЈКМ	Jabatan Kebajikan Masyarakat					
KAP	Knowledge, attitude and practice					
KAP-nOKU	Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities					
KMO	Kaiser-Meyer-Olkin					
KR	Kuder-Richardson					
LSG	Low scorer group					
Μ	Male					
МОН	Ministry of Health					
MUAC	Mid-upper arm circumference					
n	Frequency					
N/A	Non-applicable					
NCCFN	National Coordinating Committee on Food and Nutrition					
NCHS	National Centre for Health Statistics					
NHLBI	National Heart, Lung and Blood Institute					
OKU	Orang kurang upaya					
PDK	Pusat pemulihan dalam komuniti					
PWD	Persons with disabilities					
RNI	Recommended Nutrient Intake					
ROI	Return-on-investment					
SD	Standard deviation					
SPFII	Secretariat of the Permanent Forum on Indigenous Issues					
Т	Teacher					
Т0	Baseline					
T1	At 1-month interval after intervention					
T2	At 6-month interval after intervention					

UN United Nations

- WHO World Health Organization
- WHO/DAR WHO Disability and Rehabilitation Team

PEMBINAAN, PELAKSANAAN DAN PENILAIAN MODUL PEMAKANAN UNTUK JURULATIH ORANG KURANG UPAYA DI PUSAT PEMULIHAN DALAM KOMUNITI, KELANTAN

ABSTRAK

Orang kurang upaya (OKU) mudah terdedah kepada malpemakanan. Akan tetapi, maklumat tentang status pemakanan dan strategi pemakanan dalam kalangan OKU tempatan agak terhad. Kajian ini dijalankan dari Jun 2010 hingga Julai 2012. Ia bertujuan untuk menilai taraf pemakanan OKU serta membina dan menilai keberkesanan Modul Pemakanan untuk Jurulatih Orang Kurang Upaya di pusat pemulihan dalam komuniti (PDK) Kelantan. Kajian ini meliputi tiga fasa. Penilaian taraf pemakanan OKU di PDK Kelantan telah dilakukan dalam fasa pertama. Seramai 467 orang OKU (281 orang kanak-kanak dan remaja; 186 orang dewasa) telah diambil dan maklumat pemakanan serta antropometri telah dikumpulkan. Modul pemakanan dan Soal Selidik Pengetahuan, Sikap dan Amalan Petugas PDK terhadap Makanan dan Pemakanan OKU (KAP-nOKU) dibina dan disahkan dalam fasa kedua. Satu kajian intervensi telah dijalankan dalam fasa ketiga. Seramai 45 orang guru PDK dari Kelantan telah diambil sebagai kumpulan intervensi dan 42 orang guru PDK dari Terengganu sebagai kumpulan perbandingan. Kumpulan intervensi menerima latihan modul pemakanan manakala kumpulan perbandingan dididik tentang isu penjagaan kesihatan umum OKU. Penilaian awal (T0) dan penilaian susulan KAP-nOKU pada sebulan (T1) dan enam bulan (T2) selepas intervensi telah dilaksanakan dalam kedua-dua kumpulan. Temu bual separa berstruktur kualitatif telah dilakukan dalam kalangan 13 orang guru dari kumpulan

intervensi untuk mengkaji faktor-faktor yang mendorong dan menghalang guru dalam pengurusan pemakanan OKU di PDK. Hasil kajian menunjukkan bahawa prevalens kekurangan berat badan dan berlebihan berat badan dalam kalangan OKU adalah sebanyak 20.3% (17.8% kanak-kanak dan remaja; 24.2% dewasa) dan 22.8% (15.2% kanak-kanak dan remaja; 33.9% dewasa). Dalam fasa kedua, KAP-nOKU yang telah disahkan mengandungi 57 item dan domain-domain pengetahuan (KR20=0.63), sikap (α =0.67) dan amalan (α =0.82) didapati mempunyai konsistensi dalaman yang baik. Analisis ANCOVA berulangan langkah dalam fasa ketiga menunjukkan bahawa kumpulan intervensi telah mencapai skor pengetahuan (p < 0.001) dan amalan (p = 0.001) pemakanan yang lebih tinggi daripada kumpulan perbandingan pada T1 (perbezaan min pengetahuan= 6.95; perbezaan min amalan= 2.93) dan T2 (perbezaan min pengetahuan= 10.30; perbezaan min amalan= 4.78). Skor sikap (p=0.004) pemakanan yang lebih tinggi bagi kumpulan intervensi diperhatikan pada T2 (perbezaan min= 3.08). Hasil kualitatif menunjukkan bahawa sifat peribadi guru yang positif merupakan faktor penting yang mendorong amalan pengurusan pemakanan OKU di PDK. Manakala, sikap rintangan OKU terhadap strategi pemakanan yang positif serta kekurangan sokongan sosial daripada ibu bapa merupakan halangan yang utama. Kesimpulannya, modul pemakanan ini adalah berkesan dalam meningkatkan pengetahuan, sikap dan amalan guru dalam pengurusan pemakanan OKU di PDK Kelantan.

DEVELOPMENT, IMPLEMENTATION AND EVALUATION OF NUTRITION MODULE FOR TRAINERS OF PERSONS WITH DISABILITIES IN COMMUNITY-BASED REHABILITATION CENTRES, KELANTAN

ABSTRACT

Persons with disabilities (PWD) are susceptible to malnutrition. However, little is known about the nutritional status and nutrition strategies among the local disability population. This study was conducted from June 2010 to July 2012. It aimed to assess the nutritional status of PWD, develop and evaluate the effectiveness of Nutrition Module for Trainers of Persons with Disabilities in community-based rehabilitation (CBR) centres, Kelantan. This study included three phases. At Phase 1, nutrition assessment of PWD in CBR centres, Kelantan was conducted. A total of 467 PWD (281 children and adolescents; 186 adults) were recruited and their feeding and anthropometric data were collected using a structured questionnaire. The nutrition module and Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU) were developed and validated in Phase 2. At Phase 3, an intervention study was conducted. Forty-five CBR teachers from Kelantan were recruited as the intervention group and 42 CBR teachers from Terengganu as the comparison group. The intervention group received the nutrition module training while the comparison group was trained on general health care issues about PWD. Baseline (T0) and follow up KAP-nOKU assessments at 1-month (T1) and 6-month (T2) intervals after intervention were administered for both groups. Qualitative semi-structured interviews were conducted among 13 teachers

from intervention group to explore the perceived motivations and barriers to their nutrition management for PWD at CBR centres. Results showed that there was a prevalence of 20.3% of PWD (17.8% children and adolescents; 24.2% adults) who were underweight while 22.8% (15.2% children and adolescents; 33.9% adults) were overweight or obese. At Phase 2, the validated KAP-nOKU contained 57 items with good-to-excellent internal consistencies for knowledge (KR20=0.63), attitude $(\alpha=0.67)$ and practice $(\alpha=0.82)$ domains. At Phase 3, repeated measures ANCOVA demonstrated that intervention group attained significantly higher nutrition knowledge (p < 0.001) and practice (p = 0.001) scores than comparison group at T1 (mean difference for knowledge= 6.95; mean difference for practice= 2.93) and T2 (mean difference for knowledge= 10.30; mean difference for practice= 4.78). Significant higher nutrition attitude (p=0.004) score for intervention group was observed at T2 (mean difference= 3.08). Teachers' positive personal attributes were identified as the primary motivation while PWD's resistance to teachers' positive nutrition management and lack of parental social support were the major barriers. In conclusion, this nutrition module is effective on improving the nutrition knowledge, attitude and practice of CBR teachers in Kelantan.

CHAPTER 1

INTRODUCTION

1.1 Background of the study

Persons with disabilities (PWD), commonly known as *orang kurang upaya* (OKU) in Malaysian context is a complex term referring those who experience long term physical, mental, intellectual and (or) sensory impairments which result in barriers that prevent them from full and effective participation in the society (Law of Malaysia, 2008). PWD, being the most vulnerable group, are susceptible to malnutrition especially those living in rural, low-income and less-resourced areas.

International studies have been reported that PWD, especially the children and adolescents with neurological disorders, are predisposed to undernutrition and growth failure as compared to their healthy peers (Yousafzai *et al.*, 2003; Tomoum *et al.*, 2010). Studies by Marchand *et al.* (2006) in the United States and Sánchez-Lastres *et al.* (2003) in Spain had described almost 50% of children with cerebral palsy were underweight while Bertoli *et al.* (2006) reported 40% of the children with refractory epilepsy in his study at Italy were undernourished and further 24% were wasted.

On the other hand, overweight and obesity among PWD remains another issue of concern as this problem also records a remarkable higher prevalence than the general population (Marshall *et al.*, 2003; Hove, 2004). Overweight problems are

common in both children and adult population with disabilities. Alarming prevalence of around 40% of overweight or obese children and adolescents with disabilities were reported in the United States (33.5%) (Rimmer *et al.*, 2011) and Taiwan (37.7%) (Lin *et al.*, 2010). Among the adult population, data from the United Kingdom suggested that overweight problem affects almost half of the adults with intellectual disabilities (48.7%) being studied (Bhaumik *et al.*, 2008). Similar condition (42.1%) was noted in an Asian study among the institutionalised adults with intellectual disabilities (Hsu *et al.*, 2012).

Malnutrition, either undernutrition or overnutrition, can potentially exacerbate the risk of co-morbidities and further impair the residual functional ability. All these collectively impose greater suffering and burden to the disabled as well as the family, both emotionally and financially (Neyestani *et al.*, 2010).

PWD greatly rely on caregivers for nutrition intake and activities of daily living especially for those who are severely disabled. Several studies have demonstrated that increased caregiver's nutrition knowledge resulted in positive impacts on the nutritional status of children (Smith and Haddard, 2000). Besides, there is opinion suggests that school, child-care or other learning setting provides a link on promoting good nutrition practices between home and school (Hammerschmidt *et al.*, 2011) as the children routinely eat at least one meal in school every day (Freedman and Alvarez, 2010). Teachers, child-care providers or trainers in such settings, therefore, play a crucial role in defining the students' eating environment which will eventually influence the students' nutrition intake, food preferences and also food choices (Hendy and Raudenbush, 2000; Nicklas *et al.*,

2001). Hence, improving nutrition knowledge among primary care providers may show positive effect on reducing the prevalence of malnutrition in PWD.

1.2 Statement of the problem

Malnutrition has become an epidemic in recent years. Numerous cross-sectional and intervention studies globally as well as at national level have been targeted to combat the problem of malnutrition among the general population. However, little is known about the nutritional status and nutrition strategies among those with disabilities, the most vulnerable group of the population. Baseline data on the nutritional status of population with disabilities is imperative to increase the society awareness and concern. Highlighting the nutritional status of PWD can further aid in eliminating the health disparities among this subgroup.

Population with disabilities require a wide range of rehabilitation effort that currently remains challenging to furnish effectively in developing countries (Yousafzai *et al.*, 2003). PWD living in low-income, low-literacy and less-resourced rural areas are even having less opportunity to access to those facilities. Kelantan is recorded as the poorest state and has the lowest urbanization rate in Malaysia (Department of Statistics, 2011). The disadvantaged socioeconomic condition in Kelantan further predisposes the PWD to higher risk of malnutrition. To date, data on the prevalence of malnutrition among PWD in Malaysia is still limited.

In Malaysia, in order to minimize the marginalisation of this population, PWD who are too severely disabled and not accepted by the special education

mainstream will be placed at the community-based rehabilitation (CBR) centres which are under the authority of the Department of Social Welfare. Trainers, or normally known as "teachers', are the persons who teach, train and prepare meals for the PWD in the centres. To be recruited as a CBR teacher, no certification on special education endorsement, early intervention for children with disabilities, elementary or secondary education is required. According to the information obtained from the department and CBR teachers, though periodical training workshops are offered to them, there is still lack of proper and consistent nutrition training courses available to them. Due to insufficient exposure to nutrition information, CBR teachers experienced difficulties in preparing healthy and well balanced diet for the PWD. Additionally, they also faced challenges in managing the PWD with specific feeding difficulties and nutrition problems. Sharing nutrition knowledge with parents as the mutual effort to reinforce the healthful eating behaviour among the PWD is not possible. Subsequently, teaching nutrition message towards PWD in order to influence their food preferences and choices is viewed as a hard chore due to the lack of knowledge and skill.

Besides, there is no current research-tested training material, a nutrition module or guidelines specifically addressing the nutrition needs of PWD, to be used in educating the CBR teachers in Malaysia. The nutrition condition and food practice among PWD in Malaysia may differ from other countries. Moreover, there are also substantial differences in the educational, socioeconomic and cultural background of the local CBR teachers as compared to the trainers in other countries. Hence, adoption of related nutrition module from foreign countries to educate the CBR

teachers in Malaysia is not feasible and ethical. All these emphasize the rationale of having a socio-cultural sensitive and appropriate nutrition module.

1.3 Significance of the study

This study can add considerably to the body of literature pertinent to the prevalence of malnutrition and the common feeding and nutrition problems among PWD in Malaysia. This fundamental data may indicate the future research directions as it can stimulate multiple research ideas especially in the aspects of epidemiology and interventional studies towards this group. Besides, these findings can also raise the stakeholders' attention and provide important insights into the need of policy making in order to provide equal health benefits for those with disabilities.

The development of an evidence-based and research-tested nutrition module can facilitate the professionals or other health care practitioners to conduct consistent and effective nutrition education to all teachers in CBR centres in Malaysia. This module, on the other hand, can serve as the reference material in every CBR centre in Malaysia to guide the care providers in positive nutrition practices. The module might also be valued and adopted by other developing countries for the same purpose based on their condition and pertinence. This study holds relevance for the public, authorities, researchers, health care practitioners, care service providers as well as family members in garnering shared effort for the well-being of the PWD.

1.4 Conceptual framework

Figure 1.1 illustrates the conceptual framework for the present study. This framework is grounded on Bandura's Social Cognitive Theory (Bandura, 1989) for understanding the health behavioural change. The Social Cognitive Theory sets the foundation for designing, developing, implementing and evaluating a program. It incorporates three main factors: personal, environmental and behavioural for evaluating the behavioural change (round dotted triangle in Figure 1.1). In current study, this theory is appropriately relevant to the evaluation of a training using Nutrition Module for Trainers of Persons with Disabilities on improving the knowledge, attitude and practice of teachers in nutrition management for the PWD in CBR centres. This conceptual framework hypothesized that, following the nutrition module training, the teacher's personal factors such as increased nutrition knowledge and positive attitude along with the positive reinforcements from environmental factors, for instance the social support from the colleagues and the availability of facility, will motivate the teachers as well as increase their belief and self-confidence to successfully perform good nutrition practices for the PWD in the CBR centres.

1.5 Methodological framework

The methodological framework for the study is summarized in Figure 1.2. This study was conducted in three phases. Phase 1 was a descriptive cross-sectional study to assess the nutritional status of PWD in CBR centres, Kelantan. Phase 2 comprised of two components. First component was the development of Nutrition Module for Trainers of Persons with Disabilities based on the information obtained from Phase 1



Note: PWD-persons with disabilities; CBR-community-based rehabilitation

Figure 1.1 Conceptual framework of the study



Note: PWD-persons with disabilities; CBR-community-based rehabilitation; KAP-nOKU-Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities

Figure 1.2 Methodological framework of the study

The second component was the development and validation study of the Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU). The final phase, Phase 3, was an intervention study using the nutrition module among teachers from CBR centres, Kelantan with a comparison group of teachers from CBR centres, Terengganu.

1.6 Research questions

This study addresses several research questions as follows.

- 1. What is the nutritional status of PWD in CBR centres, Kelantan?
- 2. Is Nutrition Module for Trainers of Persons with Disabilities a valid training material in nutrition intervention for CBR teachers?
- 3. Is Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU) valid and reliable to assess the nutrition knowledge, attitude and practice of CBR teachers towards nutrition management for PWD?
- 4. Are the CBR teachers in Kelantan satisfied with the nutrition module training (intervention)?
- 5. Is the nutrition module intervention effective on improving the knowledge, attitude and practice of teachers in nutrition management for PWD in CBR centres, Kelantan?
- 6. What are the teachers' perceived motivations and barriers in implementing nutrition management for PWD in CBR centres, Kelantan?

1.7 Objectives of the study

1.7.1 General objective

The general objective of this study is to assess the nutritional status of PWD, develop and evaluate the effectiveness of Nutrition Module for Trainers of Persons with Disabilities in CBR centres, Kelantan.

1.7.2 Specific objectives

The specific objective for Phase 1 is:

1. To assess the nutritional status of PWD in CBR centres, Kelantan.

The specific objectives for Phase 2 are:

- To develop and validate the Nutrition Module for Trainers of Persons with Disabilities.
- 2. To develop and validate the Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU).

The specific objectives for Phase 3 are:

- To evaluate the acceptance of CBR teachers in Kelantan towards the nutrition module training.
- To evaluate the effectiveness of the nutrition module on improving the knowledge, attitude and practice of teachers in nutrition management for PWD in CBR centres, Kelantan.

3. To identify the teachers' perceived motivations and barriers in implementing nutrition management in CBR centres, Kelantan.

1.8 Hypotheses

There are four research hypotheses that are relevant to Phase 2 and 3.

The alternative hypotheses for Phase 2 are:

- H1.The Nutrition Module for Trainers of Persons with Disabilities is a valid training material in nutrition intervention for CBR teachers.
- H2.The Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU) is valid and reliable to assess the nutrition knowledge, attitude and practice of CBR teachers towards nutrition management for PWD.

The alternative hypotheses for Phase 3 are:

- H3.CBR teachers in Kelantan are satisfied with the Nutrition Module for Trainers of Persons with Disabilities training.
- H4.The Nutrition Module for Trainers of Persons with Disabilities is effective on improving the knowledge, attitude and practice of teachers in nutrition management for PWD in CBR centres, Kelantan.

Persons with disabilities (PWD): Persons who have long term limitation in physical, mental, intellectual and (or) sensory ability that can inhibit them from full and effective interaction with the society (Law of Malaysia, 2008).

Community-based rehabilitation (CBR) centre: Training centre which is managed by the local community that provides skills training and therapy services using local resources for PWD who are severely disabled.

Community-based rehabilitation (CBR) teacher: Trainer who is in charge of teaching, training, caring and preparing meals for PWD in CBR centres.

Malnutrition: A pathological state resulting from relatively or completely deficient or excess in one or more essential nutrients (Park, 2000). In this study, malnutrition refers to undernutrition and overnutrition, which are indicated by underweight and overweight or obesity, respectively.

Nutrition Knowledge, Attitude and Practice Questionnaire about Persons with Disabilities (KAP-nOKU): A nutrition questionnaire which acts as an instrument to assess the CBR teacher's nutrition knowledge, attitude and practice towards nutrition management for PWD.

Special education: Formal education modified for physically or mentally handicapped students whose needs cannot be met by the standard school curriculum.

CHAPTER 2

LITERATURE REVIEW

2.1 Disability- size of the problem

Population with disabilities accounts for a considerable portion of the world population. The latest World Report on Disability (WHO, 2011) documented that more than one billion people in the world, which represented 15% of the global population, were living with some forms of disability; and approximately 200 million of them (one in five) encountered substantial difficulties in functioning. The prevalence of disability is on the rise. The figure has been increased as much as around 60% in 10 years time (650 million were reported in 2003). PWD are known to be the world's largest minority, of whom 80% of them are living in developing countries (UN, 2012).

In Malaysia, an estimated rate of 2.8 million people lived with various forms of disability; yet, until August 2010, there were only 313,685 PWD who have registered with the Department of Social Welfare (Zulkiple, 2011). This national registry fails to present the actual picture of PWD in the country as there might be even a larger number go undetected. Among the children population, it was suggested that more than 850 thousand children below 15 years (which constituted 10% of the children population in Malaysia) had a disability with at least one third of them were severe (Amar-Singh, 2008). This data is consistent with Juanita (2011)

who stated that a total of 900 thousand children in Malaysia were diagnosed with various disabilities in 2011.

In spite of the robust healthcare strategies and disability rights movement, PWD are still prevalent to poor health. Issues on availability, accessibility, and sustainability of culturally-sensitive rehabilitation services remain questionable (WHO, 2011).

2.2 Community-based rehabilitation

Community-based rehabilitation (CBR), established by the WHO in the 1980s, is a multi-sectoral strategy that empowers the PWD in developing countries and urban poor areas to gain access to all aspects of rehabilitation services (WHO/DAR & AIFO, 2002). The idea of CBR is that services needed by the PWD should be available to them at affordable cost as well as offered in such a way that suit their regular ways of living, whether in a village or a city (Miles, 2001). Hence, CBR has been defined as a strategy within the general community development for rehabilitation, equalisation of opportunities, reduction of poverty and social inclusion of all children and adults with disabilities (WHO, 2004). It is implemented through the joined efforts of PWD themselves, their families, local communities, relevant governmental and non-governmental health, social, education, vocational and other services (WHO, 2004).

In Malaysia, CBR program is jointly managed by the Department of Social Welfare and Ministry of Health (MOH) since 1984 and is primarily focused at rural regions. It targets five main objectives, i) to enhance sense of concern and responsibility among local community in rehabilitation effort towards PWD; ii) to gather local resources for the development of rehabilitation services; iii) to encourage the use of appropriate, reasonable and effective methods or techniques which are sensitive to the need of local condition; iv) to utilise the available local infrastructures for developing the rehabilitation services; and v) to expand and decentralize comprehensive service delivery based on the needs of PWD (JKM, 2011).

The CBR centres in Malaysia, or known as *pusat pemulihan dalam komuniti* (PDK), provide rehabilitation training for PWD who are too young, too severely disabled and not accepted by the special education mainstream. No age limitation or specific requirement for PWD enrolment is indicated. A total of 428 CBR centres were established in Malaysia by 2010 (JKM, 2011). Table 2.1 lists the number of CBR centres, trainers and PWD registered in each state of Malaysia. Based on the registration criteria of the Department of Social Welfare, PWD are categorized into one of the seven disability categories upon registration (JKM, 2012). The disability categories are hearing, visual, speech, physical, learning, mental and multiple disabilities. Table 2.2 details the explanation for each category.

The CBR centres function as registration, referral, rehabilitation, training and resource centres (Isa *et al.*, 2013). Training services offered including gross motor and fine motor skills, social and language development, activities of daily living, writing, reading and counting, vocational, sports and recreation (JKM, 2011).

State	Number			
-	CBR centres	Supervisors	Trainers	PWD
Perlis	5	5	17	242
Kedah	27	27	126	1471
Pulau Pinang	20	20	69	647
Perak	36	36	114	1286
Selangor	39	39	162	2257
WP Kuala Lumpur	9	9	45	380
Negeri Sembilan	40	40	231	1639
Melaka	17	17	77	715
Johor	63	63	271	2853
Pahang	45	45	131	1424
Kelantan	33	33	103	1218
Terengganu	40	40	123	1354
Sarawak	25	25	150	1932
Sabah	27	27	88	1465
WP Labuan	2	2	7	60
Total	428	428	1714	18943

 Table 2.1
 Number of CBR centres, trainers and PWD registered in Malaysia

Source: JKM (2011)

In Kelantan, there are 33 CBR centres and each centre is operated by three to four trainers. The PWD (*Pelatih OKU*) attend classes from 8am to 1pm, from Sunday to Wednesday. On Thursdays, the CBR trainers pay home visits to those home-based PWD, mostly the severely disabled children. Each PWD receives a monthly allowance of RM150 from the Department of Social Welfare. PWD take one lunch meal which is prepared by the trainers in the centres each day. Food budget is obtained from the deduction on PWD's allowance and the amount deducted is based on the consensus of the authority and parents of PWD in each centre. In general, an amount of RM20 to RM30 is collected from each PWD for food expenses.

The CBR trainers (*Petugas PDK*), or commonly known as CBR teachers (*Cikgu PDK*), are recruited on voluntary basis to provide service at the centres with a monthly salary of RM800. They are usually belonged to the local community where the centre locates. Recruitment is neither based on their educational level nor age.

No.	Category	Explanation
1	Hearing	Inability to hear clearly in both ears with or without the use of
	•	hearing aids.
		Hearing disability can be divided into four levels:
		 Minimum- 15-<30dB (children) and 20-<30dB (adult)
		• Intermediate- 30-<60dB
		• Severe- 60-< 90dB
		• Profound- \geq 90dB
2	Visual	Poor vision in both eyes; blindness in one or both eyes.
		Poor vision is defined as visual acuity of less than 6/18 but equal to
		or better than 3/60 or corresponding visual field loss to less than
		20°.
		Blindness is referred to visual acuity of less than 3/60 or
		corresponding visual field loss to less than 10°.
3	Speech	Inability to produce sound correctly or fluently; and the speech
		cannot be understood by others during communication. Diagnosis
	Dhavelagel	for children to be assessed only after five years old.
4	Physical	Permanent physical disability, either caused by loss of limbs or any
		limba For instance
		Limb defects (concentral or acquired)
		 Enino defects (congenital of acquired) Spinol cord injunt
		• Spinar cord injury
		 Shoke Troumatic brain injury
		• A chondronlasia
		Cerebral palay
	Learning	• Celebral parsy Disorder in learning, cognition and intelligence that is inconsistent
5	Leanning	with the biological age. For example
		 Global developmental delayed (less than 5 years old)
		 Down's syndrome
		 Intellectual disability (more than 5 years old)
		 Autism spectrum disorders
		 Attention deficit hyperactive disorder
		 Specific (dyslexia, dyscalculia and dysgraphia)
6	Mental	Severe mental illness or disorder that impairs one's ability to
0	Wiental	function effectively in own living or interaction with the society
		For instance.
		• Organic mental disorder
		• Schizophrenic, paranoid and other psychotic disorder
		 Mood disorder (depression and bipolar)
7	Multiple	Having a combination of two or more disabilities and generally
		inappropriate to be classified into category 1 to 6
	·····	

Source: JKM (2012)

In addition, no qualification on special education or early childhood program is required (Chen et al., 2012) and most of the teachers only attain their highest educational level at lower (Form 3) or upper secondary school (Form 5). The teachers are exposed to the training courses on skills in handling PWD only after their recruitment. These training courses are usually organized by the Ministry of Health, higher education institutions and non-governmental organizations. Most of the trainings offered are focusing on physiotherapy or occupational therapy, sexual education and vocational training for the PWD. In the centres, the teachers play a crucial role in training, teaching and preparing meals for the PWD as well as supporting the caregivers in taking care of their children (Chen et al., 2012; Isa et al., 2013). The PWD and their families highly depend on the CBR teachers for rehabilitation services and information to promote good health, especially those in rural and low-literacy areas. According to the information given, food preparation or menu standardization for all CBR centres in Kelantan was not indicated as there was no specific guideline provided by the authorities. Food prepared was usually based on the PWD's preferences, teachers' convenience and availability of food sources.

Various health strategies have been provided by the Malaysia Ministry of Health to ensure a comprehensive health care for PWD to achieve optimal health and maximum independence. Since 1986, Early Intervention Programme for Children with Special Needs was initiated and mainly focused on early detection of disability population at health clinics and referral to hospitals for rehabilitation. Beginning 1996, rehabilitation services are started at selected health clinics and managed by trained public health nurses. The nurses undergo various in-service trainings which mostly emphasize on basic techniques for handling PWD with gross motor and fine motor delay; visual impairment; activities of daily living, communication, personal social and behavioural, sexual and reproductive health problems. The Health Care for PWD Plan of Action 2011-2010 was also developed in line with the Convention on the Rights of PWD and the PWD Act 2008 to outline the health service plans for PWD. In 2011, together with *Badan Amal dan Kebajikan Tenaga Isteri-Isteri* (BAKTI), MOH developed and launched a manual on healthy lifestyle, "*PDK KU SIHAT: Garis Panduan & Manual Aktiviti*", which focuses on healthy eating and regular exercise for PWD. This manual was only distributed among CBR centres in Kelantan at August 2012. In this program, health personnel such as public health nurses, nutritionists and occupational therapists are in charge of providing outreach services such as advice on healthy menu, healthy meal preparation and safe physical activity to CBR centres within operational area of the health clinics. However, issues of accessibility and sustainability of the above strategies remain uncertain.

2.3 Disabilities and nutritional concerns

A disability can occur at prenatal, perinatal, neonatal or at any stage of an individual's life. These disabilities may be brought about by diseases harming mother during pregnancy; genetic abnormalities; oxygen deprivation and acquired brain injury during birth; accidents, infections or diseases acquired after birth. Disabilities involving brain and central nervous system are permanent, irreversible and affecting a wide range of developmental, motor and muscular, sensory, learning and cognitive abilities (Almond *et al.*, 2007).

PWD frequently have nutritional concerns (Van Riper, 2010) resulted from their disablement that may lead them to poor health. Besides, they are also prone to develop co-morbid conditions such as obesity, diabetes mellitus or cardiovascular diseases that will further worsen their functional abilities (Traci *et al.*, 2002). The following section discusses the types of disability most prevalent in the CBR centres in Malaysia and their related nutritional concerns.

2.3.1 Down's syndrome

Down's syndrome is a genetic disorder caused by the presence of an extra chromosome 21 or known as Trisomy 21. Down's syndrome occurs in approximately 1 per 660 live births in Malaysia (Zulkiple, 2012). Children with Down's syndrome are affected with cognitive deficit and growth impairment (Nehring, 2010). They are also at risk for multiple congenital anomalies, such as congenital heart defects, respiratory infections, gastrointestinal problems (e.g. constipation and celiac disease) and endocrine abnormalities (Roizen and Patterson, 2003; Almond *et al.*, 2007; Van Riper, 2010).

Feeding difficulties are common among children with Down's syndrome (Lewis and Kritzinger, 2004; Almond *et al.*, 2007). These feeding problems are associated with decreased muscle tones- hypotonia, which affects the strength and mobility of oral muscles (Kumin *et al.*, 1991) and results in weak lip closure, poor sucking, chewing and swallowing (Almond *et al.*, 2007). Additionally, the genetic anomaly causes multiple cranial skeletal abnormalities (Lewis and Kritzinger, 2004) which in turn contribute to their feeding difficulties. Small oral cavity, large tongue,

short and narrow palate, enlarged tonsils along with decreased nasal passages may interfere the oral feeding in children with Down's syndrome (Lewis and Kritzinger, 2004). Congenital heart defects occur in 40 to 50% of children, are important risk factors for the feeding problems (Pueschel, 1990; Lewis and Kritzinger, 2004; Freeman *et al.*, 2008). Children with heart defects may experience breathing difficulty, vomiting and easily fatigue during eating (Lewis and Kritzinger, 2004). All these combined conditions can lead to inadequate food intake and poor weight gain in children with Down's syndrome.

On the other hand, there is also a high tendency for persons with Down's syndrome to become overweight or obese (Cronk *et al.*, 1988; Rubin *et al.*, 1998; Murray and Ryan-Krause, 2010). Some studies suggest that a rate of 30 to 50% of these children are obese (Harris *et al.*, 2003). Endocrine dysfunctions such as hypothyroidism which leads to decreased basal metabolic rate (Luke *et al.*, 1994; Bauer *et al.*, 2003) and abnormal leptin level that impairs regulation of satiety (Cento *et al.*, 1999; Magni *et al.*, 2004), serve as the main culprits. Furthermore, delayed gross motor abilities (Holcomb *et al.*, 2009) and negative behavioural tendency, for instance the inattention, non-compliance and oppositional behaviours (Jahromi *et al.*, 2008) among persons with Down's syndrome cause them to live in a sedentary lifestyle and have less engagement in physical activity. These collectively increase the prevalence of overweight and obesity.

2.3.2 Cerebral palsy

According to Blair *et al.* (2007), cerebral palsy (CP) refers to a group of permanent disorders involving movement and posture which are resulted from non-progressive interference or lesion in the developing brain. CP is estimated to occur in 2 to 3 per 1000 live born infants (Yeargin-Allsopp *et al.*, 2008; Benfer *et al.*, 2012). A host of disabilities co-occur with CP. These include intellectual impairment, seizure, sensory (visual, hearing, touch) and communication disorders (Odding *et al.*, 2006; Rosenbaum *et al.*, 2007; Van Riper, 2010).

The neurologic lesion greatly impairs the oral motor functions and manifests a range of complex feeding problems (Andrew et al., 2011; Benfer et al., 2012). Reilly and Skuse (1992) explained that feeding dysfunction in persons with CP is always due to three interacting conditions: i) the impaired oral motor functions (hypotonia, persistent tongue thrust, weak lip closure); ii) diminished neurologic maturation (hyperactive gag reflexes and reduced swallowing coordination); and iii) poor truncal support and head control result in inappropriate seating posture that make mealtime an unpleasant experience. They frequently suffer from dysphagia or swallowing difficulties, coughing, choking, vomiting or sensory impairment to clearly communicate pain, hunger or satiety (Tomoum et al., 2010; Van Riper, 2010; Andrew et al., 2011). In addition, their reduced feeding skills can be further aggravated by specific medical conditions such as gastro-esophageal reflux disease (GERD), aspiration, constipation and seizure (Almond et al., 2007; Tomoum et al., 2010; Andrew et al., 2011). Furthermore, drooling which is defined as the unintentional loss of saliva (Reddihough et al., 2010), is another common problem in children with CP. Dehydration, impaired mastication function, oral mouth infections (Van der Burg *et al.*, 2006) and aspiration pneumonia (Walshe *et al.*, 2010) are the nutritional concerns inherent to the drooling problem.

The aforementioned oral-motor involvement causes lengthy mealtimes, increases fatigue and mealtime resistance and further contributes to insufficient energy intake and eventually growth failure (Tomoum *et al.*, 2010; Van Riper, 2010; Andrew *et al.*, 2011). In a multicentre study involving 230 children and adolescents with CP, Fung *et al.* (2002) had demonstrated that the extent of feeding difficulties is directly associated with the degree of undernutrition. Gisel *et al.* (2000) and Fung *et al.* (2002) further added even children who experienced mild feeding dysfunction had inadequate food intake and poor growth.

2.3.3 Autism spectrum disorders

Autism spectrum disorders (ASD) is defined as a set of lifelong neurodevelopment disabilities characterized by social and communication deficits and presence of stereotyped and repetitive behaviours (Matson, 2007; Soden *et al.*, 2012; Zimmer *et al.*, 2012). In Malaysia, ASD has a prevalence of 1 in 625 children (Dolah *et al.*, 2011). Common co-morbidities identified among persons with ASD are learning disabilities (but surprisingly, some may have average or above Intelligence Quotient level), attention deficit hypersensitivity disorder (ADHD), altered gastrointestinal functions, fear or anxiety, and epilepsy (Connor, 2007; Twachtman-Reilly *et al.*, 2008; Parellada *et al.*, 2011).

Based on an early report by DeMyer (1979), more than 90% of children with ASD have some feeding and mealtime problems. These problems have been documented by several investigators over the years and are classified into three main categories: i) food selectivity or "picky eating"; ii) food refusal; and iii) disruptive mealtime behaviours (Ahearn et al., 2001; Schreck and Williams, 2006; Bandini et al., 2010). Some children with ASD are overly selective and only eat a restricted range of food (Bandini et al., 2010). They can have strong aversions to specific colours, smells, textures or temperature. This rigidity can extend to food brands or packaging, food presentation and mealtime environments (Williams et al., 2000; Schreck et al., 2004; Schreck and Williams, 2006; Connor, 2007; Bandini et al., 2010). Hence, introduction of new food may be challenging and distressing. Field et al. (2003) further added that some may even refuse to eat the entire food group, for instance the milk and dairy products (Zimmer et al., 2012). These negative food behaviours are detrimental to their nutritional status. Nevertheless, numerous studies demonstrated conflicting findings about the nutrition deficiencies of this population (Zimmer et al., 2012). Macronutrients intake of children with ASD is not different from their healthy peers, as reported by Emond et al. (2010) and Johnson et al. (2008). However, micronutrient deficiencies are evident especially for calcium, iron, Vitamin A, Vitamin C and Vitamin D (Lindsay et al., 2006; Herndon et al., 2009; Bandini et al., 2010; Zimmer et al., 2012).

Problematic mealtime behaviours are another matter of concern. Persons with ASD have minimal verbal skills and have difficulties to communicate experience, feelings or symptoms (Buie *et al.*, 2010). Additionally, the sensory processing deficits restrict their ability to identify the source of body discomfort or pain