



Final Examination  
2018/2019 Academic Session

June 2019

**JIM106 – Elementary Statistics  
(Asas Statistik)**

Duration: 3 hours  
(Masa: 3 jam)

---

Please check that this examination paper consists of **TWENTY FOUR (24)** pages of printed material before you begin the examination.

*[Sila pastikan bahawa kertas peperiksaan ini mengandungi **DUA PULUH EMPAT (24)** muka surat yang bercetak sebelum anda memulakan peperiksaan ini].*

**Instructions** : Answer **ALL** questions.

**Arahan** : Jawab **SEMUA** soalan].

In the event of any discrepancies, the English version shall be used.

*[Sekiranya terdapat sebarang percanggahan pada soalan peperiksaan, versi Bahasa Inggeris hendaklah digunapakai].*

1. (a). A study was done to determine the numbers of days that it took to receive a shipment from the day it was ordered. The following data was recorded.

Days	Frequency
1-3	6
4-6	8
7-9	10
10-12	7
13-15	0
16-18	5

- (i). Draw the histogram and the frequency polygon.  
 (ii). Find the mean, median, mode and standard deviation.  
 (iii). Comment on the distribution of the data.

(70 marks)

- (b). Events  $A$ ,  $B$ ,  $C$  are independent with  $P(A) = 1/2$ ,  $P(B) = 1/4$  and  $P(C) = 1/8$ .

Find

- (i).  $P(\bar{A} \cup (B \cap C))$ .  
 (ii).  $P(\overline{A \cup B \cup C})$ .

(30 marks)

2. (a). A car plate number consists of 5 letters followed by 2 digits.

- (i). How many plate numbers can be made if repetitions are allowed?  
 (ii). How many plate numbers can be made if repetitions are NOT allowed?

(30 marks)

- (b). A population is known to be normally distributed with  $\sigma^2 = 25$ . Find the sample size so as to be 95% confident that the difference between the sample mean and the population mean is less than 1 unit.

(30 marks)

...3/-

- (c). The number of calls received per day at a crisis center is distributed as follows:

Number $X$	30	31	32	33	34
Probability $p(x)$	0.05	0.21	$k$	0.25	0.11

Find

- (i).  $k$   
 (ii).  $E(X)$   
 (iii).  $Var(X)$

(40 marks)

3. (a). A recent study of 75 workers found that 53 people rode the bus to work each day. Find the 95% confidence interval of the proportion of all workers who rode the bus to work.

(35 marks)

- (b). A sample of labor costs per hour to assemble a certain product has a mean of \$2.60 and a standard deviation of \$0.15. Using Chebyshev's theorem, find the range in which at least 88.89% of the data will lie.

(30 marks)

- (c). A switchboard handles 600 calls/hr during peak hours. The switched board fuse will blow if more than 20 connections/min are made. Use the Poisson distribution to find the probability that the fuse will blow.

(35 marks)

4. (a). A biologist knows that the average length of a certain leaf is 4 inches with a standard deviation of 0.6 inch. A sample size of 20 leaves after being given a new plant food had an average length of 4.2 inches. Assume that the variable length is normally distributed. Is there reason to believe that the new food is responsible for a change in the growth of the leaves? Use  $\alpha = 0.01$ .

(60 marks)

...4/-

- (b).  $X$  is normally distributed with mean  $\mu$  and variance  $\sigma^2$ . Given  $P(X > 1000) = 0.9772$  and  $P(X > 13000) = 0.0228$ , find  $\mu$  and  $\sigma$ .

(40 marks)

5. (a). A researcher wanted to compare the pulse rates of identical twins to see whether there was any difference. Eight sets of twins were selected. The rates (beats/min) are given below.

Twin A	87	92	78	83	88	90	84	93
Twin B	83	95	79	83	86	93	80	86

At  $\alpha = 0.01$ , is there a significant difference in the average pulse rates of the twins? Use the  $P$ -value method.

(70 marks)

- (b). A box contains 6 blue balls, 3 red balls and 1 yellow ball. Two balls are taken from the box.
- (i). Write the sample space for this experiment.
- (ii). Find the probability of getting both balls of different colour.

(30 marks)

1. (a). Satu kajian dilakukan untuk menentu bilangan hari yang diperlu untuk menerima penghantaran dari hari pesanan dibuat. Data berikut telah direkodkan.

Hari	Frekuensi
1-3	6
4-6	8
7-9	10
10-12	7
13-15	0
16-18	5

- (i). Lakarkan histogram dan poligon frekuensi.  
(ii). Cari min, median, mod dan sisihan piawai.  
(iii). Komen tentang taburan data.

(70 markah)

- (b). Peristiwa  $A$ ,  $B$ ,  $C$  adalah tak bersandar dengan  $P(A)=1/2$ ,  $P(B)=1/4$ , and  $P(C)=1/8$ .

Cari

- (i).  $P(\bar{A} \cup (B \cap C))$ .  
(ii).  $P(\overline{A \cup B \cup C})$ .

(30 markah)

2. (a). Nombor plat kereta terdiri daripada 5 abjad diikuti dengan 2 digit.  
(i). Berapakah nombor plat dapat dibuat jika ulangan dibenarkan?  
(ii). Berapakah nombor plat dapat dibuat jika ulangan TIDAK dibenarkan?  
(30 markah)

- (b). Suatu populasi diketahui bertabur secara normal dengan  $\sigma^2 = 25$ . Cari saiz sampelnya supaya mempunyai keyakinan 95% bahawa perbezaan di antara purata sampel dengan purata sebenarnya adalah kurang dari 1 unit.

(30 markah)

...6/-

- (c). Bilangan panggilan yang diterima setiap hari di pusat krisis bertabur seperti berikut:

Nombor $X$	30	31	32	33	34
Kemungkinan $p(x)$	0.05	0.21	$k$	0.25	0.11

Cari

- (i).  $k$ .  
 (ii).  $E(X)$ .  
 (iii).  $Var(X)$ .

(40 markah)

3. (a). Satu kajian terkini yang terdiri daripada 75 orang pekerja mendapati bahawa 53 pekerja menaiki bas ke tempat kerja setiap hari. Cari selang keyakinan 95% bagi kadaran pekerja yang menaiki bas ke tempat kerja.

(35 markah)

- (b). Satu sampel kos buruh per jam untuk memasang produk tertentu mempunyai min \$2.60 dan sisihan piawai \$0.15. Dengan menggunakan teorem Chebyshev, cari julat di mana sekurang-kurangnya 88.89% data akan berada.

(30 markah)

- (c). Suatu papan suis mengendalikan 600 panggilan/jam pada waktu yang paling sibuk. Fius papan suis akan terbakar jika ia membuat lebih daripada 20 sambungan/minit. Gunakan taburan Poisson untuk mendapatkan kebarangkalian fius ini akan terbakar.

(35 markah)

4. (a). Seorang ahli biologi mengetahui bahawa purata panjang sejenis daun ialah 4 inci dengan sisihan piawai 0.6 inci. Satu sampel bersaiz 20 daun yang telah diberi makanan pokok yang baru mempunyai purata panjang 4.2 inci. Andaikan pembolehubah panjang daun bertabur secara normal. Adakah sebab untuk mempercayai bahawa makanan pokok yang baru itu bertanggungjawab ke atas perubahan dalam pertumbuhan daun tersebut? Guna  $\alpha = 0.01$ .

(60 markah)

...7/-

- (b).  $X$  bertabur secara normal dengan min  $\mu$  dan varians  $\sigma^2$ . Diberi  $P(X > 1000) = 0.9772$  dan  $P(X > 13000) = 0.0228$ , cari  $\mu$  dan  $\sigma$ .

(40 markah)

5. (a). Seorang penyelidik ingin membandingkan kadar denyutan nadi kembar seiras untuk melihat sama ada terdapat perbezaan. Lapan kembar seiras dipilih. Kadar (denyutan/min) diberi seperti berikut:

Kembar A	87	92	78	83	88	90	84	93
Kembar B	83	95	79	83	86	93	80	86

Pada  $\alpha = 0.01$ , adakah terdapat perbezaan signifikan dalam denyutan nadi kembar seiras? Gunakan kaedah nilai- $P$ .

(70 markah)

- (b). Satu kotak mempunyai 6 biji bola biru, 3 biji bola merah dan 1 biji bola kuning. Dua biji bola dikeluarkan daripada kotak ini.

- (i). Tulis ruang sampel bagi eksperimen ini.  
(ii). Cari kebarangkalian kedua-dua biji bola yang dikeluarkan adalah berlainan warna.

(30 markah)

**FORMULAS****Chapter 3 Data Description**

Mean for individual data:  $\bar{X} = \frac{\sum X}{n}$

Mean for grouped data:  $\bar{X} = \frac{\sum f \cdot X_m}{n}$

Standard deviation for a sample:

$$s = \sqrt{\frac{n(\sum X^2) - (\sum X)^2}{n(n-1)}}$$

Standard deviation for grouped data:

$$s = \sqrt{\frac{n(\sum f \cdot X_m^2) - (\sum f \cdot X_m)^2}{n(n-1)}}$$

Range rule of thumb:  $s \approx \frac{\text{range}}{4}$

Median for grouped data:

$$MD = \frac{(n/2) - cf}{f}(w) + L_m$$

where

$n$  = sum of frequencies

$cf$  = cumulative frequency of class immediately preceding the median class

$w$  = width of median class

$f$  = frequency of median class

$L_m$  = lower boundary of median class



**Chapter 4 Probability of Counting Rules**

Addition rule 1 (mutually exclusive events):

$$P(A \text{ or } B) = P(A) + P(B)$$

Addition rule 2 (events not mutually exclusive):

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

Multiplication rule 1 (independent events):

$$P(A \text{ and } B) = P(A) \cdot P(B)$$

Multiplication rule 2 (dependent events):

$$P(A \text{ and } B) = P(A) \cdot P(B|A)$$

Conditional probability:  $P(B|A) = \frac{P(A \text{ and } B)}{P(A)}$

Complementary events:  $P(\bar{E}) = 1 - P(E)$

Fundamental counting rule: Total number of outcomes of a sequence when each event has a different number of possibilities:  $k_1 \cdot k_2 \cdot k_3 \cdots k_n$

Permutation rule: Number of permutations of  $n$  objects taking  $r$  at a time is

$${}_n P_r = \frac{n!}{(n-r)!}$$

Combination rule: Number of combinations of  $r$  objects selected from  $n$  objects is

$${}_n C_r = \frac{n!}{(n-r)!r!}$$

**Chapter 5 Discrete Probability Distributions**

Mean for a probability distribution:  $\mu = \sum [X \cdot P(X)]$

Variance and standard deviation for a probability distribution:

$$\sigma^2 = \sum [X^2 \cdot P(X)] - \mu^2$$

$$\sigma = \sqrt{\sum [X^2 \cdot P(X)] - \mu^2}$$

Expectation:  $E(X) = \sum [X \cdot P(X)]$

Binomial probability:  $P(X) = \frac{n!}{(n-X)!X!} \cdot p^X \cdot q^{n-X}$

Mean for binomial distribution:  $\mu = n \cdot p$

Variance and standard deviation for the binomial distribution:

$$\sigma^2 = n \cdot p \cdot q \qquad \sigma = \sqrt{n \cdot p \cdot q}$$

Multinomial probability:

$$P(X) = \frac{n!}{X_1! X_2! X_3! \dots X_k!} \cdot p_1^{X_1} \cdot p_2^{X_2} \cdot p_3^{X_3} \dots p_k^{X_k}$$

Poisson probability:  $P(X; \lambda) = \frac{e^{-\lambda} \lambda^X}{X!}$  where  $X = 0, 1, 2, \dots$

Hypergeometric probability:  $P(X) = \frac{{}_a C_X \cdot {}_b C_{n-X}}{{}_{a+b} C_n}$

### Chapter 6 The Normal Distribution

Standard score:  $z = \frac{X - \mu}{\sigma}$  or  $\frac{X - \bar{X}}{s}$

Mean of sample means:  $\mu_{\bar{X}} = \mu$

Standard error of the mean:  $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$

Central limit theorem formula:  $z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$

### Chapter 7 Confidence Intervals and Sample Size

z confidence interval for means:

$$\bar{X} - z_{\alpha/2} \left( \frac{\sigma}{\sqrt{n}} \right) < \mu < \bar{X} + z_{\alpha/2} \left( \frac{\sigma}{\sqrt{n}} \right)$$

t confidence interval for means:

$$\bar{X} - t_{\alpha/2} \left( \frac{s}{\sqrt{n}} \right) < \mu < \bar{X} + t_{\alpha/2} \left( \frac{s}{\sqrt{n}} \right)$$

Sample size for means:  $n = \left( \frac{z_{\alpha/2} \cdot \sigma}{E} \right)^2$  where  $E$  is the maximum error of estimate

Confidence interval for a proportion:

$$\hat{p} - (z_{\alpha/2}) \sqrt{\frac{\hat{p}\hat{q}}{n}} < p < \hat{p} + (z_{\alpha/2}) \sqrt{\frac{\hat{p}\hat{q}}{n}}$$

Sample size for a proportion:  $n = \hat{p}\hat{q}\left(\frac{z_{\alpha/2}}{E}\right)^2$

where  $\hat{p} = \frac{X}{n}$  and  $\hat{q} = 1 - \hat{p}$

Confidence interval for variance:

$$\frac{(n-1)s^2}{\chi_{\text{right}}^2} < \sigma^2 < \frac{(n-1)s^2}{\chi_{\text{left}}^2}$$

Confidence interval for standard deviation:

$$\sqrt{\frac{(n-1)s^2}{\chi_{\text{right}}^2}} < \sigma < \sqrt{\frac{(n-1)s^2}{\chi_{\text{left}}^2}}$$

## Chapter 8 Hypothesis Testing

$z$  test:  $z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$  for any value  $n$ . If  $n < 30$ , population must be normally distributed.

$$z = \frac{\bar{X} - \mu}{s/\sqrt{n}} \text{ for } \sigma \text{ unknown and } n \geq 30$$

$t$  test:  $t = \frac{\bar{X} - \mu}{s/\sqrt{n}}$  for  $n < 30$  (d.f. =  $n - 1$ )

$z$  test for proportions:  $z = \frac{\hat{p} - p}{\sqrt{pq/n}}$

Chi-square test for a single variance:  $\chi^2 = \frac{(n-1)s^2}{\sigma^2}$

(d.f. =  $n - 1$ )

### Chapter 9 Testing the Difference Between Two Means, Two Variances and Two Proportions

z test for comparing two means (independent samples);

$$z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}$$

Formula for the confidence interval for difference of two means (large samples):

$$(\bar{X}_1 - \bar{X}_2) - z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} < \mu_1 - \mu_2 < (\bar{X}_1 - \bar{X}_2) + z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

*Note:*  $s_1^2$  and  $s_2^2$  can be used when  $n_1 \geq 30$  and  $n_2 \geq 30$ .

F test for comparing two variances:  $F = \frac{s_1^2}{s_2^2}$

where  $s_1^2$  is the larger variance and

$$\text{d.f.N.} = n_1 - 1, \text{ d.f.D} = n_2 - 1$$

t test for comparing two means (independent samples, variances not equal):

$$t = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

(d.f. = the smaller on  $n_1 - 1$  or  $n_2 - 1$ )

Formula for the confidence interval for difference of two means (small independent samples, variance unequal):

$$(\bar{X}_1 - \bar{X}_2) - t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} < \mu_1 - \mu_2 < (\bar{X}_1 - \bar{X}_2) + t_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

(d.f. = smaller of  $n_1 - 1$  and  $n_2 - 1$ )

$t$  test for comparing two means (independent samples, variances equal):

$$t = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{(n_1 + n_2 - 2)} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}}$$

(d.f. =  $n_1 + n_2 - 2$ )

Formula for the confidence interval for difference of two means (small independent samples, variances equal):

$$(\bar{X}_1 - \bar{X}_2) - t_{\alpha/2} \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$(\bar{X}_1 - \bar{X}_2) + t_{\alpha/2} \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

and d.f. =  $n_1 + n_2 - 2$ .

$t$  test for comparing two means for dependent samples:

$$t = \frac{\bar{D} - \mu_D}{s_D / \sqrt{n}} \quad \text{where } \bar{D} = \frac{\sum D}{n} \quad \text{and}$$

$$s_D = \sqrt{\frac{n \sum D^2 - (\sum D)^2}{n(n-1)}} \quad (\text{d.f.} = n - 1)$$

Formula for confidence interval for the mean of the difference for dependent samples:

$$\bar{D} - t_{\alpha/2} \frac{S_D}{\sqrt{n}} < \mu_D < \bar{D} + t_{\alpha/2} \frac{S_D}{\sqrt{n}}$$

(d.f. =  $n - 1$ )

$t$  test for comparing two proportions:

$$z = \frac{(\hat{p}_1 - \hat{p}_2) - (p_1 - p_2)}{\sqrt{\bar{p}\bar{q}\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$$

$$\text{where } \bar{p} = \frac{X_1 + X_2}{n_1 + n_2} \quad \hat{p}_1 = \frac{X_1}{n_1}$$

$$\bar{q} = 1 - \bar{p} \quad \hat{p}_2 = \frac{X_2}{n_2}$$

Formula for the confidence interval for the difference of two proportions:

$$(\hat{p}_1 - \hat{p}_2) - z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}} < p_1 - p_2 < (\hat{p}_1 - \hat{p}_2) + z_{\alpha/2} \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}$$

TABLES

**Table E The Standard Normal Distribution**

**Cumulative Standard Normal Distribution**

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

For z values less than -3.49, use 0.0001.






Table E (continued)										
Cumulative Standard Normal Distribution										
z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

For z values greater than 3.49, use 0.9999.




Table C The Poisson Distribution										
x	$\lambda$									
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0
0	.9048	.8187	.7408	.6703	.6065	.5488	.4966	.4493	.4066	.3679
1	.0905	.1637	.2222	.2681	.3033	.3293	.3476	.3595	.3659	.3679
2	.0045	.0164	.0333	.0536	.0758	.0988	.1217	.1438	.1647	.1839
3	.0002	.0011	.0033	.0072	.0126	.0198	.0284	.0383	.0494	.0613
4	.0000	.0001	.0003	.0007	.0016	.0030	.0050	.0077	.0111	.0153
5	.0000	.0000	.0000	.0001	.0002	.0004	.0007	.0012	.0020	.0031
6	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0002	.0003	.0005
7	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001
x	$\lambda$									
	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
0	.3329	.3012	.2725	.2466	.2231	.2019	.1827	.1653	.1496	.1353
1	.3662	.3614	.3543	.3452	.3347	.3230	.3106	.2975	.2842	.2707
2	.2014	.2169	.2303	.2417	.2510	.2584	.2640	.2678	.2700	.2707
3	.0738	.0867	.0998	.1128	.1255	.1378	.1496	.1607	.1710	.1804
4	.0203	.0260	.0324	.0395	.0471	.0551	.0636	.0723	.0812	.0902
5	.0045	.0062	.0084	.0111	.0141	.0176	.0216	.0260	.0309	.0361
6	.0008	.0012	.0018	.0026	.0035	.0047	.0061	.0078	.0098	.0120
7	.0001	.0002	.0003	.0005	.0008	.0011	.0015	.0020	.0027	.0034
8	.0000	.0000	.0001	.0001	.0001	.0002	.0003	.0005	.0006	.0009
9	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0002
x	$\lambda$									
	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
0	.1225	.1108	.1003	.0907	.0821	.0743	.0672	.0608	.0550	.0498
1	.2572	.2438	.2306	.2177	.2052	.1931	.1815	.1703	.1596	.1494
2	.2700	.2681	.2652	.2613	.2565	.2510	.2450	.2384	.2314	.2240
3	.1890	.1966	.2033	.2090	.2138	.2176	.2205	.2225	.2237	.2240
4	.0992	.1082	.1169	.1254	.1336	.1414	.1488	.1557	.1622	.1680
5	.0417	.0476	.0538	.0602	.0668	.0735	.0804	.0872	.0940	.1008
6	.0146	.0174	.0206	.0241	.0278	.0319	.0362	.0407	.0455	.0504
7	.0044	.0055	.0068	.0083	.0099	.0118	.0139	.0163	.0188	.0216
8	.0011	.0015	.0019	.0025	.0031	.0038	.0047	.0057	.0068	.0081
9	.0003	.0004	.0005	.0007	.0009	.0011	.0014	.0018	.0022	.0027
10	.0001	.0001	.0001	.0002	.0002	.0003	.0004	.0005	.0006	.0008
11	.0000	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0002	.0002
12	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001
x	$\lambda$									
	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0
0	.0450	.0408	.0369	.0334	.0302	.0273	.0247	.0224	.0202	.0183
1	.1397	.1304	.1217	.1135	.1057	.0984	.0915	.0850	.0789	.0733
2	.2165	.2087	.2008	.1929	.1850	.1771	.1692	.1615	.1539	.1465
3	.2237	.2226	.2209	.2186	.2158	.2125	.2087	.2046	.2001	.1954
4	.1734	.1781	.1823	.1858	.1888	.1912	.1931	.1944	.1951	.1954

Table C		<i>(continued)</i>									
		$\lambda$									
$x$	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	
5	.1075	.1140	.1203	.1264	.1322	.1377	.1429	.1477	.1522	.1563	
6	.0555	.0608	.0662	.0716	.0771	.0826	.0881	.0936	.0989	.1042	
7	.0246	.0278	.0312	.0348	.0385	.0425	.0466	.0508	.0551	.0595	
8	.0095	.0111	.0129	.0148	.0169	.0191	.0215	.0241	.0269	.0298	
9	.0033	.0040	.0047	.0056	.0066	.0076	.0089	.0102	.0116	.0132	
10	.0010	.0013	.0016	.0019	.0023	.0028	.0033	.0039	.0045	.0053	
11	.0003	.0004	.0005	.0006	.0007	.0009	.0011	.0013	.0016	.0019	
12	.0001	.0001	.0001	.0002	.0002	.0003	.0003	.0004	.0005	.0006	
13	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0001	.0002	.0002	
14	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001	
		$\lambda$									
$x$	4.1	4.2	4.3	4.4	4.5	4.6	4.7	4.8	4.9	5.0	
0	.0166	.0150	.0136	.0123	.0111	.0101	.0091	.0082	.0074	.0067	
1	.0679	.0630	.0583	.0540	.0500	.0462	.0427	.0395	.0365	.0337	
2	.1393	.1323	.1254	.1188	.1125	.1063	.1005	.0948	.0894	.0842	
3	.1904	.1852	.1798	.1743	.1687	.1631	.1574	.1517	.1460	.1404	
4	.1951	.1944	.1933	.1917	.1898	.1875	.1849	.1820	.1789	.1755	
5	.1600	.1633	.1662	.1687	.1708	.1725	.1738	.1747	.1753	.1755	
6	.1093	.1143	.1191	.1237	.1281	.1323	.1362	.1398	.1432	.1462	
7	.0640	.0686	.0732	.0778	.0824	.0869	.0914	.0959	.1002	.1044	
8	.0328	.0360	.0393	.0428	.0463	.0500	.0537	.0575	.0614	.0653	
9	.0150	.0168	.0188	.0209	.0232	.0255	.0280	.0307	.0334	.0363	
10	.0061	.0071	.0081	.0092	.0104	.0118	.0132	.0147	.0164	.0181	
11	.0023	.0027	.0032	.0037	.0043	.0049	.0056	.0064	.0073	.0082	
12	.0008	.0009	.0011	.0014	.0016	.0019	.0022	.0026	.0030	.0034	
13	.0002	.0003	.0004	.0005	.0006	.0007	.0008	.0009	.0011	.0013	
14	.0001	.0001	.0001	.0001	.0002	.0002	.0003	.0003	.0004	.0005	
15	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0001	.0001	.0002	
		$\lambda$									
$x$	5.1	5.2	5.3	5.4	5.5	5.6	5.7	5.8	5.9	6.0	
0	.0061	.0055	.0050	.0045	.0041	.0037	.0033	.0030	.0027	.0025	
1	.0311	.0287	.0265	.0244	.0225	.0207	.0191	.0176	.0162	.0149	
2	.0793	.0746	.0701	.0659	.0618	.0580	.0544	.0509	.0477	.0446	
3	.1348	.1293	.1239	.1185	.1133	.1082	.1033	.0985	.0938	.0892	
4	.1719	.1681	.1641	.1600	.1558	.1515	.1472	.1428	.1383	.1339	

Table C		<i>(continued)</i>									
x	$\lambda$										
	5.1	5.2	5.3	5.4	5.5	5.6	5.7	5.8	5.9	6.0	
5	.1753	.1748	.1740	.1728	.1714	.1697	.1678	.1656	.1632	.1606	
6	.1490	.1515	.1537	.1555	.1571	.1584	.1594	.1601	.1605	.1606	
7	.1086	.1125	.1163	.1200	.1234	.1267	.1298	.1326	.1353	.1377	
8	.0692	.0731	.0771	.0810	.0849	.0887	.0925	.0962	.0998	.1033	
9	.0392	.0423	.0454	.0486	.0519	.0552	.0586	.0620	.0654	.0688	
10	.0200	.0220	.0241	.0262	.0285	.0309	.0334	.0359	.0386	.0413	
11	.0093	.0104	.0116	.0129	.0143	.0157	.0173	.0190	.0207	.0225	
12	.0039	.0045	.0051	.0058	.0065	.0073	.0082	.0092	.0102	.0113	
13	.0015	.0018	.0021	.0024	.0028	.0032	.0036	.0041	.0046	.0052	
14	.0006	.0007	.0008	.0009	.0011	.0013	.0015	.0017	.0019	.0022	
15	.0002	.0002	.0003	.0003	.0004	.0005	.0006	.0007	.0008	.0009	
16	.0001	.0001	.0001	.0001	.0001	.0002	.0002	.0002	.0003	.0003	
17	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0001	
x	$\lambda$										
	6.1	6.2	6.3	6.4	6.5	6.6	6.7	6.8	6.9	7.0	
0	.0022	.0020	.0018	.0017	.0015	.0014	.0012	.0011	.0010	.0009	
1	.0137	.0126	.0116	.0106	.0098	.0090	.0082	.0076	.0070	.0064	
2	.0417	.0390	.0364	.0340	.0318	.0296	.0276	.0258	.0240	.0223	
3	.0848	.0806	.0765	.0726	.0688	.0652	.0617	.0584	.0552	.0521	
4	.1294	.1249	.1205	.1162	.1118	.1076	.1034	.0992	.0952	.0912	
5	.1579	.1549	.1519	.1487	.1454	.1420	.1385	.1349	.1314	.1277	
6	.1605	.1601	.1595	.1586	.1575	.1562	.1546	.1529	.1511	.1490	
7	.1399	.1418	.1435	.1450	.1462	.1472	.1480	.1486	.1489	.1490	
8	.1066	.1099	.1130	.1160	.1188	.1215	.1240	.1263	.1284	.1304	
9	.0723	.0757	.0791	.0825	.0858	.0891	.0923	.0954	.0985	.1014	
10	.0441	.0469	.0498	.0528	.0558	.0588	.0618	.0649	.0679	.0710	
11	.0245	.0265	.0285	.0307	.0330	.0353	.0377	.0401	.0426	.0452	
12	.0124	.0137	.0150	.0164	.0179	.0194	.0210	.0227	.0245	.0264	
13	.0058	.0065	.0073	.0081	.0089	.0098	.0108	.0119	.0130	.0142	
14	.0025	.0029	.0033	.0037	.0041	.0046	.0052	.0058	.0064	.0071	
15	.0010	.0012	.0014	.0016	.0018	.0020	.0023	.0026	.0029	.0033	
16	.0004	.0005	.0005	.0006	.0007	.0008	.0010	.0011	.0013	.0014	
17	.0001	.0002	.0002	.0002	.0003	.0003	.0004	.0004	.0005	.0006	
18	.0000	.0001	.0001	.0001	.0001	.0001	.0001	.0002	.0002	.0002	
19	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0001	.0001	

Table C		<i>(continued)</i>									
x	$\lambda$										
	7.1	7.2	7.3	7.4	7.5	7.6	7.7	7.8	7.9	8.0	
0	.0008	.0007	.0007	.0006	.0006	.0005	.0005	.0004	.0004	.0003	
1	.0059	.0054	.0049	.0045	.0041	.0038	.0035	.0032	.0029	.0027	
2	.0208	.0194	.0180	.0167	.0156	.0145	.0134	.0125	.0116	.0107	
3	.0492	.0464	.0438	.0413	.0389	.0366	.0345	.0324	.0305	.0286	
4	.0874	.0836	.0799	.0764	.0729	.0696	.0663	.0632	.0602	.0573	
5	.1241	.1204	.1167	.1130	.1094	.1057	.1021	.0986	.0951	.0916	
6	.1468	.1445	.1420	.1394	.1367	.1339	.1311	.1282	.1252	.1221	
7	.1489	.1486	.1481	.1474	.1465	.1454	.1442	.1428	.1413	.1396	
8	.1321	.1337	.1351	.1363	.1373	.1382	.1388	.1392	.1395	.1396	
9	.1042	.1070	.1096	.1121	.1144	.1167	.1187	.1207	.1224	.1241	
10	.0740	.0770	.0800	.0829	.0858	.0887	.0914	.0941	.0967	.0993	
11	.0478	.0504	.0531	.0558	.0585	.0613	.0640	.0667	.0695	.0722	
12	.0283	.0303	.0323	.0344	.0366	.0388	.0411	.0434	.0457	.0481	
13	.0154	.0168	.0181	.0196	.0211	.0227	.0243	.0260	.0278	.0296	
14	.0078	.0086	.0095	.0104	.0113	.0123	.0134	.0145	.0157	.0169	
15	.0037	.0041	.0046	.0051	.0057	.0062	.0069	.0075	.0083	.0090	
16	.0016	.0019	.0021	.0024	.0026	.0030	.0033	.0037	.0041	.0045	
17	.0007	.0008	.0009	.0010	.0012	.0013	.0015	.0017	.0019	.0021	
18	.0003	.0003	.0004	.0004	.0005	.0006	.0006	.0007	.0008	.0009	
19	.0001	.0001	.0001	.0002	.0002	.0002	.0003	.0003	.0003	.0004	
20	.0000	.0000	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0002	
21	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0001	

x	$\lambda$									
	8.1	8.2	8.3	8.4	8.5	8.6	8.7	8.8	8.9	9.0
0	.0003	.0003	.0002	.0002	.0002	.0002	.0002	.0002	.0001	.0001
1	.0025	.0023	.0021	.0019	.0017	.0016	.0014	.0013	.0012	.0011
2	.0100	.0092	.0086	.0079	.0074	.0068	.0063	.0058	.0054	.0050
3	.0269	.0252	.0237	.0222	.0208	.0195	.0183	.0171	.0160	.0150
4	.0544	.0517	.0491	.0466	.0443	.0420	.0398	.0377	.0357	.0337
5	.0882	.0849	.0816	.0784	.0752	.0722	.0692	.0663	.0635	.0607
6	.1191	.1160	.1128	.1097	.1066	.1034	.1003	.0972	.0941	.0911
7	.1378	.1358	.1338	.1317	.1294	.1271	.1247	.1222	.1197	.1171
8	.1395	.1392	.1388	.1382	.1375	.1366	.1356	.1344	.1332	.1318
9	.1256	.1269	.1280	.1290	.1299	.1306	.1311	.1315	.1317	.1318

<b>Table C</b>		<i>(continued)</i>									
<i>x</i>	$\lambda$										
	8.1	8.2	8.3	8.4	8.5	8.6	8.7	8.8	8.9	9.0	
10	.1017	.1040	.1063	.1084	.1104	.1123	.1140	.1157	.1172	.1186	
11	.0749	.0776	.0802	.0828	.0853	.0878	.0902	.0925	.0948	.0970	
12	.0505	.0530	.0555	.0579	.0604	.0629	.0654	.0679	.0703	.0728	
13	.0315	.0334	.0354	.0374	.0395	.0416	.0438	.0459	.0481	.0504	
14	.0182	.0196	.0210	.0225	.0240	.0256	.0272	.0289	.0306	.0324	
15	.0098	.0107	.0116	.0126	.0136	.0147	.0158	.0169	.0182	.0194	
16	.0050	.0055	.0060	.0066	.0072	.0079	.0086	.0093	.0101	.0109	
17	.0024	.0026	.0029	.0033	.0036	.0040	.0044	.0048	.0053	.0058	
18	.0011	.0012	.0014	.0015	.0017	.0019	.0021	.0024	.0026	.0029	
19	.0005	.0005	.0006	.0007	.0008	.0009	.0010	.0011	.0012	.0014	
20	.0002	.0002	.0002	.0003	.0003	.0004	.0004	.0005	.0005	.0006	
21	.0001	.0001	.0001	.0001	.0001	.0002	.0002	.0002	.0002	.0003	
22	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0001	.0001	.0001	
<i>x</i>	$\lambda$										
	9.1	9.2	9.3	9.4	9.5	9.6	9.7	9.8	9.9	10.0	
0	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0000	
1	.0010	.0009	.0009	.0008	.0007	.0007	.0006	.0005	.0005	.0005	
2	.0046	.0043	.0040	.0037	.0034	.0031	.0029	.0027	.0025	.0023	
3	.0140	.0131	.0123	.0115	.0107	.0100	.0093	.0087	.0081	.0076	
4	.0319	.0302	.0285	.0269	.0254	.0240	.0226	.0213	.0201	.0189	
5	.0581	.0555	.0530	.0506	.0483	.0460	.0439	.0418	.0398	.0378	
6	.0881	.0851	.0822	.0793	.0764	.0736	.0709	.0682	.0656	.0631	
7	.1145	.1118	.1091	.1064	.1037	.1010	.0982	.0955	.0928	.0901	
8	.1302	.1286	.1269	.1251	.1232	.1212	.1191	.1170	.1148	.1126	
9	.1317	.1315	.1311	.1306	.1300	.1293	.1284	.1274	.1263	.1251	
10	.1198	.1210	.1219	.1228	.1235	.1241	.1245	.1249	.1250	.1251	
11	.0991	.1012	.1031	.1049	.1067	.1083	.1098	.1112	.1125	.1137	
12	.0752	.0776	.0799	.0822	.0844	.0866	.0888	.0908	.0928	.0948	
13	.0526	.0549	.0572	.0594	.0617	.0640	.0662	.0685	.0707	.0729	
14	.0342	.0361	.0380	.0399	.0419	.0439	.0459	.0479	.0500	.0521	
15	.0208	.0221	.0235	.0250	.0265	.0281	.0297	.0313	.0330	.0347	
16	.0118	.0127	.0137	.0147	.0157	.0168	.0180	.0192	.0204	.0217	
17	.0063	.0069	.0075	.0081	.0088	.0095	.0103	.0111	.0119	.0128	
18	.0032	.0035	.0039	.0042	.0046	.0051	.0055	.0060	.0065	.0071	
19	.0015	.0017	.0019	.0021	.0023	.0026	.0028	.0031	.0034	.0037	

<b>Table C</b>		<i>(continued)</i>									
<i>x</i>	$\lambda$										
	8.1	8.2	8.3	8.4	8.5	8.6	8.7	8.8	8.9	9.0	
10	.1017	.1040	.1063	.1084	.1104	.1123	.1140	.1157	.1172	.1186	
11	.0749	.0776	.0802	.0828	.0853	.0878	.0902	.0925	.0948	.0970	
12	.0505	.0530	.0555	.0579	.0604	.0629	.0654	.0679	.0703	.0728	
13	.0315	.0334	.0354	.0374	.0395	.0416	.0438	.0459	.0481	.0504	
14	.0182	.0196	.0210	.0225	.0240	.0256	.0272	.0289	.0306	.0324	
15	.0098	.0107	.0116	.0126	.0136	.0147	.0158	.0169	.0182	.0194	
16	.0050	.0055	.0060	.0066	.0072	.0079	.0086	.0093	.0101	.0109	
17	.0024	.0026	.0029	.0033	.0036	.0040	.0044	.0048	.0053	.0058	
18	.0011	.0012	.0014	.0015	.0017	.0019	.0021	.0024	.0026	.0029	
19	.0005	.0005	.0006	.0007	.0008	.0009	.0010	.0011	.0012	.0014	
20	.0002	.0002	.0002	.0003	.0003	.0004	.0004	.0005	.0005	.0006	
21	.0001	.0001	.0001	.0001	.0001	.0002	.0002	.0002	.0002	.0003	
22	.0000	.0000	.0000	.0000	.0001	.0001	.0001	.0001	.0001	.0001	
<i>x</i>	$\lambda$										
	9.1	9.2	9.3	9.4	9.5	9.6	9.7	9.8	9.9	10.0	
0	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0000	
1	.0010	.0009	.0009	.0008	.0007	.0007	.0006	.0005	.0005	.0005	
2	.0046	.0043	.0040	.0037	.0034	.0031	.0029	.0027	.0025	.0023	
3	.0140	.0131	.0123	.0115	.0107	.0100	.0093	.0087	.0081	.0076	
4	.0319	.0302	.0285	.0269	.0254	.0240	.0226	.0213	.0201	.0189	
5	.0581	.0555	.0530	.0506	.0483	.0460	.0439	.0418	.0398	.0378	
6	.0881	.0851	.0822	.0793	.0764	.0736	.0709	.0682	.0656	.0631	
7	.1145	.1118	.1091	.1064	.1037	.1010	.0982	.0955	.0928	.0901	
8	.1302	.1286	.1269	.1251	.1232	.1212	.1191	.1170	.1148	.1126	
9	.1317	.1315	.1311	.1306	.1300	.1293	.1284	.1274	.1263	.1251	
10	.1198	.1210	.1219	.1228	.1235	.1241	.1245	.1249	.1250	.1251	
11	.0991	.1012	.1031	.1049	.1067	.1083	.1098	.1112	.1125	.1137	
12	.0752	.0776	.0799	.0822	.0844	.0866	.0888	.0908	.0928	.0948	
13	.0526	.0549	.0572	.0594	.0617	.0640	.0662	.0685	.0707	.0729	
14	.0342	.0361	.0380	.0399	.0419	.0439	.0459	.0479	.0500	.0521	
15	.0208	.0221	.0235	.0250	.0265	.0281	.0297	.0313	.0330	.0347	
16	.0118	.0127	.0137	.0147	.0157	.0168	.0180	.0192	.0204	.0217	
17	.0063	.0069	.0075	.0081	.0088	.0095	.0103	.0111	.0119	.0128	
18	.0032	.0035	.0039	.0042	.0046	.0051	.0055	.0060	.0065	.0071	
19	.0015	.0017	.0019	.0021	.0023	.0026	.0028	.0031	.0034	.0037	

Table C (continued)		$\lambda$									
$x$	9.1	9.2	9.3	9.4	9.5	9.6	9.7	9.8	9.9	10.0	
20	.0007	.0008	.0009	.0010	.0011	.0012	.0014	.0015	.0017	.0019	
21	.0003	.0003	.0004	.0004	.0005	.0006	.0006	.0007	.0008	.0009	
22	.0001	.0001	.0002	.0002	.0002	.0002	.0003	.0003	.0004	.0004	
23	.0000	.0001	.0001	.0001	.0001	.0001	.0001	.0001	.0002	.0002	
24	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0001	.0001	.0001	
$x$	11	12	13	14	15	16	17	18	19	20	
0	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	
1	.0002	.0001	.0000	.0000	.0000	.0000	.0000	.0000	.0000	.0000	
2	.0010	.0004	.0002	.0001	.0000	.0000	.0000	.0000	.0000	.0000	
3	.0037	.0018	.0008	.0004	.0002	.0001	.0000	.0000	.0000	.0000	
4	.0102	.0053	.0027	.0013	.0006	.0003	.0001	.0001	.0000	.0000	
5	.0224	.0127	.0070	.0037	.0019	.0010	.0005	.0002	.0001	.0001	
6	.0411	.0255	.0152	.0087	.0048	.0026	.0014	.0007	.0004	.0002	
7	.0646	.0437	.0281	.0174	.0104	.0060	.0034	.0018	.0010	.0005	
8	.0888	.0655	.0457	.0304	.0194	.0120	.0072	.0042	.0024	.0013	
9	.1085	.0874	.0661	.0473	.0324	.0213	.0135	.0083	.0050	.0029	
10	.1194	.1048	.0859	.0663	.0486	.0341	.0230	.0150	.0095	.0058	
11	.1194	.1144	.1015	.0844	.0663	.0496	.0355	.0245	.0164	.0106	
12	.1094	.1144	.1099	.0984	.0829	.0661	.0504	.0368	.0259	.0176	
13	.0926	.1056	.1099	.1060	.0956	.0814	.0658	.0509	.0378	.0271	
14	.0728	.0905	.1021	.1060	.1024	.0930	.0800	.0655	.0514	.0387	
15	.0534	.0724	.0885	.0989	.1024	.0992	.0906	.0786	.0650	.0516	
16	.0367	.0543	.0719	.0866	.0960	.0992	.0963	.0884	.0772	.0646	
17	.0237	.0383	.0550	.0713	.0847	.0934	.0963	.0936	.0863	.0760	
18	.0145	.0256	.0397	.0554	.0706	.0830	.0909	.0936	.0911	.0844	
19	.0084	.0161	.0272	.0409	.0557	.0699	.0814	.0887	.0911	.0888	
20	.0046	.0097	.0177	.0286	.0418	.0559	.0692	.0798	.0866	.0888	
21	.0024	.0055	.0109	.0191	.0299	.0426	.0560	.0684	.0783	.0846	
22	.0012	.0030	.0065	.0121	.0204	.0310	.0433	.0560	.0676	.0769	
23	.0006	.0016	.0037	.0074	.0133	.0216	.0320	.0438	.0559	.0669	
24	.0003	.0008	.0020	.0043	.0083	.0144	.0226	.0328	.0442	.0557	
25	.0001	.0004	.0010	.0024	.0050	.0092	.0154	.0237	.0336	.0446	
26	.0000	.0002	.0005	.0013	.0029	.0057	.0101	.0164	.0246	.0343	
27	.0000	.0001	.0002	.0007	.0016	.0034	.0063	.0109	.0173	.0254	
28	.0000	.0000	.0001	.0003	.0009	.0019	.0038	.0070	.0117	.0181	
29	.0000	.0000	.0001	.0002	.0004	.0011	.0023	.0044	.0077	.0125	