

UNIVERSITI SAINS MALAYSIA

First Semester Examination
Academic Session 2004/2005

October 2004

ZCT 307E/3 - Solid State Physics I
[Fizik Keadaan Pepejal I]

Duration 3 hours
[Masa 3 jam]

Please check that this examination paper consists of **THREE** pages of printed material before you begin the examination

[Sila pastikan bahawa kertas peperiksaan ini mengandungi TIGA muka surat yang bercetak sebelum anda memulakan peperiksaan ini.]

Instructions Answer all **FOUR** (4) questions. Students are allowed to answer all questions in Bahasa Malaysia or in English

Arahan: *Jawab kesemua EMPAT (4) soalan. Pelajar dibenarkan menjawab semua soalan sama ada dalam Bahasa Malaysia atau Bahasa Inggeris.]*

- 1 (a) Discuss the reasons for the failure of Dulong and Petit's law to predict the specific heat at low temperatures. Why should the law be valid at high temperatures?
 [(a) *Bincang mengapa hukum Dulong dan Petit tidak mampu meramal haba tentu pada suhu rendah. Kenapa hukum itu berlaku pada suhu tinggi?*]
- (b) Discuss how Debye's theory managed to overcome the shortcomings of the classical theory
 [(b) *Bincangkan bagaimana teori Debye mengatasi masalah di atas*]
- (c) Debye temperature of carbon is 1850 K. Compute the Debye frequency involved in the Debye theory
 [(c) *Suhu Debye bagi karbon adalah 1850 K. Tentukan frekuensi Debye bagi kes teori Debye*]
- (20/100)
- 2 (a) Discuss the failure of the classical free electron model with special reference to the specific heat of metals
 [(a) *Bincangkan kegagalan model klasik elektron bebas berasaskan haba tentu logam*]
- (b) Explain how Sommerfeld modified the free electron theory to remove this shortcomings
 [(b) *Jelaskan bagaimana model Sommerfeld mengubahsuaikan teori elektron bebas bagi menjelaskan perkara di atas*]
- (c) Estimate the fractions of electrons excited about the Fermi level at room temperature ($T = 300$ K). $E_F = 3.1$ eV for Na
 [(c) *Tentukan pecahan elektron yang teruja sekitar paras Fermi pada suhu bilik ($T = 300$ K). $E_F = 3.1$ eV bagi Na*]
- (25/100)
- 3 (a) The Equation below is derived from the Kronig-Penney Model for an electron in a periodic field,
 [(a) *Persamaan berikut telah diterbitkan daripada model Kronig-Penney.*]

$$P \frac{\sin \alpha a}{\alpha a} + \cos \alpha a = \cos ka$$

where [di mana] $P = \frac{mV_0ba}{\hbar^2}$ and [dan] $\alpha^2 = \frac{2mE}{\hbar^2}$

- (1) Explain the physical meaning of all the terms in the equation above
 [(i) *Jelaskan maksud fizikal bagi semua sebutan persamaan di atas*]

[11] Plot a graph of $P \frac{\sin \alpha a}{\alpha a} + \cos \alpha a$ versus αa Discuss what you can deduce from the graph

[(ii) *Plot graf $P \frac{\sin \alpha a}{\alpha a} + \cos \alpha a$ lawan αa Bincang apa yang anda perolehi dari graf tersebut*]

- (b) Show from the E-k graph that materials can be classified into conductors, insulators and semiconductors

[(b) *Dari graf E-k bagaimanakah anda boleh mengelaskan bahan konduktor, penebat dan semikonduktor*]

- [c] Prove that for the Kronig Penney model the energy of the lowest energy

band at $k = 0$ is $E = \frac{\hbar^2 P}{ma^2}$

[(c) *Buktikan dari model Kronig-Penney bahawa tenaga bagi jalur terendah*

$k = 0$ adalah $E = \frac{\hbar^2 P}{ma^2}$]

(30/100)

- 4 (a) Show that for the case of intrinsic semiconductors, the carrier concentration is given by

[(a) *Tunjukkan bagi kes semikonduktor intrinsik, kepekatan pembawa diberi oleh*

$$n = p = n_i = 2 \left(\frac{k_B T}{2\pi \hbar^2} \right)^{3/2} (m_e m_h)^{3/4} e^{-E_g / 2k_B T}$$

- [b] Mobilities of electrons and holes in a sample of intrinsic germanium at 300 K are $0.36 \text{ m}^2 \text{V}^{-1} \text{s}^{-1}$ and $0.17 \text{ m}^2 \text{V}^{-1} \text{s}^{-1}$ respectively. If the conductivity of the specimen is $2.12 \text{ } \Omega^{-1} \text{m}^{-1}$, compute the forbidden energy gap

[(b) *Kelincahan elektron dan lohong bagi sampel germanium intrinsik pada suhu 300 K adalah $0.36 \text{ m}^2 \text{V}^{-1} \text{s}^{-1}$ dan $0.17 \text{ m}^2 \text{V}^{-1} \text{s}^{-1}$ secara berturutan. Jika kekonduksian germanium adalah $2.12 \text{ } \Omega^{-1} \text{m}^{-1}$ hitung jurang tenaga germanium*]

(25/100)