





**LAPORAN AKHIR PROJEK PENYELIDIKAN
R & D JANGKA PENDEK**

"In vivo toxicity study of a plant *Phyllanthus amarus*, in rat"



Investigators:

**Dr. K.N.S. Sirajudeen
Prof. Madya Dr. Siti Amrah Sulaiman
Prof. Madya Dr. M. Madhavan
Prof. Madya Dr. Zabidah Ismail**



Grant No: 304/PPSP/6131166

| | |
|--|--------------------------------|
| BAHAGIAN PENYELIDIKAN PUSAT PENGAJIAN SAINS PERUBATAN | |
| SALINAN : | |
| <input type="checkbox"/> | Bhg. Penyelidikan, PPSP |
| <input type="checkbox"/> | Perpustakaan Perubatan, USM/KK |
| <input checked="" type="checkbox"/> | RCMO |
| Tangan : Tarikh : 2/5/05 | |

TABLE OF CONTENTS

| | PAGE |
|-------------------------|------|
| ABSTRACT | 1 |
| 1 INTRODUCTION | 3 |
| 2 MATERIALS AND METHODS | 6 |
| 3 RESULTS | 12 |
| 4 DISCUSSION | 25 |
| 5. CONCLUSION | 27 |
| ACKNOWLEDGEMENTS | 28 |
| REFERENCES | 29 |

“Kajian toksikiti in vivo *Phyllanthus amarus* terhadap tikus”

ABSTRAK

Pengenalan

Di Malaysia dan juga di negara lain, *Phyllanthus amarus* (*P. amarus*), sejenis spesies pokok dari keluarga Euphorbiaceae digunakan oleh pengamal perubatan tradisional bagi merawat jaundis dan penyakit lain. Adalah dilaporkan efikasi spesies ini berbeza dari segi kawasan geografi dan kepelbagaian dan perbezaan ini banyak dikaitkan dengan kandungan bahan di dalamnya. Bagaimanapun, buat masa ini tiada kajian toksikiti yang lengkap dengan dokumentasi dijalankan di Malaysia.

Objektif

Tujuan kajian awal ini ialah untuk menentukan kesan toksik, jika ada, juga profil keselamatan ekstrak daun berakues *P. amarus* yang diberikan secara oral ke atas hepar tikus dengan menaksirkan perubahan morfologi, biokimia dan histologi.

Bahan dan metod

Tikus Sprague-Dawley (berat 180-230 g) digunakan sebagai model haiwan untuk kajian ini. Untuk kajian toksikiti akut, tikus jantan diberikan ekstrak daun akues *P. amarus* (5g/kg) dan untuk kajian toksikiti kronik, tikus jantan dan betina diberikan dos 100g, 400g dan 800g/kg selama 6 minggu. Haiwan kawalan juga disediakan tanpa diberikan sediaan ekstrak. Berat badan tikus diambil pada permulaan kajian dan berskala sehingga tamat kajian. Selepas masa kajian (6 minggu), haiwan akan dikorbankan dan homogenat hepar tikus dianalisa untuk mendapatkan sebarang petanda biokimia kecederaan hepar, alanin transaminas (ALT), aspartat transaminas (AST), alkalin fosfatas (ALP), laktat dehidrogenas (LDH) dan jumlah protein serum. Bahagian hepar juga diambil untuk kajian histologi – mikroskopi ringan, kajian antigen nuklear sel proliferatif (PCNA) dan kajian apoptik dengan menggunakan Tag kit apop.

Keputusan

Pemberian akut ekstrak *P. amarus* (5g/kg) tidak memberikan sebarang tanda toksikiti atau kematian. Manakala kajian toksikiti kronik tiada menunjukkan perbezaan signifikan ($P > 0.05$) dari segi berat badan juga petanda biokimia (ALT, AST, ALP, LDH dan jumlah protein) di antara haiwan kawalan dengan haiwan yang diberikan ekstrak *P. amarus* sepanjang masa kajian. Ekstrak *P. amarus* yang tidak toksik dibuktikan oleh kajian histologi i.e. tiada kelihatan perbezaan di antara tikus kawalan dengan tikus yang memakan ekstrak *P. amarus*.

Rumusan

Pemberian akut ekstrak *P. amarus* pada dos 5g/kg tidak memberikan kesan toksik kepada hepar tikus. Kajian toksikiti kronik pula menunjukkan tiada toksikiti kumulatif seperti yang ditunjukkan oleh tiada perubahan signifikan ke atas parameter yang dikaji, juga keputusan kajian histologi.

Kata kunci: *Phyllanthus amarus*, ekstrak akues, hepar, toksikiti, enzim petanda, kajian histologi

ABSTRACT

Introduction: *Phyllanthus amarus* (*P. amarus*), a plant species of *Euphorbiaceae* family, is used as a folk medicine for jaundice and other diseases in Malaysia and other countries. The variation in their efficacy with geographical location and varieties has been reported which has been attributed to their constituent composition. But, so far no toxicity studies have been carried out on this plant with clear documentation, especially with those plants growing in Malaysia.

Objective: The aim of this preliminary study was to determine the toxic side effects of aqueous extract of leaves of *P. amarus* (grown in Malaysia) following oral administration in rats by assessing the morphological, biochemical and histological changes in them.

Materials & Methods: Sprague-Dawley rats weighing (180-230 grams) were used as animal models in this study. Aqueous extract of leaves of *P. amarus* was administered orally to male rats for acute toxicity study (5 gram / Kg body weight) and to the male and female rats for chronic toxicity study (at the doses of 100, 400 and 800 mg/ Kg body weight/day for six weeks). Control of male and female rats was also maintained without plant extract administration. Body weight of the rats was taken initially and periodically once a week, till the end of experimental period. At the end of experimental period (6 weeks), the rats were sacrificed and analyzed for biochemical markers of liver injury - alanine transaminase (ALT), aspartate transaminase (AST), alkaline phosphatase (ALP), lactate dehydrogenase (LDH) and total protein in the serum collected and the marker enzymes were assayed in the homogenates of the rat liver. Liver sections were taken for histological studies; Light microscopy, proliferative cell nuclear antigen (PCNA) study and apoptotic study by using ApopTag kit .

Results: Acute administration of *P. amarus* extract, orally, even at a dose of 5 gram/Kg body weight) did not produce any signs of toxicity or mortality. In the chronic study, no significant differences ($p>0.05$) were observed between the control and *P. amarus* extract administered rats in the body weight gain during the experimental period as well as in the biochemical markers analyzed (ALT, AST, ALP, LDH and total protein) in serum and liver homogenates. The non-toxic nature of *P. amarus* extract administration was confirmed by histological studies i.e., no observable changes were found between control and *P. amarus* extract administered rats.

Conclusion: Acute oral administration of *P. amarus* extract is non-toxic to the rat liver, even at a dose of 5 gram /kg body weight. The chronic toxicity studies of *P. amarus* extracts administration (of 100-800 mg/kg body wt) showed the absence of cumulative toxicity as reflected by the non-significant change in the parameters studied as well as from the results of the histological studies.

Key words: *Phyllanthus amarus*, Aqueous extract, Liver, Toxicity, Marker enzymes, Histological studies

1. INTRODUCTION

Herbal medicines has been used since ancient times. Most of the population of the underdeveloped and developing countries depend on some form of traditional medicine. Malaysia is one of the important nations of biodiversity in the world and enriched with natural plant resources (Muhamad bin Zakaria and Mustafa Ali Mohd, 1994).

One of the plant genres widely used traditionally for the treatment of different diseases is *Phyllanthus* (Family: *Euphoribaceae*) and is distributed in most tropical and subtropical countries and comprise approximately of 550-750 species throughout the world. Among them, one of the most studied species is *P.amarus* and is widely used in Malaysia and other countries (Calixto *et al.*, 1998; Muhamad bin Zakaria and Mustafa Ali Mohd, 1994).

Some of the vernacular names of the plant include: Dukong anak (Malay), Kilanelli (Tamil), Bhumyamalakai (Sanskrit), Chanca piedra, Quebra pedra (Brazil).

P.amarus has bitter, astringent, cooling, diuretic, stomachic, antiseptic, antiviral, antidiabetic, hypotensive, antinociceptive, febrifuge properties and is traditionally used in the treatment of jaundice, diarrhea, dysentery, diabetes, fevers, uro-genital diseases, ulcers and wounds (Santos *et al*, 1995; Calixto *et al*, 1998). Recent years, a growing interest shown towards the *Phyllanthus* with respect to their potential on management of several diseases (Odetola and

Akojenu, 2000; Rajeshkumar *et al.*, 2002; Srividya and Periwai, 1995) More research was done on its antiviral effect. Some reports have shown the antiviral effect of this plant by reducing the detectable hepatitis B surface antigen (HbsAg) of HBV positive patients (Thyagarajan *et al.*, 1988; Ott *et al.*, 1997). But other studies like from China, Thailand , India showed the failure of *P. amarus* in eradicating the HbsAg in patients with chronic hepatitis B virus (Leelarasamee *et al.*, 1990; Wang *et al.*, 1991, Doshi *et al.*, 1997). The variation in the clinical effect of these studies has been attributed to many factors such as different species, differences in growing condition and different processing methods (Thyagarajan *et al.*, 2002; Wang, 2000).

Usually herbal medicines are widely perceived by the public as being natural, healthful and free from side effects, but that is not 100 % true. Plants contain hundreds of constituents and some of them may elicit a toxic side effects. A number of studies exist reporting the toxic effect of herbal medicines (Shaw *et al.*, 1997; Kaplowitz, 1997; Calixto, 2000). Therefore efficacy and safety study should be performed in these herbs.

Eventhough a large number of clinical trials has been done on *P. amarus* and reporting their benefits, so far no systematic toxicological investigation has been reported on this plant, especially the *P. amarus* growing in Malaysia.

Since the efficacy of *Phyllanthus* species varies with geographical location and varieties and because of their constituent composition variation, the

P. amarus grown in Malaysia (widely used here as the folk medicine for jaundice) has to be ascertained for their biosafety by conducting acute and chronic toxicity study .

So the aim of this preliminary study was an attempt to determine the toxic effects of aqueous extract of leaves of *P. amarus* (grown in Malaysia) following oral administration in rats by assessing the morphological, biochemical and histological changes.

2 MATERIALS AND METHODS

2.1 Plant material

P. amarus was collected from MARDI (Malaysian Agricultural Research and Development Institute, Telung, Bachok, Kelantan) during the months of July-October and identified by FRIM (Forest Research Institute of Malaysia, Kepong, Selangor). The aqueous extraction of *P. amarus* leaves was carried out at the Department of Pharmacology, School of Medical Sciences, Universiti Sains Malaysia.

2.2 Preparation of plant extract

The plant material was cleaned and the leaves were separated, oven dried at 50°C. Then macerated into dry powder. Approximately 4.5 grams of fine leaves powder of *P. amarus* were extracted with distilled water (2:25) using soxhlet apparatus. After extraction, concentrated by rotary evaporator at 65°C. Then transferred into a suitable container and freeze dried by freeze dryer. The yield of the final crude aqueous *P. amarus* extract was 6-9% (~380 grams). The dried extract was stored in a desicator until its use. The extract was dissolved in distilled water to the desired concentration just before the study.

2.3 Animal and experimental design

Male and female Sprague-Dawley rats weighing (180-220 grams) were used as the experimental animal in this study and were obtained from Animal House Facility Unit, Health Campus, Universiti Sains Malaysia and acclimatized for one week prior to start the experiment. Animals were

housed in a standard cages at a temperature of $23\pm 2^{\circ}\text{C}$, 45-55 % relative humidity with a 12h light/12h dark cycle. The animals were fed with commercial pellet diet and water *ad libitum*.

Our study protocol was approved by the animal ethical committee, Health Campus, Universiti Sains Malaysia.

Two types of studies were carried out (WHO, 1993; Arnold, 1990) Acute and chronic study and in both studies the rats were administered with *P.amarus* extract by gavage (orally).

2.4 Acute study

Aqueous leaves extract of *P.amarus* at a single dose of 5g/Kg body weight was administered orally to two male rats. Another two male rats served as control (without *P.amarus* extract administration). The animals were observed carefully for any visible signs of toxicity and mortality immediately after dosing, at 4h, 24h intervals, during the recovery period of 48h and twice daily upto 14 days. After 14 days, the rats were sacrificed under ether anesthesia. A thorough autopsy was carried out and all organs were observed for any macroscopic changes.

2.5 Chronic study

2.5.1 Animal groups

For the chronic study, the animals were grouped (ten rats in each group) as follows

| Male rats | Female rats |
|--|--|
| Group I – Control Group II, III & IV – Rats administered with <i>P.amarus</i> extract at the doses of 100,400 and 800 mg/Kg body wt/day for 6 weeks by gavage, respectively | Group A – Control Group B, C & D – Rats administered with <i>P.amarus</i> extract at the doses of 100,400 and 800 mg/Kg body wt/day for 6 weeks by gavage, respectively |

The animals were observed daily for any signs of morbidity and mortality and their body weights were measured periodically in the experimental period.

2.5.2 Collection of serum and liver samples for analysis

At the end of the experimental period (6 weeks), after an overnight fasting, both the male and female rat controls and *P.amarus* extract administered groups were sacrificed by decapitation. Blood was collected, allowed to clot and then centrifuged at 3,000 rpm for 15 minutes. Serum samples were separated and used for biochemical analysis. The samples were stored at -80°C, if not used immediately.

Liver tissues were excised and washed in ice-cold saline and an weighed amount was homogenized with 0.1M Tris buffer at pH 7.4 under cold

condition. The supernatant of the liver homogenates was used for the biochemical estimations.

A portion of the liver tissue of all the (control and *P.amarus* extract administered) groups were fixed in 10% formal saline for the histological studies.

2.5.3 Biochemical analysis

In the collected serum, the total protein and the activities of the liver marker enzymes such as ALT,AST,ALP and LDH were assayed by using standard kit (ALT Randox kit , Randox Total protein biuret reagent- Randox Laboratories Ltd, UK; AST,ALP and LDH Roche Kits - Roche diagnostics, GmbH, Germany) in a Hitachi-912 autoanalyser available in our department of chemical Pathology, School of Medical Sciences, Universiti Sains Malaysia.

Similarly the activities of the marker enzymes – ALT,AST, ALP and LDH were assayed in the liver homogenates prepared.

2.5.4 Histological studies

This study was carried out in the department of Pathology, School of Medical Sciences, Universiti Sains Malaysia.

A portion of all the liver specimens fixed in 10% formal saline were processed routinely overnight using histokinette. Then they were embedded in paraffin was. Three sections, each four micron in thickness were cut from each paraffin block.

Light Microscopic study

One section from each sample was stained with Haematoxylin & Eosin (H&E) stain by the standard method for light microscopic histological examination

Proliferative Cell Nuclear Antigen (PCNA) study

Another section from each sample was stained with anti PCNA (DAKO, Denmark) using standard immunohistochemistry procedure and the positivity was visualized with the chromogen diaminobenzidine (DAB) (DAKO, Denmark) and expressed in percentage.

Apoptosis study

The third section from each sample was assessed for apoptosis. They were stained with ApopTag –Apoptosis detection kit (Chemicon, USA) whereby the reagents are designed to label the free 3'OH DNA termini in situ with chemically labelled and unlabelled nucleotides. The nucleotides contained in the Reaction Buffer are enzymatically added to the DNA by terminal deoxynucleotidyl transferase (TdT). TdT catalyzes a template-independent addition of nucleotide triphosphates to the 3'OH ends of

double stranded or single stranded DNA. The incorporated nucleotides form an oligomer composed of digoxigenin-conjugated nucleotide and unlabelled nucleotide in a random sequence. The ratio of labelled to unlabelled nucleotide in ApopTag Peroxidase Kits is optimized to promote anti digoxigenin antibody binding. DNA fragments which have been labelled with digoxigenin-nucleotide are then allowed to bind an anti-digoxigenin antibody that is conjugated with to a peroxidase reporter molecule. The bound peroxidase antibody conjugate enzymatically generates a permanent, intense, localized stain from chromogenic substance, providing sensitive detection in immunohistochemistry. This mixed molecular biological-histochemical system allows for sensitive and specific staining of very high concentrations of 3'OH ends that are localised in apoptotic bodies.

2.6 Statistical Analysis

The results of the biochemical analysis are expressed as mean \pm S.D. for ten animals in each group. The difference between the control and P.amarus extract administered groups (for each sex of animals) have been analysed by student's t-test. p value <0.05 , considered as significant.

3. RESULTS

3.1 ACUTE STUDY

During the course of the acute study, no treatment related effect was observed on the general condition or behaviour of the experimental animals. All the rats appeared to be normal, survived during the experimental period and none of them showed any visible signs of toxicity. A thorough autopsy of the *P.amarus* administered rats revealed no treatment related macroscopic changes.

3.2 CHRONIC STUDY

3.2.1 Body weight

Mean body weight gain of the *P.amarus* extract administered groups (at the doses of 100,400 and 800 mg/Kg body weight/day) have shown no appreciable difference when compared to their control (both male and female rats) after 6 weeks duration of the study (Figure 1).

The macroscopic appearance of all the organs observed was found to be normal and there were no appreciable changes in the liver weight of *P.amarus* extract administered groups and control group (data not shown).

3.2.2 Biochemical study

There were no significant differences ($p>0.05$) observed in the biochemical parameters studied (ALT,AST,ALP,LDH and total protein) in

the serum between controls and *P.amarus* extract administered groups (of both male and female) (Figure 2 & 3).

Similarly, the parameters assayed (ALT,AST,ALP and LDH) in the liver homogenates of controls and *P.amarus* extract administered groups also showed no significant difference ($p>0.05$) (Figure 4 & 5).

Fig. 1 : Mean body weight gain of control and *P. amarus* administered groups, Male and Female rats at the end of experimental period






Figure 2: Level of total protein and activities of serum marker enzymes of control and *P. amarus* extract administered groups (male rats)




Values are mean \pm S.D. for ten animals in each group and analyzed by Student's t-test. Group II, III and IV compared with Group I (control) and p value <0.05 considered significant.
^{NS} $p>0.05$ - not significant

Figure 3: Level of total protein and activities of serum marker enzymes of control and *P. amarus* extract administered groups (female rats)



Values are mean \pm S.D. for ten animals in each group and analyzed by Student's t-test. Group B,C and D compared with Group A (control) and p value <0.05 considered significant.
NS $p > 0.05$ - not significant


Figure 4: Activities of ALT, AST, ALP and LDH in the liver homogenates of control and *P.amarus* extract administered groups (male rats)



Group I –Control;
Group II, III and IV – *P. amarus* extract administered groups at the doses of 100mg/Kg body wt/day; 400mg/Kg body wt/day; 800mg/Kg body wt/day, orally, for 6 weeks respectively.

Values are mean \pm S.D. for ten animals in each group and analyzed by Student's t-test.
Group II,III and IV compared with Group I (control) and p value <0.05 considered significant.
^{NS} p>0.05- not significant

Figure 5: Activities of ALT, AST, ALP and LDH in the liver homogenate of control and *P. amarus* extract administered groups (female rats)



Group A –Control;
Group B,C and D – *P. amarus* extract administered groups at the doses of 100mg/Kg body wt/day; 400mg/Kg body wt/day; 800mg/Kg body wt/day, orally, for 6 weeks respectively.

Values are mean \pm S.D. for ten animals in each group & analyzed by Student's t-test.
Group B,C and D compared with Group A (control) and p value <0.05 considered significant.
^{NS} $p>0.05$ - not significant

3.3 Histological study

3.3.3. Light microscopic study

Histological examination of the liver under light microscope showed no significant pathological changes in the *P.amarus* extract administered groups (at the acute dose of 5g/Kg body weight and at the doses of 100, 400 & 800 mg/Kg body wt/ day, orally, for six weeks to both male & female rats) (figures 8-14) when compared to their controls (Figures 6 & 7).




Fig. 6: Normal liver from control male rat (H&E)




Fig.7: Normal liver from control female rat (H&E)




Fig. 8: Liver appears normal in male rat with acute administration of *P.amarus* extract 5g/Kg body wt. (H&E)




Fig. 9: Liver from male rat administered with *P. amarus* extract 100 mg/Kg body wt/ day for 6 weeks (H&E)




Fig.10: Liver from female rat administered with *P. amarus* extract 100 mg/Kg body wt/ day for 6 weeks (H&E)




Fig. 11: Liver from male rat administered with *P. amarus* extract 400 mg/Kg body wt/ day for 6 weeks (H&E)




Fig.12: Liver from female rat administered with *P. amarus* extract 400 mg/Kg body wt/ day for 6 weeks (H&E)




Fig. 13: Liver from male rat administered with *P. amarus* extract 800 mg/Kg body wt/ day for 6 weeks (H&E)




Fig.14: Liver from female rat administered with *P. amarus* extract 800 mg/Kg body wt/ day for 6 weeks (H&E)

3.3.2 PCNA Study

Anti PCNA antibody was used to measure the proliferation of hepatocytes by immunohistochemistry. The cells that are in the proliferative pool shows brown colour staining (Figure 15). PCNA of liver from control rat was 1.9% (Figures 16 &17) and for the acute dose (5g/Kg body weight) *P.amarus* extract administered rat was 1.8% (Figure18). For the *P.amarus* extract administered groups at the doses of 100, 400 & 800 mg/Kg body wt/ day , orally, for 6 weeks (both male & female rat livers) showed PCNA of 0.2% -2.1% (Figures 19-24) which are similar to that of their controls.




Fig. 15: Lymphnode – positive control – shows increased proliferation in the germinal centre (GC) (anti PCNA)




Fig.16: Liver from control male rat shows only occasional cells in proliferative cycle (anti PCNA)




Fig.17: Liver from control female rat shows only occasional cells in proliferative cycle (anti PCNA)




Fig.18: Liver from male rat with acute administration of *P.amarus* extract 5g/Kg body wt. shows only occasional cells in proliferative cycle (anti PCNA)




Fig.19: Liver from male rat administered with *P.amarus* extract 100 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)




Fig.21: Liver from male rat administered with *P.amarus* extract 400 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)



Fig.23: Liver from male rat administered with *P.amarus* extract 800 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)




Fig.20: Liver from female rat administered with *P.amarus* extract 100 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)




Fig.22: Liver from female rat administered with *P.amarus* extract 400 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)




Fig.24: Liver from female rat administered with *P.amarus* extract 800 mg/Kg body wt/ day for 6 weeks shows only occasional cells in proliferative cycle (anti PCNA)

3.3.3 Apoptosis study

Apoptosis in the liver was measured in the hepatocytes using ApopTag Kit. The positivity was indicated by the brown colour in nucleus (Figure 25). Apoptosis in the livers from control (both male and female rats) were 0.2% (Figure 26 & 27). Apoptosis in the livers from *P.amarus* extract administered groups were ranged from 0.2% to 1.5% (Figures 28-34) which are similar to that of their controls.



Fig. 25: Apoptotic bodies are stained brown in a positive control slide (ApopTag kit)




Fig. 26: Normal control male liver shows occasional apoptotic bodies (ApopTag kit)




Fig. 27: Normal control female liver shows occasional apoptotic bodies (ApopTag kit)




Fig. 28: Male rat liver administered with acute *P.amarus* extract at a dose of 5g/Kg body wt, shows occasional apoptotic bodies (ApopTag kit)




Fig. 29: Liver from male rat administered with *P.amarus* extract 100 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)



Fig. 30: Liver from female rats administered with *P.amarus* extract 100 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)



Fig. 31: Liver from male rat administered with *P.amarus* extract 400 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)



Fig.32: Liver from female rat administered with *P.amarus* extract 400 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)




Fig. 33: Liver from male rat administered with *P.amarus* extract 800 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)




Fig. 34: Liver from female rat administered with *P.amarus* extract 800 mg/Kg body wt/ day for 6 weeks shows occasional apoptotic bodies (ApopTag kit)

4. DISCUSSION

Herbal medicines have played a major role in world health and are very popular in developing and underdeveloped countries with a long tradition in the use of medicinal plants. Reports indicate that the idea, herbal drugs are very safe and free from side effect is false (Shaw *et al.*, 1997; Kaplowitz, 1997; Calixto, 2000). Therefore, clear understanding of potential adverse effect of herbs used by human population is necessary for implementing safety measures to the public.

Since in the case of *P. amarus*, conflicting reports have been obtained regarding the efficacy of the plant grown in different countries and hence, a study on their toxicity is required. In our present work, we have studied the toxicity of *P. amarus* (growing in Malaysia) in rats .

The results of our acute study with the absence of lethality or toxic side effect on oral administration of *P. amarus* extract even at a dose of 5g/kg body wt clearly indicate the non-toxic nature of the plant extract. Toxicologists agree that any test substance that is not lethal on acute administration at a concentration of 5g/Kg body wt, is essentially non-toxic (OECD, 1981; Brock *et al.*, 1995)

In the chronic toxicity study, we did biochemical analysis and histological studies to determine the cumulative toxic effect *P. amarus* extract on rat liver .

Liver is an organ involved in many metabolic functions and is prone to xenobiotic-induced injury because of their central role in xenobiotic metabolism (Sturgill and Lambert, 1997). Liver contains a host of enzymes such as ALT, AST, LDH and ALP. The activities of these enzymes are used to assess the functional status of the liver and as the biochemical markers of liver damage (Moss and Ralph Handerson, 1999). Hepatotoxic drugs cause damage to the liver cell membrane and these enzymes are leaked out into serum and shows increased activities (Kumar et al, 2004; sturgill and Lambert, 1997). Our study results showed that there were no increased activities of ALT, AST, LDH and ALP on *P. amarus* extract administration indicating that the plant is non-hepatotoxic in nature.

The results of our biochemical estimations were confirmed by the histological studies (light microscopy, PCNA study and apoptosis study).

Histological examination of liver specimen is useful in identifying the type of injury present in it (Friedman *et al.*, 1996). Our light microscopic study showed no observable changes between control and *P. amarus* administered groups.

Researchers assessed the role of cell proliferation and compensatory tissue repair in response to initial liver injury by PCNA study (Wang *et al.*, 2000). The expression of PCNA by immunohistochemistry is a reliable indicator of both the proliferation rate and location of the compartment (Yamada *et al.*, 1992). We also found no difference between control and *P. amarus* administered groups in the PCNA study.

In assessing cell proliferation, a strong correlation was found between the PCNA study and ^3H -Thymidine incorporation study. PCNA expression identifies more cells than just those in S-Phase (Yamada *et al.*, 1992). Since there were no difference between control and *P. amarus* administered groups in the PCNA study, the necessity to carryout the ^3H -Thymidine incorporation study did not occur.

Experimentally induced T-cell mediated hepatic injury is assessed by apoptosis (Tiegs *et al.*, 1992). In our study, the assessment of apoptosis by using Apop Tag Kit showed similar pattern between control and *P. amarus* administered groups.

Therefore, the biochemical and histological studies showed no measurable toxic side effect of *P. amarus* extract on rat liver.

CONCLUSIONS

Acute oral administration of *P. amarus* extract is non-toxic to the rat liver, even at a dose of 5 gram /kg body weight. The chronic toxicity studies of *P. amarus* extracts administration (of 100-800 mg/kg body wt) showed the absence of cumulative toxicity as reflected by the non-significant change in the parameters studied as well as from the results of the histological studies.

So, further studies can be carried out in this plant to assess their antihepatotoxic, antidiabetic, anticancerous and antidiarrheal action.

ACKNOWLEDGEMENTS

We would like to thank En. Lukmi Ismail for his great help in plant extract preparation for our study and also wish to acknowledge the help provided by the technical staff from Department of Pharmacology, Department of Pathology and Department of Chemical Pathology. The help provided by Dr.Mummedy swamy during the experimental study is kindly acknowledged. This project was carried out under the financial support of USM short-term grant (304/PPSP/6131166).

REFERENCES

- Arnold, D.L., 1990. Oral Ingestion studies. In: Handbook of In vivo toxicity testing, Academic Press Inc., pp.167-188.
- Brock, W.D., Trochimowicz, H.J., Millischer, R.JH., Farr, C., Kawano, T., Rusch, G.M., 1995. Acute and subchronic toxicity of 1,1-dichloro-1-fluoroethane (HCFC-141 b). Food and Chemical Toxicology 33, 483-490.
- Calixto, J.B., 2000. Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). Brazilian Journal of Medical and Biological Research 33, 179-189.
- Calixto, J.B., Santos, A.R.S, Cechinel-Filho, V., Yunes, R.A., 1998. A review of the plants of the genus *phyllanthus*: their chemistry, pharmacology and therapeutic potential. Medicinal Research Reviews 18, 225-258.
- Doshi, J.C., Vaidya, A.B., Antarkar, D.S., Deolalikar, R., Antani, D.H., 1994. A two stage clinical trial of *phyllanthus amarus* in hepatitis B carriers: failure to eradicate the surface antigen. Indian Journal of Gastroenterology 13, 7-8.
- Friedman, L.S., Martin, P., Munoz, S.J., 1996. Liver function tests and the objective evaluation of the patient with liver disease. In : D. Zakin and T.D. Boyer (Eds.), Hepatology: a text book of liver disease. 3rd edition, WB saunders, Philadelphia, pp 791-833.
- Kaplowitz, N., 1997. Hepatotoxicity of herbal remedies. Insight into the intricacies of Plant-animal warfare and cell death. Gastroenterology, 113, 1408-1412.
- Kumar, G., Sharmila banu, G., Vanitha pappa, P., Sundararajan, M., Rajasekara Pandian, M., 2004. Hepatoprotective activity of *Trianthera portulacastrum* L. against paracetamol and thioacetamide intoxication in albino rats. Journal of Ethnopharmacology 92, 37-40.
- Leelarasamee, A., Trakulsomboon, S., Maunwongyathi, P., Somanabandhu, A., Pidetcha, P., Matrakool, b., Lebnak, T., Ridthimat, W., Chandanayingyong, D., 1990. Failure of *Phyllanthus amarus* to eradicate hepatitis B surface antigen from symptomless carriers. Lancet 335, 1600-1601.
- Moss, D.W., Ralph Handerson, A., 1999. Clinical Enzymology. In: Burtis, C.A., Ashword, E.R., (Eds.), Tietz Text book of clinical chemistry, WB Saunders Company, 3rd edition, Philadelphia, pp 651-683.
- Muhamad bin Zakaria and Mustafa Ali Mohd., 1994. Penerbit Fajar Bakti Sdn. Bhd, Kualalumpur, Malaysia.
- Odetola, A.A., Akojenu, S.M., 2000. Anti-diarrhoeal and gastro-intestinal potentials of the aqueous extract of *Phyllanthus amarus* (Euphorbiaceae). African Journal of Medical Sciences 29, 119-122.

OECD, 1981. Guidelines for the Testing of Chemicals. Organization for Economic Cooperation and Development, Paris.

Ott, M., Thyagarajan, S.P., Gupta, S., 1997. *Phyllanthus amarus* suppresses hepatitis B virus by interrupting interactions between HBV-enhancer I and cellular transcription factors. *European Journal of Clinical Investigation* 27, 908-915.

Rajeshkumar, N.V., Joy, K.L., Kuttan, G., Ramsewak, R.S., Nair, M.G., Kuttan, R., 2002. Antitumor and anticarcinogenic activity of *Phyllanthus amarus* extract. *Journal of Ethnopharmacology* 81, 17-22.

Santos, A.R.S., Ailho, V.C., Yunes, R.A., Calixto, J.B., 1995. Analysis of the mechanism underlying the antinociceptive effect of the extracts of plants from the genus *phyllanthus*. *General Pharmacology* 26, 1499-1506.

Shaw, D., Leon, C., Koleu, S., Murray, V., 1997. Traditional remedies and Food Supplements. A five-year toxicological study (1991-1995), *Drug safety*, 17, 342-356.

Srividya, N., Periwal, S., 1995. Diuretic, hypotensive and hypoglycaemic effect of *Phyllanthus amarus*. *Indian Journal of Experimental Biology* 33, 861-864.

Sturgill, M.G., Lambert, G.H., 1997. Xenobiotics-induced hepatotoxicity; Mechanism of liver injury and method of monitoring hepatic function. *Clinical Chemistry* 43, 1512-1526.

Thyagarajan, S.P., Jayaram, S., Gopalakrishnan, V., Hari, R., Jeyakumar, P., Sripathi, M.S., 2002. Herbal medicines for liver diseases in India. *Journal of Gastroenterology and Hepatology* 17, S370-S376.

Thyagarajan, S.P., Subramanian, S., Thirunalasundari, T., Venkateswaran, P.S., Blumberg, B.S., 1988. Effect of *Phyllanthus amarus* on chronic carriers of hepatitis B virus. *Lancet* 2, 764-766.

Tiegs, G., Hentschel, J., Wendel, A., 1992. A T cell-dependent experimental liver injury in mice inducible by concanavalin A. *American Journal of Clinical Investigation* 90, 196.

Wang, B.E., 2000. Treatment of chronic liver diseases with traditional Chinese medicine. *Journal of Gastroenterology and Hepatology*, 15, E67-E70.

Wang, M., Zhou, H.B., Zhao, G.I., Zhao, S., Mai, K., 1991. *Phyllanthus amarus* cannot eliminate HbsAg in chronic hepatitis B virus infection. *Hepatology* 21, 22-24.

Wang, T., Fontenot, R.D., Soni, G.M., Buccin, T.J., Mehendale, H.M., 2000. Enhanced hepatotoxicity and toxic outcome of thioacetamide in streptozotocin-induced diabetic rats. *Toxicology and Applied Pharmacology* 166, 92-100.

WHO, 1993. Research Guidelines for evaluating the safety and efficacy of herbal medicine. Regional office for the western pacific, Manila.

Yamada, K., Youshitake, K., Sato M., Ahnen, D.J., 1992. Proliferating Cell Nuclear Antigen Expression in Normal, Preneoplastic and Neoplastic Colonic Epithelium of the Rat. *Gastroenterology* 103, 160-167.

Manuscript to be submitted to Journal of Ethnopharmacology